



Heriot-Watt University
Research Gateway

Insights on Quorum-Quenching Properties of *Lysinibacillus fusiformis* Strain RB21, a Malaysian Municipal Solid-Waste Landfill Soil Isolate, via Complete Genome Sequence Analysis

Citation for published version:

Yong, D, Ee, R, Lim, Y-L, Chang, C-Y, Yin, W-F & Chan, K-G 2015, 'Insights on Quorum-Quenching Properties of *Lysinibacillus fusiformis* Strain RB21, a Malaysian Municipal Solid-Waste Landfill Soil Isolate, via Complete Genome Sequence Analysis', *Genome Announcements*, vol. 3, no. 3, e00409-15.
<https://doi.org/10.1128/genomeA.00409-15>

Digital Object Identifier (DOI):

[10.1128/genomeA.00409-15](https://doi.org/10.1128/genomeA.00409-15)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Genome Announcements

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Insights on Quorum-Quenching Properties of *Lysinibacillus fusiformis* Strain RB21, a Malaysian Municipal Solid-Waste Landfill Soil Isolate, via Complete Genome Sequence Analysis

Delicia Yong,^a Robson Ee,^a Yan-Lue Lim,^a Chien-Yi Chang,^{b,c} Wai-Fong Yin,^a  Kok-Gan Chan^a

Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia^a; Interdisciplinary Computing and Complex BioSystems (ICOS) Research Group, School of Computing Science, Newcastle University, Claremont Tower, Newcastle upon Tyne, United Kingdom^b; The Centre for Bacterial Cell Biology, Medical School, Newcastle University, Newcastle upon Tyne, United Kingdom^c

***Lysinibacillus fusiformis* strain RB21 is a quorum-quenching bacterium that is able to degrade quorum-sensing signaling molecules. Here, we present the first complete genome sequence of *L. fusiformis* strain RB21. The finished genome is 4.8 Mbp in size, and the quorum-quenching gene was identified.**

Received 21 March 2015 Accepted 25 March 2015 Published 7 May 2015

Citation Yong D, Ee R, Lim Y-L, Chang C-Y, Yin W-F, Chan K-G. 2015. Insights on quorum-quenching properties of *Lysinibacillus fusiformis* strain RB21, a Malaysian municipal solid-waste landfill soil isolate, via complete genome sequence analysis. *Genome Announc* 3(3):e00409-15. doi:10.1128/genomeA.00409-15.

Copyright © 2015 Yong et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Kok-Gan Chan, kokgan@um.edu.my.

Lysinibacillus fusiformis strain RB21 was isolated from a nonoperational municipal solid waste landfill site in Malaysia. Equipped with quorum-quenching (QQ) ability, *L. fusiformis* strain RB21 is able to degrade quorum-sensing (QS) signaling molecules, namely, *N*-acyl-homoserine lactone (AHL). As QS regulates the expression of various genes that are vital for bacterial pathogenicity (1), QQ is garnering attention as a potential alternative for the treatment of bacterial infections (2). In this study, we sequenced and here report the first complete genome sequence of *L. fusiformis*, with the objectives of studying this bacterium comprehensively and characterizing its QQ gene.

The MasterPure DNA purification kit (Epicentre) was used to extract the genomic DNA according to the manufacturer's instructions. The Qubit 2.0 fluorometer (Invitrogen, USA) and NanoDrop spectrophotometer (Thermo Fisher Scientific, USA) were used to determine the concentration and purity of the extracted genomic DNA, respectively. A 10-kb SMRTbell template library was then constructed and sequenced using the PacBio RSII single-molecule real-time (SMRT) sequencing platform (Pacific Biosciences, USA) (3, 4). From the sequencing output, a total of 124,500 reads with a mean read length of 4,993 bp were generated. The filtered insert reads were subsequently assembled into two scaffolds using the Hierarchical Genome Assembly Process (HGAP) *de novo* assembly workflow (version 2) (Pacific Biosciences) (5).

Gene prediction and genome annotation were performed using RAST (6), Prokka (7), Integrated Microbial Genomes-Expert Review (IMG-ER) (8), RNAmmer (9), tRNAscan-SE (10), and the NCBI Prokaryotic Genome Annotation Pipeline (PGAP).

The genome of *L. fusiformis* strain RB21 comprises a circular chromosomal genome (4,843,789 bp, 37.62% G+C content, and 104.3-fold coverage) and a circular plasmid (17,280 bp, 53.67% G+C content, and 673.45-fold coverage). Annotation analysis revealed the presence of 4,900 genes, of which 4,721 genes were protein-coding genes, whereas another 179 genes were RNA genes

(42 rRNA and 107 tRNA genes). In addition, proteins encoded by 3,018 genes were assigned to 21 Clusters of Orthologous Groups (COGs). A QQ gene, *N*-acyl-homoserine lactonase, was also found in the genome. The complete genome sequence of *L. fusiformis* strain RB21 provides the first genetic evidence of its QQ activity.

Nucleotide sequence accession numbers. This complete genome sequence has been deposited in DDBJ/ENA/GenBank under the accession numbers CP010820 (chromosome) and CP010821 (plasmid). The versions described in this paper are the first versions, CP010820.1 and CP010821.1.

ACKNOWLEDGMENTS

This work was supported by the University of Malaya for High Impact Research (UM-MOHE HIR) grant UM C/625/1/HIR/MOHE/CHAN/01, no. A000001-50001 and grant UM C/625/1/HIR/MOHE/CHAN/14/1, H-50001-A000027, awarded to Kok-Gan Chan, which is gratefully acknowledged.

REFERENCES

- Williams P, Winzer K, Chan WC, Cámara M. 2007. Look who's talking: communication and quorum sensing in the bacterial world. *Philos Trans R Soc Lond B Biol Sci* 362:1119–1134. <http://dx.doi.org/10.1098/rstb.2007.2039>.
- Hong KW, Koh CL, Sam CK, Yin WF, Chan KG. 2012. Quorum quenching revisited—from signal decays to signalling confusion. *Sensors* 12:4661–4696. <http://dx.doi.org/10.3390/s120404661>.
- Ee R, Lim YL, Yin WF, Chan KG. 2014. *De novo* assembly of the quorum-sensing *Pandoraea* sp. strain RB-44 complete genome sequence using PacBio single-molecule real-time sequencing technology. *Genome Announc* 2(2):e00245-14. <http://dx.doi.org/10.1128/genomeA.00245-14>.
- Chan KG, Yin WF, Lim YL. 2014. Complete genome sequence of *Pseudomonas aeruginosa* strain YL84, a quorum-sensing strain isolated from compost. *Genome Announc* 2(2):e00246-14. <http://dx.doi.org/10.1128/genomeA.00246-14>.
- Chan KG, Tan KH, Yin WF, Tan JY. 2014. Complete genome sequence of *Cedecea neteri* strain ssm04, a bacterium isolated from pickled mackerel sashimi. *Genome Announc* 2(6):e01339-14. <http://dx.doi.org/10.1128/genomeA.01339-14>.

6. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
7. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
8. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. 2009. IMG-ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 25:2271–2278. <http://dx.doi.org/10.1093/bioinformatics/btp393>.
9. Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
10. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.