



Heriot-Watt University  
Research Gateway

## Generalised cosine functions, basis and regularity properties

### Citation for published version:

Boulton, L & Melkonian, H 2016, 'Generalised cosine functions, basis and regularity properties', *Journal of Mathematical Analysis and Applications*, vol. 444, no. 1, pp. 25–46.  
<https://doi.org/10.1016/j.jmaa.2016.06.024>

### Digital Object Identifier (DOI):

[10.1016/j.jmaa.2016.06.024](https://doi.org/10.1016/j.jmaa.2016.06.024)

### Link:

[Link to publication record in Heriot-Watt Research Portal](#)

### Document Version:

Peer reviewed version

### Published In:

Journal of Mathematical Analysis and Applications

### General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [open.access@hw.ac.uk](mailto:open.access@hw.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Accepted Manuscript

Generalised cosine functions, basis and regularity properties

Lyonell Boulton, Houry Melkonian


PII: S0022-247X(16)30254-2  
DOI: <http://dx.doi.org/10.1016/j.jmaa.2016.06.024>  
Reference: YJMAA 20512

To appear in: *Journal of Mathematical Analysis and Applications*

Received date: 13 November 2015

Please cite this article in press as: L. Boulton, H. Melkonian, Generalised cosine functions, basis and regularity properties, *J. Math. Anal. Appl.* (2016), <http://dx.doi.org/10.1016/j.jmaa.2016.06.024>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



# Generalised cosine functions, basis and regularity properties

Lyonell Boulton<sup>1</sup> & Houry Melkonian<sup>2</sup>

*Department of Mathematics and  
Maxwell Institute for Mathematical Sciences  
Heriot-Watt University, Edinburgh EH14 4AS, UK.*

June 2016

## Abstract

We examine regularity and basis properties of the family of rescaled  $p$ -cosine functions. We find sharp estimates for their Fourier coefficients. We then determine two thresholds,  $p_0 < 2$  and  $p_1 > 2$ , such that this family is a Schauder basis of  $L_s(0, 1)$  for all  $s > 1$  and  $p \in [p_0, p_1]$ .

## 1 Introduction

The contents of this paper can be summarised as follows. Consider a continuous 2-periodic function  $f : \mathbb{R} \rightarrow \mathbb{C}$ . Denote by  $\mathcal{F}$  the family of rescalings  $\mathcal{F} = \{f(nx)\}_{n \in \mathbb{N}}$ . When does  $\mathcal{F}$  form a Schauder basis of  $L_s \equiv L_s(0, 1)$  for all  $s > 1$ ? This question can be traced back to a 1945 note by Arne Beurling [1]. However, quite remarkably, there are still a number of open problems associated to it. As it turns, finding a concrete answer can be extremely difficult, even for apparently simple functions  $f$ .

In a series of recent papers the above question has been addressed for the particular choice  $f(x) = \sin_p(\pi_p x)$ , the  $p$ -sine functions. Let  $p > 1$ . Let the increasing function  $F_p : [0, 1] \rightarrow [0, \frac{\pi_p}{2}]$  be defined by means of the integral

$$(1) \quad F_p(y) := \int_0^y (1 - t^p)^{-\frac{1}{p}} dt$$

where

$$\pi_p := 2F_p(1) = \frac{2\pi}{p \sin(\frac{\pi}{p})}.$$

<sup>1</sup>Email address: L.Boulton@hw.ac.uk

<sup>2</sup>Email address: hm189@hw.ac.uk

Denote the inverse of  $F_p$  by  $\sin_p$ , which is increasing in the segment  $[0, \frac{\pi_p}{2}]$ . Extend to the whole of  $\mathbb{R}$  by means of the rules

$$(2) \quad \sin_p(-x) = -\sin_p(x) \quad \text{and} \quad \sin_p\left(\frac{\pi_p}{2} - x\right) = \sin_p\left(\frac{\pi_p}{2} + x\right),$$

which makes it  $2\pi_p$ -periodic and continuous in  $\mathbb{R}$ . The choice  $p = 2$  corresponds to the standard trigonometric setting  $\sin_2 \equiv \sin$ ,  $\pi_2 = \pi$  and in this case  $\mathcal{F}$  is a Schauder basis of  $L_s$  for all  $s > 1$  as a consequence of Fourier's Theorem.

The study of generalised trigonometric functions has a long history which dates back to the XIX century, [14] and [9, Note 4.1]. The study of the  $p$ -sine functions is closely related to the one-dimensional  $p$ -Laplacian non-linear eigenvalue problem, see the work of Elbert [10] and Ôtani [15]. Their basis properties were first examined in [2], where it was announced that the family  $\{\sin_p(n\pi_p \cdot)\}_{n \in \mathbb{N}}$  forms a Schauder basis of  $L_s$  for all  $s > 1$  and  $p \geq \frac{12}{11}$ . Further development in this respect were settled in [5], [6] and [4]. Currently we know that this family is a Schauder basis of  $L_s$  for all  $s > 1$  when  $p > \tilde{p}_0$ , and also a Riesz basis of  $L_2$  for  $p \in (\hat{p}_0, \tilde{p}_0]$ , where  $\tilde{p}_0 \approx 1.087$  and  $\hat{p}_0 \approx 1.044$  satisfy complicated identities involving hypergeometric functions [4].

Let

$$(3) \quad \cos_p x := \frac{d}{dx} \sin_p x \quad \forall x \in \mathbb{R}$$

and set  $f(x) = \cos_p(\pi_p x)$ , the  $p$ -cosine functions. From the various results established in the recent paper [7], it follows that  $\mathcal{F} \cup \{1\}$  is a Schauder basis of  $L_s$  for all  $s > 1$  and  $p \in (p_0^\dagger, 2]$  where  $p_0^\dagger \approx 1.75$ . In the present work we establish that this basis property in fact holds true for  $p$  in a wider segment. To be precise, we show the following.

**Theorem 1.** *There exist  $p_0 < \frac{3}{2}$  and  $p_1 > \frac{11}{5}$ , such that  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^\infty$  is a Schauder basis of  $L_s$  for all  $s > 1$  and  $p \in [p_0, p_1]$ .*

The constants  $p_0$  and  $p_1$  will be given analytically as the zeros of corresponding equations involving the parameter  $p$ . Their approximated values turn out to be  $p_0 \approx 1.46$  and  $p_1 \approx 2.43$ .

The proof of Theorem 1 is naturally divided into the cases  $1 < p < 2$  and  $p > 2$ . The different parts of the paper follow this division. In Section 2 we collect various properties of the  $p$ -trigonometric functions which will be useful later on. In Section 3 we establish precise upper bounds on the asymptotic behaviour of the Fourier coefficients of  $\cos_p(\pi_p \cdot)$ . In Section 4 we recall the framework for determining invertibility of the change of coordinates map between the families  $\{\cos(n\pi \cdot)\}_{n=0}^\infty$  and  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^\infty$ . In Section 5 we assemble the proof of Theorem 1, by combining the crucial criterion (12) of Section 4 with the estimates of Section 3. In the final Section 6 we describe the relation between the results announced here and other existing work.

## 2 The generalised trigonometric functions

We begin by recalling various elementary properties of the  $p$ -cosine functions. A more complete account on this matter can be found in [5, Section 2] and [9, Chapter 2].

Throughout we shall assume that  $1 < p < \infty$ . Note that  $\pi_p$  is a decreasing function, smooth in  $p > 1$ , such that

$$\begin{cases} \pi_p \rightarrow \infty & p \rightarrow 1^+ \\ \pi_p = \pi & p = 2 \\ \pi_p \rightarrow 2 & p \rightarrow \infty. \end{cases}$$

Here and everywhere below we write  $p' := p/(p-1)$ . According to [5, (2.3)], we know that

$$(4) \quad p' \pi_{p'} = p \pi_p.$$

From (2) and (3) it immediately follows that  $\cos_p$  is  $2\pi_p$ -periodic,

$$\cos_p(x) = \cos_p(-x) \quad \text{and} \quad \cos_p\left(x + \frac{\pi_p}{2}\right) = -\cos_p\left(x - \frac{\pi_p}{2}\right) \quad \forall x \in \mathbb{R}.$$

Moreover, setting  $y = \sin_p(x)$  for  $x \in [0, \pi_p/2]$  in the formula for the derivative of the inverse function of (1), gives

$$(5) \quad \cos_p(x) = (1 - y^p)^{1/p} = (1 - \sin_p(x)^p)^{1/p}.$$

Thus,  $\cos_p$  is decreasing in  $(0, \pi_p/2]$ ,  $\cos_p(0) = 1$  and  $\cos_p(\pi_p/2) = 0$ . In fact we have,

$$|\sin_p x|^p + |\cos_p x|^p = 1 \quad \forall x \in \mathbb{R}.$$

See [5, (2.7)].

**Lemma 1.** For all  $x \in [0, \frac{1}{2})$ ,

a.

$$\cos_p(\pi_p x) = \sin_{p'}\left(\pi_{p'}\left(\frac{1}{2} - x\right)\right)^{p'-1}$$

b.

$$\frac{d}{dx} \cos_p(x) = -\sin_p(x)^{p-1} \cos_p(x)^{2-p}$$

c.

$$\frac{d^2}{dx^2} \cos_p(x) = \sin_p(x)^{p-2} \cos_p(x)^{3-2p} [2 - p - \cos_p(x)^p].$$

*Proof.* The calculations leading to “a” and “b” can be found in the proofs of [5, Proposition 2.2] and [5, Proposition 2.1], respectively. From (5) we get

$$\begin{aligned} \frac{d^2}{dx^2} \cos_p(x) &= (2-p) \sin_p(x)^{2p-2} \cos_p(x)^{3-2p} - (p-1) \sin_p(x)^{p-2} \cos_p(x)^{3-p} \\ &= \sin_p(x)^{p-2} \cos_p(x)^{3-2p} [(2-p) \sin_p(x)^p - (p-1) \cos_p(x)^p], \end{aligned}$$

which is “c”.  $\square$

The following inequalities will be important below.

**Lemma 2.** *Let  $1 < p \leq q < \infty$  and  $x \in [0, \frac{1}{2}]$ . Then*

a.  $\sin_p(\pi_p x) \geq \sin_q(\pi_q x)$

b.  $\cos_p(\pi_p x) \leq \cos_q(\pi_q x)$ .

*Proof.* Statement “a” is [5, Corollary 4.4-(iii)].

Let us show “b”. A direct evaluation at  $x = 0$  and  $x = 1/2$  gives equality for all  $p$  and  $q$  at these points, so these two cases are immediate. Let  $x \in (0, \frac{1}{2})$  be fixed. Since  $p'$  is decreasing in  $p > 1$ , from part “a” it follows that

$$\frac{d}{dp} \sin_{p'} \left( \pi_{p'} \left( \frac{1}{2} - x \right) \right) \geq 0 \quad \forall p \in (1, \infty).$$

Note that,  $0 < \sin_{p'}(\pi_{p'}(\frac{1}{2} - x)) < 1$  and hence  $\ln(\sin_{p'}(\pi_{p'}(\frac{1}{2} - x))) < 0$ . Substituting the identity from Lemma 1(a), yields

$$\begin{aligned} \frac{d}{dp} \cos_p(\pi_p x) &= \frac{d}{dp} \left[ \sin_{p'} \left( \pi_{p'} \left( \frac{1}{2} - x \right) \right) \right]^{\frac{1}{p-1}} \\ &= \left[ -\frac{\ln(\sin_{p'}(\pi_{p'}(\frac{1}{2} - x)))}{(p-1)^2} + \frac{\frac{d}{dp} [\sin_{p'}(\pi_{p'}(\frac{1}{2} - x))]}{(p-1) \sin_{p'}(\pi_{p'}(\frac{1}{2} - x))} \right] \cos_p(\pi_p x) > 0. \end{aligned}$$

This implies “b”.  $\square$

## 2.1 The case $1 < p < 2$

For  $1 < p < 2$ , let  $u_p : [0, \frac{1}{2}] \rightarrow \mathbb{R}$  be given by

$$u_p(x) := \cos'_p(\pi_p x) = -\sin_p(\pi_p x)^{p-1} \cos_p(\pi_p x)^{2-p}.$$

This function will simplify the notation when we determine estimates for the Fourier coefficients of the  $p$ -cosine functions in Section 3.1. Here and everywhere below we write

$$(6) \quad c_p := (p-1)^{\frac{p-1}{p}} (2-p)^{\frac{2-p}{p}}.$$

**Lemma 3.** *Let  $1 < p < 2$ . Then*

- a.  $u_p(x) \leq 0$  for all  $x \in [0, \frac{1}{2}]$
- b.  $u_p(x) = 0$  if and only if  $x = 0$  or  $x = \frac{1}{2}$
- c.  $u_p(x) = -c_p$  for  $x \in [0, \frac{1}{2}]$  if and only if  $x = m_p \in (0, \frac{1}{2})$ , where  $m_p$  is the unique point such that  $\cos_p(\pi_p m_p)^p = 2 - p$
- d.  $u_p : [0, m_p] \rightarrow [-c_p, 0]$  is decreasing
- e.  $u_p : [m_p, \frac{1}{2}] \rightarrow [-c_p, 0]$  is increasing
- f.  $\min_{x \in [0, \frac{1}{2}]} u_p(x) = -c_p$ .

*Proof.* Since  $\sin_p(\pi_p x)$  and  $\cos_p(\pi_p x)$  are non-negative over  $[0, \frac{1}{2}]$ , then “a” holds true. Since  $\sin_p(\pi_p x)$  only vanishes at  $x = 0$  and  $\cos_p(\pi_p x)$  only vanishes at  $x = \frac{1}{2}$  in this interval, then “b” holds true.

Lemma 1-c gives

$$u'_p(x) = \pi_p \sin_p(\pi_p x)^{p-2} \cos_p(\pi_p x)^{3-2p} [2 - p - \cos_p(\pi_p x)^p].$$

Neither  $\sin_p$  nor  $\cos_p$  vanish in  $(0, \frac{1}{2})$ . On the other hand,  $\cos_p(0) = 1 > 2 - p$ ,  $\cos_p(\frac{x_p}{2}) = 0 < 2 - p$  and  $\cos_p(\pi_p x)^p$  is decreasing for  $x \in (0, \frac{1}{2})$ . Then the term  $\cos_p(\pi_p x)^p + p - 2$  indeed vanishes at the unique point  $m_p \in (0, \frac{1}{2})$  as stated in “c”.

At  $m_p$ ,

$$\begin{aligned} u_p(m_p) &= -\sin_p(\pi_p m_p)^{p-1} \cos_p(\pi_p m_p)^{2-p} \\ &= -(1 - \cos_p(\pi_p m_p)^p)^{\frac{p-1}{p}} \cos_p(\pi_p m_p)^{2-p} = -c_p. \end{aligned}$$

Hence, the proof of “d” and “e”, and thus of “f”, is achieved as follows. Just observe that in the expression for  $u'_p(x)$  above,  $\cos_p(\pi_p x)^p > 2 - p$  for  $x \in [0, m_p]$  and  $\cos_p(\pi_p x)^p < 2 - p$  for  $x \in (m_p, \frac{1}{2})$ , because  $\cos_p(\pi_p x)$  is decreasing in  $x \in (0, \frac{1}{2})$ .  $\square$

According to parts “d” and “e” of Lemma 3, the function  $u_p$  is invertible, when restricted to the segments  $[0, m_p]$  and  $[m_p, \frac{1}{2}]$ . We denote the inverses by  $w_{1,p} : [-c_p, 0] \rightarrow [0, m_p]$  and  $w_{2,p} : [-c_p, 0] \rightarrow [m_p, \frac{1}{2}]$ , respectively, so that

$$u_p(w_{k,p}(x)) = x \quad \forall x \in [-c_p, 0] \quad k = 1, 2.$$

## 2.2 The case $p > 2$

For  $p > 2$ , let  $v_p : (0, \frac{1}{2}] \rightarrow [0, \infty)$  be given by

$$v_p(x) := (p' - 1) \sin_{p'}(\pi_{p'}x)^{p'-2} \cos_{p'}(\pi_{p'}x).$$

Let us summarise various properties of this function, which will be employed in Section 3.2.

**Lemma 4.** *Let  $p > 2$ . Then*

- a.  $v_p$  is decreasing in  $(0, \frac{1}{2}]$
- b.  $\lim_{x \rightarrow 0^+} x v_p(x) = 0$
- c.  $\lim_{x \rightarrow 0^+} v_p(x) = +\infty$  and  $v_p(\frac{1}{2}) = 0$
- d.  $\lim_{x \rightarrow 0^+} v_p'(x) = -\infty$  and  $v_p'(\frac{1}{2}) = 0$ .

*Proof.* For  $p > 2$ ,  $p' \in (1, 2)$  and so  $p' - 2 < 0$ . Since,  $\sin_{p'}(\pi_{p'}x)$  is increasing and  $\cos_{p'}(\pi_{p'}x)$  is decreasing in  $x \in (0, \frac{1}{2})$ , then “a” holds true.

Let us show “b”. L’Hôpital’s Rule gives

$$\lim_{x \rightarrow 0^+} \frac{x}{[\sin_{p'}(\pi_{p'}x)]^{2-p'}} = \lim_{x \rightarrow 0^+} \frac{[\sin_{p'}(\pi_{p'}x)]^{p'-1}}{(2-p')\pi_{p'} \cos_{p'}(\pi_{p'}x)} = 0.$$

Then,

$$\lim_{x \rightarrow 0^+} x v_p(x) = \lim_{x \rightarrow 0^+} (p' - 1) \frac{x \cos_{p'}(\pi_{p'}x)}{[\sin_{p'}(\pi_{p'}x)]^{2-p'}} = 0,$$

as claimed in “b”.

Both statements “c” and “d” follow directly from (5), the expression

$$v_p'(x) = (p' - 1)\pi_{p'} \sin_{p'}(\pi_{p'}x)^{p'-3} \cos_{p'}(\pi_{p'}x)^{2-p'} \left[ (p' - 1) \cos_{p'}(\pi_{p'}x)^{p'} - 1 \right],$$

and continuity of  $\sin_p$  and  $\cos_p$  at  $x = 0$ .  $\square$

According to this lemma, there exists a function  $z_p : [0, \infty) \rightarrow (0, \frac{1}{2}]$  such that  $z_p$  is the inverse function of  $v_p$ . This inverse function has the following characteristics.

- a.  $z_p$  is decreasing in  $[0, \infty)$
- b.  $z_p(0) = \frac{1}{2}$  and  $\lim_{x \rightarrow \infty} z_p(x) = 0$
- c.  $\lim_{x \rightarrow 0^+} z_p'(x) = +\infty$  and  $\lim_{x \rightarrow \infty} z_p'(x) = 0$ .



### 3 The Fourier coefficients of the $p$ -cosine functions

Let

$$a_j(p) \equiv a_j := 2 \int_0^1 \sin_p(\pi_p x) \sin(j\pi x) dx \quad \forall j \in \mathbb{N}$$

be the Fourier sine coefficients of  $\sin_p(\pi_p x)$ . Let

$$b_j(p) \equiv b_j := 2 \int_0^1 \cos_p(\pi_p x) \cos(j\pi x) dx \quad \forall j \in \mathbb{N} \cup \{0\}$$

be the Fourier cosine coefficients of  $\cos_p(\pi_p x)$ . Since  $\sin_p$  is an odd function and  $\cos_p$  is an even function,  $a_j = b_j = 0$  for all  $j \equiv_2 0$ . Here and elsewhere below we will write  $j \equiv_2 k$  to denote that  $j \equiv k \pmod{2}$ .

**Lemma 5.** For  $j \in \mathbb{N}$ ,

$$b_j(p) = \frac{j\pi}{\pi_p} a_j(p).$$

*Proof.* Let  $j \equiv_2 1$ . Integration by parts alongside with the fact that  $\cos_p(\pi_p x)$  and  $\cos(j\pi x)$  are odd with respect to  $\frac{1}{2}$ , yield

$$\begin{aligned} b_j &= 2 \int_0^1 \cos_p(\pi_p x) \cos(j\pi x) dx = 4 \int_0^{\frac{1}{2}} \cos_p(\pi_p x) \cos(j\pi x) dx \\ &= \frac{4}{\pi_p} \cos(j\pi x) \sin_p(\pi_p x) \Big|_0^{\frac{1}{2}} + \frac{4j\pi}{\pi_p} \int_0^{\frac{1}{2}} \sin_p(\pi_p x) \sin(j\pi x) dx \\ &= \frac{j\pi}{\pi_p} a_j. \end{aligned}$$

□

We now find estimates on  $|b_j(p)|$  in terms of the parameter  $p > 1$ .

#### 3.1 The case $1 < p < 2$

**Lemma 6.** For  $1 < p < 2$ , let  $c_p > 0$  be given by (6). Then

$$|b_j(p)| < \frac{8\pi_p}{j^2\pi^2} c_p \quad \forall j \geq 1.$$

*Proof.* Integrate by parts twice to get

$$\begin{aligned}
b_j &= 4 \int_0^{\frac{1}{2}} \cos_p(\pi_p x) \cos(j\pi x) dx \\
&= \frac{4}{j\pi} \cos_p(\pi_p x) \sin(j\pi x) \Big|_0^{\frac{1}{2}} - \frac{4\pi_p}{j\pi} \int_0^{\frac{1}{2}} \cos'_p(\pi_p x) \sin(j\pi x) dx \\
&= -\frac{4\pi_p}{j\pi} \int_0^{\frac{1}{2}} \cos'_p(\pi_p x) \sin(j\pi x) dx \\
&= \frac{4\pi_p}{j^2\pi^2} \cos'_p(\pi_p x) \cos(j\pi x) \Big|_0^{\frac{1}{2}} - \frac{4\pi_p}{j^2\pi^2} \int_0^{\frac{1}{2}} \frac{d}{dx} [\cos'_p(\pi_p x)] \cos(j\pi x) dx.
\end{aligned}$$

From the identities in Lemma 3(b), it follows that the boundary term in the fourth equality always vanishes. Thus,

$$\begin{aligned}
b_j &= -\frac{4\pi_p}{j^2\pi^2} \int_0^{\frac{1}{2}} u'_p(x) \cos(j\pi x) dx \\
&= -\frac{4\pi_p}{j^2\pi^2} \left( \int_0^{m_p} u'_p(x) \cos(j\pi x) dx + \int_{m_p}^{\frac{1}{2}} u'_p(x) \cos(j\pi x) dx \right) \\
&= -\frac{4\pi_p}{j^2\pi^2} \left( \int_0^{-c_p} \cos(j\pi w_{1,p}(s)) ds + \int_{-c_p}^0 \cos(j\pi w_{2,p}(s)) ds \right).
\end{aligned}$$

Hence,

$$\begin{aligned}
|b_j| &\leq \frac{4\pi_p}{j^2\pi^2} \left[ \int_{-c_p}^0 |\cos(j\pi w_{1,p}(s))| ds + \int_{-c_p}^0 |\cos(j\pi w_{2,p}(s))| ds \right] \\
&< \frac{8\pi_p}{j^2\pi^2} c_p,
\end{aligned}$$

because the functions inside the integrals are not constants identically equal to 1.  $\square$

### 3.2 The case $p > 2$

Let  $p > 2$ . According to Lemma 1(a),

$$b_j(p) = 4 \int_0^{\frac{1}{2}} \sin_{p'} \left( \pi_{p'} \left( \frac{1}{2} - x \right) \right)^{\frac{1}{p-1}} \cos(j\pi x) dx.$$

Since  $\cos(j\pi(\frac{1}{2} - t)) = (-1)^{\frac{j-1}{2}} \sin(j\pi t)$  for  $j \equiv_2 1$ , changing variables to  $t = \frac{1}{2} - x$  gives

$$b_j = (-1)^{\frac{j-1}{2}} 4 \int_0^{\frac{1}{2}} \sin_{p'}(\pi_{p'} t)^{\frac{1}{p-1}} \sin(j\pi t) dt.$$

By virtue of Lemma 4 and integration by parts twice, then

$$\begin{aligned}
b_j &= (-1)^{\frac{j-1}{2}} \frac{4\pi p'}{j\pi} \int_0^{\frac{1}{2}} v_p(t) \cos(j\pi t) dt \\
&= (-1)^{\frac{j-1}{2}} \frac{4\pi p'}{j\pi} \left[ \frac{1}{j\pi} v_p(t) \sin(j\pi t) \Big|_0^{\frac{1}{2}} - \frac{1}{j\pi} \int_0^{\frac{1}{2}} v_p'(t) \sin(j\pi t) dt \right] \\
&= (-1)^{\frac{j+1}{2}} \frac{4\pi p'}{j^2 \pi^2} \int_0^{\frac{1}{2}} v_p'(t) \sin(j\pi t) dt \\
(7) \quad &= (-1)^{\frac{j+3}{2}} \frac{4\pi p'}{j^2 \pi^2} \int_0^\infty \sin(j\pi z_p(y)) dy.
\end{aligned}$$

**Lemma 7.** *Let  $p > 2$ . Then*

$$|b_j(p)| < \frac{2\pi p'}{\pi^2(p-1)} \left[ 2 + \frac{\pi^2}{2}(p-2) \right] j^{-p'} \quad \forall j \geq 3.$$

*Proof.* Since  $p > 2$ , then  $1 < p' < 2$ . Let  $r = p' - 1$ . In view of Lemma 2, we have

$$v_p(t) \leq r [\sin_{p'}(\pi p' t)]^{r-1} \leq r [\sin(\pi t)]^{r-1}$$

and so

$$(8) \quad z_p(y) \leq \frac{1}{\pi} \arcsin \left[ \left( \frac{y}{r} \right)^{\frac{1}{r-1}} \right] =: r_p(y) \quad \forall y \in [r, \infty).$$

Set

$$\eta(j) := r \sin \left( \frac{\pi}{2j} \right)^{r-1}.$$

Then,

$$r_p(\eta(j)) = \frac{1}{2j} < \frac{1}{2}.$$

Here we use the requirement  $j \geq 3$ , in order to make sure that the arc-sine does not change branches.

Set

$$J_1 = \int_0^{\eta(j)} dx = \eta(j)$$

and

$$J_2 = \int_{\eta(j)}^\infty \sin(j\pi r_p(y)) dy.$$

Then, (7) yields

$$|b_j| \leq \frac{4\pi p'}{j^2 \pi^2} (J_1 + J_2).$$

Here  $J_2$  is guaranteed to be on the right hand side, because

$$0 < j\pi z_p(y) \leq j\pi z_p(\eta(j)) \leq j\pi r_p(\eta(j)) = \frac{\pi}{2},$$

so that  $0 < \sin(j\pi z_p(y)) \leq \sin(j\pi r_p(y))$  for  $y \in [\eta(j), \infty)$ .

Let us estimate an upper bound for  $J_2$ . Changing variables to

$$t = j\pi r_p(y) \iff y = r \sin\left(\frac{t}{j}\right)^{r-1}$$

gives

$$\begin{aligned} J_2 &= \int_0^{\frac{\pi}{2}} \frac{r(1-r)}{j} \sin\left(\frac{t}{j}\right)^{r-2} \cos\left(\frac{t}{j}\right) \sin(t) dt \\ &= r(1-r) \int_0^{\frac{\pi}{2}} \sin\left(\frac{t}{j}\right)^{r-1} \left[ \frac{\frac{t}{j}}{\sin\left(\frac{t}{j}\right)} \right] \left( \frac{\sin t}{t} \right) \cos\left(\frac{t}{j}\right) dt. \end{aligned}$$

Note that,

$$(9) \quad \max_{0 < \theta \leq \frac{\pi}{2}} \frac{\theta}{\sin \theta} = \frac{\pi}{2}, \quad \max_{0 < \theta \leq \frac{\pi}{2}} \frac{\sin \theta}{\theta} = 1$$

and

$$0 < t < j\pi r_p(\eta(j)) = \frac{\pi}{2}.$$

Here we are using once again the fact that  $j \geq 3$ . Then

$$J_2 < \frac{\pi}{2} r(1-r) \int_0^{\frac{\pi}{2}} \sin\left(\frac{t}{j}\right)^{r-1} \cos\left(\frac{t}{j}\right) dt.$$

Changing variables to

$$\tau = \sin\left(\frac{t}{j}\right),$$

yields

$$J_2 < \frac{j\pi}{2} r(1-r) \int_0^{\sin \frac{\pi}{2j}} \tau^{r-1} d\tau = \frac{j\pi}{2} (1-r) \sin\left(\frac{\pi}{2j}\right)^r.$$

Then

$$|b_j| < \frac{2\pi_{p'}}{j^2\pi^2} \left[ 2 + \frac{j\pi(1-r)}{r} \sin\left(\frac{\pi}{2j}\right) \right] \eta(j).$$

According to (9), we get

$$\eta(j) \leq rj^{1-r}$$

and

$$(10) \quad |b_j| < \frac{2\pi_{p'}r}{j^2\pi^2} \left[ 2 + \frac{j\pi(1-r)}{r} \frac{\pi}{2j} \right] j^{1-r}.$$

Simplifying the expression on the right hand side, ensures the validity of the lemma.  $\square$

## 4 The change of coordinates map

We now derive various properties of the change of coordinates maps that take the 2-cosine functions into the  $p$ -cosine functions. Most of the material in this section can also be found in [2], [5], [7] and [4]. We keep a self-contained presentation here by including details of the main arguments.

Given any  $g \in L_s$ , denote the even extension of  $g$  with respect to 1 by

$$\tilde{g}(x) = \begin{cases} g(x) & x \in [0, 1] \\ g(2-x) & x \in (1, 2]. \end{cases}$$

A 2-periodic extension of  $g$  to the whole of  $\mathbb{R}$  is then written as

$$g^*(x) = \tilde{g}\left(x - 2 \left\lfloor \frac{x}{2} \right\rfloor\right).$$

The floor function  $[y] \in \mathbb{Z}$  is the unique integer such that  $y - [y] \in [0, 1)$ . For any  $n \in \mathbb{N}$ , let

$$M_n g(x) := g^*(nx).$$

**Lemma 8.** *The operators  $M_n : L_s \rightarrow L_s$  are linear isometries.*

*Proof.* Indeed,

$$\begin{aligned} \|M_n g\|_{L_s}^s &= \int_0^1 |M_n g(x)|^s dx = \int_0^1 |g^*(nx)|^s dx = \int_0^1 |\tilde{g}(nx - 2 \lfloor \frac{nx}{2} \rfloor)|^s dx \\ &= \frac{1}{n} \int_0^n |\tilde{g}(y - 2 \lfloor \frac{y}{2} \rfloor)|^s dy = \frac{1}{n} \sum_{l=0}^{n-1} \int_l^{l+1} |\tilde{g}(y - 2 \lfloor \frac{y}{2} \rfloor)|^s dy \\ &= \frac{1}{n} \left[ \sum_{\substack{l=0 \\ l \equiv_2 0}}^{n-1} \int_l^{l+1} |\tilde{g}(y - 2 \lfloor \frac{y}{2} \rfloor)|^s dy + \sum_{\substack{l=1 \\ l \equiv_2 1}}^{n-1} \int_l^{l+1} |\tilde{g}(y - 2 \lfloor \frac{y}{2} \rfloor)|^s dy \right]. \end{aligned}$$

Changing variables to  $w = y - l$  for  $l \equiv_2 0$  and  $z = y - (l - 1)$  for  $l \equiv_2 1$ , gives

$$\lfloor \frac{y}{2} \rfloor = \begin{cases} \frac{l}{2} & \text{whenever } l \equiv_2 0 \\ \frac{l-1}{2} & \text{whenever } l \equiv_2 1. \end{cases}$$

Hence,

$$\|M_n g\|_{L_s}^s = \frac{1}{n} \left[ \sum_{\substack{l=0 \\ l \equiv_2 0}}^{n-1} \int_0^1 |g(w)|^s dw + \sum_{\substack{l=1 \\ l \equiv_2 1}}^{n-1} \int_1^2 |\tilde{g}(z)|^s dz \right].$$

Another change of variables  $z = 2 - w$ , then yields

$$\|M_n g\|_{L_s}^s = \frac{1}{n} \left[ n \int_0^1 |g(w)|^s dw \right] = \|g\|_{L_s}^s$$

as claimed.  $\square$

Let  $e_n(x) := \cos(n\pi x)$ . If

$$g = \frac{\widehat{g}(0)}{2}e_0 + \sum_{j=1}^{\infty} \widehat{g}(j)e_j \in L_s$$

where

$$\widehat{g}(k) := 2 \int_0^1 g(x)e_k(x)dx \quad \forall k \in \mathbb{N} \cup \{0\}$$

are the corresponding cosine Fourier coefficients, then

$$M_n g = \frac{\widehat{g}(0)}{2}e_0 + \sum_{j=1}^{\infty} \widehat{g}(j)M_n e_j = \frac{\widehat{g}(0)}{2}e_0 + \sum_{j=1}^{\infty} \widehat{g}(j)e_{nj} \in L_s.$$

Now, let  $f_n(x) := \cos_p(n\pi_p x)$ . Note that  $e_0(x) = f_0(x) = 1$  for all  $x \in \mathbb{R}$ . Suitable linear extensions of the map  $A : e_n \mapsto f_n$  are the changes of coordinates between  $\{e_n\}_{n=0}^{\infty}$  and  $\{f_n\}_{n=0}^{\infty}$ . Our next goal is to find a canonical decomposition for  $A$  in terms of  $M_n$  and the Fourier coefficients  $b_n(p)$ . After that, we show that these are bounded operators of the Banach spaces  $L_s$  for all  $s > 1$ .

**Proposition 1.** *For all  $p > 1$ ,*

$$\sum_{j=1}^{\infty} |b_j(p)| < \infty.$$

*Proof.* This is a direct consequence of lemmas 6 and 7. See (14) and (23) below.  $\square$

In the notation of Section 3, we have  $\widehat{f}_1(k) = b_k(p)$  for all  $k \in \mathbb{N} \cup \{0\}$ . Recall that  $b_k = 0$  for  $k \equiv_2 0$ . Since any of the functions  $f_n(x)$  is continuous, then they all have a Fourier cosine expansion

$$f_n(x) = \frac{1}{2}\widehat{f}_n(0)e_0(x) + \sum_{k=1}^{\infty} \widehat{f}_n(k)e_k(x)$$

which is both pointwise convergent for all  $x \in [0, 1]$  and also convergent in the norm of  $L_s$  for all  $s > 1$ . Then, for all  $n > 1$ ,

$$\begin{aligned} \widehat{f}_n(k) &= 2 \int_0^1 f_1(nx) \cos(k\pi x)dx \\ &= 2 \int_0^1 \left( \sum_{m=1}^{\infty} \widehat{f}_1(m) \cos(m\pi nx) \right) \cos(k\pi x)dx \\ &= 2 \sum_{m=1}^{\infty} \widehat{f}_1(m) \int_0^1 \cos(mn\pi x) \cos(k\pi x)dx \\ &= \begin{cases} b_m(p) & \text{for } mn = k, \ m \equiv_2 1 \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Here we can exchange the infinite summation with the integral sign, due to the pointwise convergence of the series, Proposition 1 and the Dominated Convergence theorem.

Let

$$(11) \quad A := \sum_{j=1}^{\infty} b_j(p) M_j.$$

By virtue of Proposition 1, Lemma 8 and the triangle inequality, it follows that the expression (11) is convergent in the operator norm of  $L_s$  and that  $A : L_s \rightarrow L_s$  is a bounded linear operator such that

$$\|A\|_{L_s \rightarrow L_s} \leq \sum_{j=1}^{\infty} |b_j| \|M_j\|_{L_s \rightarrow L_s} = \sum_{j=1}^{\infty} |b_j|.$$

Moreover,

$$Ae_0 = \sum_{j=1}^{\infty} b_j M_j e_0 = \sum_{j=1}^{\infty} b_j e_0 = \sum_{j=1}^{\infty} b_j e_j(0) = \cos_p(\pi_p 0) = 1 = f_0$$

and

$$Ae_n = \sum_{j=1}^{\infty} b_j M_j e_n = \sum_{j=1}^{\infty} \hat{f}_1(j) e_{nj} = \sum_{k=1}^{\infty} \hat{f}_n(k) e_k = f_n \quad \forall n \in \mathbb{N}.$$

These are the change of basis maps between  $\{e_n\}_{n=0}^{\infty}$  and  $\{f_n\}_{n=0}^{\infty}$ .

The operator  $A$  is an homeomorphism of  $L_s$  if and only if the family  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^{\infty}$  is a Schauder basis of  $L_s$ , cf. [12] or [16]. Then we have the following criterion, which is a consequence of [13, Theorem IV-1.16],

$$(12) \quad \sum_{\substack{j=3 \\ j \equiv 21}}^{\infty} |b_j(p)| < |b_1(p)| \quad \Rightarrow \quad \begin{cases} \{\cos_p(n\pi_p \cdot)\}_{n=0}^{\infty} \text{ is a Schauder} \\ \text{basis of } L_s \text{ for all } s > 1. \end{cases}$$

We employ this criterion below in order to determine the basis thresholds for the family  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^{\infty}$  claimed in Theorem 1.

## 5 Proof of Theorem 1

The proof is separated into two cases.

### 5.1 The case $1 < p < 2$

Recall the expression for  $c_p$  given in (6) and consider the identity

$$(13) \quad \pi_p^2 c_p = \frac{\pi^3}{\pi^2 - 8}.$$

**Lemma 9.** *There exists  $1 < p_0 < 2$  such that (13) holds true for  $p = p_0$ . Moreover,*

$$\pi_p^2 c_p < \frac{\pi^3}{\pi^2 - 8} \quad \forall p \in (p_0, 2).$$

*Proof.* It will be enough to prove that  $\pi_p^2 c_p$  is a convex function of the parameter  $p$  for all  $1 < p < 2$ . Indeed, since

$$\lim_{p \rightarrow 1^+} \pi_p^2 c_p = \infty \quad \text{and} \quad \lim_{p \rightarrow 2^-} \pi_p^2 c_p = \pi^2 < \frac{\pi^3}{\pi^2 - 8},$$

both statements will immediately follow from this property.

Firstly note that

$$\frac{d}{dp} \ln(p-1)^{\frac{p-1}{p}} = \frac{1}{p^2} \ln(p-1) + \frac{1}{p}$$

and

$$\frac{d^2}{dp^2} \ln(p-1)^{\frac{p-1}{p}} = \frac{2-p}{p^2(p-1)} - 2 \frac{\ln(p-1)}{p^3} > 0.$$

Then  $\ln(p-1)^{\frac{p-1}{p}}$  is convex for  $1 < p < 2$ .

Similarly, we have

$$\frac{d}{dp} \ln(2-p)^{\frac{2-p}{p}} = \frac{-2}{p^2} \ln(2-p) - \frac{1}{p}$$

and

$$\frac{d^2}{dp^2} \ln(2-p)^{\frac{2-p}{p}} = \frac{4-p}{p^2(2-p)} + 4 \frac{\ln(2-p)}{p^3} > 0.$$

Then, also  $\ln(2-p)^{\frac{2-p}{p}}$  is convex for  $1 < p < 2$ .

Furthermore,

$$\frac{d}{dp} [\ln \pi_p] = \frac{\pi \cot\left(\frac{\pi}{p}\right)}{p^2} - \frac{1}{p}$$

and

$$\frac{d^2}{dp^2} \ln \pi_p = \frac{(p^2 + \pi^2)}{p^4} - \frac{2\pi}{p^3} \cot\left(\frac{\pi}{p}\right) + \frac{\pi^2}{p^4} \cot^2\left(\frac{\pi}{p}\right) > 0.$$

The latter is a consequence of the fact that  $\cos \frac{\pi}{p} < 0$  and  $\sin \frac{\pi}{p} > 0$ . Hence, also  $\ln \pi_p^2$  is convex for  $1 < p < 2$ .

The convexity of the logarithm of each one of the multiplying terms in the expression for  $\pi_p^2 c_p$ , implies that  $\ln \pi_p^2 c_p$  is convex for  $1 < p < 2$ . This ensures that indeed  $\pi_p^2 c_p$  is convex in the same segment and the validity of the statement is ensured.  $\square$



**Corollary 1.** Let  $1 < p_0 < 2$  be such that (13) holds true for  $p = p_0$ . The family  $\{\cos_p(n\pi_p)\}_{n=0}^{\infty}$  is a Schauder basis of  $L_s$  for all  $s > 1$  and  $p_0 \leq p \leq 2$ .

*Proof.* According to Lemma 6,

$$(14) \quad \sum_{\substack{j=3 \\ j \neq 2^1}}^{\infty} |b_j(p)| < \frac{8\pi_p c_p}{\pi^2} \sum_{\substack{j=3 \\ j \neq 2^1}}^{\infty} \frac{1}{j^2} = \frac{\pi_p^2 c_p (\pi^2 - 8)}{\pi^2 \pi_p}.$$

On the other hand, in view of Lemma 5 and Lemma 2(a), we have

$$\begin{aligned} b_1(p) &= \frac{\pi}{\pi_p} a_1 = \frac{4\pi}{\pi_p} \int_0^{\frac{1}{2}} \sin_p(\pi_p x) \sin(\pi x) dx \\ &\geq \frac{4\pi}{\pi_p} \int_0^{\frac{1}{2}} \sin(\pi x)^2 dx = \frac{\pi}{\pi_p}. \end{aligned}$$

Then, Lemma 9 yields

$$\sum_{\substack{j=3 \\ j \neq 2^1}}^{\infty} |b_j(p)| < b_1(p)$$

for all  $p \in [p_0, 2)$ . By virtue of (12) the claimed conclusion follows.  $\square$

Since

$$\pi_{\frac{4}{3}}^2 c_{\frac{4}{3}} = \frac{\pi^2 3^{\frac{5}{4}} \sqrt{2}}{2} > \frac{\pi^3}{\pi^2 - 8}$$

and

$$\pi_{\frac{2}{3}}^2 c_{\frac{2}{3}} = \frac{64\pi^2}{27\sqrt[3]{4}} < \frac{\pi^3}{\pi^2 - 8},$$

then  $\frac{4}{3} < p_0 < \frac{3}{2}$ . This settles the proof of Theorem 1 for  $1 < p < 2$ .

**Remark 1.** An implementation of the Newton method gives  $p_0 \approx 1.458801$  as an approximated solution of (13) with all digits correct.

## 5.2 Case $p > 2$

Recall the following identities involving the Riemann Zeta function [11, 3.411, 9.522 & 9.524],

$$(15) \quad \zeta(q) = \frac{1}{\Gamma(q)} \int_0^{\infty} \frac{t^{q-1}}{e^t - 1} dt \quad \operatorname{Re}(q) > 1,$$

$$(16) \quad \sum_{\substack{j=1 \\ j \neq 2^0}}^{\infty} \frac{1}{j^q} = \left(1 - \frac{1}{2^q}\right) \zeta(q)$$

and

$$(17) \quad \frac{\zeta'(q)}{\zeta(q)} = - \sum_{k=1}^{\infty} \frac{\Delta(k)}{k^q}$$

where

$$\Delta(k) = \begin{cases} \ln(r) & \text{if } k = r^m \text{ for some } r \text{ prime and } m \in \mathbb{N} \\ 0 & \text{otherwise.} \end{cases}$$

**Lemma 10.** *Let*

$$t_0 = \frac{2(e^2 - 3e + 1)}{(e^2 - 2e - 1)}.$$

*Then*

$$(18) \quad \zeta\left(\frac{3}{2}\right) < \frac{2}{\sqrt{\pi}} \left( 2\sqrt{2} \arctan \frac{1}{\sqrt{2}} + \frac{\pi^2}{6} + \frac{t_0^2}{4} - \frac{(t_0 - 1)^2}{2(e-1)^2} - \frac{t_0(e-2)+1}{e-1} \right).$$

*Proof.* Since  $\Gamma(1 + \frac{1}{2}) = \frac{\sqrt{\pi}}{2} 1!! = \frac{\sqrt{\pi}}{2}$ , the representation (15) gives

$$\begin{aligned} \zeta\left(\frac{3}{2}\right) &= \frac{2}{\sqrt{\pi}} \int_0^{\infty} \frac{t^{1/2}}{e^t - 1} dt \\ &= \frac{2}{\sqrt{\pi}} \left( \int_0^1 + \int_1^{\infty} \frac{t^{1/2}}{e^t - 1} dt \right) = \frac{2}{\sqrt{\pi}} (J_1 + J_2). \end{aligned}$$

We estimate separately upper bounds for  $J_1$  and  $J_2$ .

The change of variables  $t = u^2$ , yields

$$\begin{aligned} J_1 &= \int_0^1 \frac{t^{1/2}}{e^t - 1} dt < \int_0^1 \frac{t^{1/2}}{t + \frac{t^2}{2}} dt \\ &= \int_0^1 \frac{2u^2}{u^2 + \frac{u^4}{2}} du = 2\sqrt{2} \arctan \frac{1}{\sqrt{2}}. \end{aligned}$$

On the other hand, we know that  $\zeta(2) = \int_0^{\infty} \frac{t}{e^t - 1} dt = \frac{\pi^2}{6}$ , so

$$J_2 \leq \int_1^{\infty} \frac{t}{e^t - 1} dt = \frac{\pi^2}{6} - \int_0^1 \frac{t}{e^t - 1} dt.$$

We find lower bound for the integral on the right hand side, by interpolating the curve  $c(t) = \frac{t}{e^t - 1}$  at two points,  $t = 0$  and  $t = 1$ . Firstly observe that  $c(t) \rightarrow 1$  as  $t \rightarrow 0$ ,  $c(t)$  is decreasing and  $c''(t) \geq 0$  for  $t \in [0, 1]$ . Let  $t_0$  be as in the hypothesis and let

$$\tilde{c}(t) = \begin{cases} 1 - \frac{1}{2}t & 0 \leq t \leq t_0 \\ \frac{1}{(e-1)^2}(1-t) + \frac{1}{e-1} & t_0 \leq t \leq 1 \end{cases}$$

be the piecewise linear interpolant of  $c(t)$  in the two segments  $[0, t_0]$  and  $[t_0, 1]$ , which is continuous at  $t_0$ . Note that  $\tilde{c}(t)$  and  $c(t)$  are tangent at  $t = 0$  and  $t = 1$ . Then

$$c(t) \geq \tilde{c}(t) \quad \forall t \in [0, 1].$$

Hence

$$\begin{aligned} \int_0^1 c(t) dt &\geq \int_0^{t_0} \left(1 - \frac{1}{2}t\right) dt + \int_{t_0}^1 \left(\frac{1}{(e-1)^2}(1-t) + \frac{1}{e-1}\right) dt \\ &= -\frac{t_0^2}{4} + \frac{(t_0-1)^2}{2(e-1)^2} + \frac{t_0(e-2)+1}{e-1}. \end{aligned}$$

Thus

$$J_2 \leq \frac{\pi^2}{6} + \frac{t_0^2}{4} - \frac{(t_0-1)^2}{2(e-1)^2} - \frac{t_0(e-2)+1}{e-1}.$$

Alongside with the upper bound above for  $J_1$ , this ensures the validity of the claimed statement.  $\square$

Now, consider the equation

$$(19) \quad \frac{2\pi p'}{\pi^2(p-1)} \left[2 + \frac{\pi^2}{2}(p-2)\right] \left[\left(1 - \frac{1}{2p'}\right) \zeta(p') - 1\right] = \frac{8}{\pi\pi_p}.$$

**Lemma 11.** *There exists  $p_1 \in (\frac{11}{5}, 3)$  such that (19) holds true for  $p = p_1$ . Moreover,*

$$\frac{2\pi p'}{\pi^2(p-1)} \left[2 + \frac{\pi^2}{2}(p-2)\right] \left[\left(1 - \frac{1}{2p'}\right) \zeta(p') - 1\right] < \frac{8}{\pi\pi_p} \quad \forall p \in [2, p_1].$$

*Proof.* From (4) it follows that the identity (19) reduces to

$$(20) \quad \frac{\pi}{p^2 \sin(\frac{\pi}{p})^2} \left(2 + \frac{\pi^2}{2}(p-2)\right) \left[\left(1 - \frac{1}{2p'}\right) \zeta(p') - 1\right] = 1.$$

Denote by  $h(p)$  the left hand side of (20). Then  $h : (1, \infty) \rightarrow \mathbb{R}$  is continuous and

$$h(2) = \frac{\pi}{2} \left(\frac{\pi^2}{8} - 1\right) < 1.$$

Since

$$\zeta\left(\frac{3}{2}\right) > 1 + \frac{\sqrt{2}}{4} + \sqrt{3} \sum_{k=3}^{\infty} \frac{1}{k^2} = \frac{4 + \sqrt{2}}{4} + \sqrt{3} \left(\frac{\pi^2}{6} - \frac{5}{4}\right),$$

we get

$$\begin{aligned} h(3) &= \frac{\pi}{9 \sin(\frac{\pi}{3})^2} \left[2 + \frac{\pi^2}{2}\right] \left[\left(1 - \frac{1}{2 \cdot \frac{3}{2}}\right) \zeta\left(\frac{3}{2}\right) - 1\right] \\ &> \frac{\pi}{9 \sin(\frac{\pi}{3})^2} \left[2 + \frac{\pi^2}{2}\right] \left[\left(1 - \frac{1}{2 \cdot \frac{3}{2}}\right) \left(\frac{4 + \sqrt{2}}{4} + \sqrt{3} \left(\frac{\pi^2}{6} - \frac{5}{4}\right)\right) - 1\right] \\ &> 1. \end{aligned}$$

Hence, there exists  $p_1 \in (2, 3)$  such that  $h(p_1) = 1$ .

The derivative

$$\frac{d}{dq} \left[ \left(1 - \frac{1}{2^q}\right) \zeta(q) \right] = \frac{\ln(2)}{2^q} \zeta(q) + \left(1 - \frac{1}{2^q}\right) \zeta'(q)$$

is negative for any  $q \in (1, 2)$ . Indeed the identity (17) gives

$$\begin{aligned} \frac{\zeta'(q)}{\zeta(q)} &< -\frac{\ln(2)}{2^q} - \frac{\ln(3)}{3^q} - \frac{\ln(2)}{4^q} \\ &< -\ln(2) \left[ \frac{1}{2^q} + \frac{1}{3^q} + \frac{1}{4^q} \right] < \frac{\ln(2)}{1 - 2^q}, \end{aligned}$$

so that

$$\frac{d}{dq} \left[ \left(1 - \frac{1}{2^q}\right) \zeta(q) \right] = \zeta(q) \left[ \frac{\ln(2)}{2^q} + \frac{2^q - 1}{2^q} \frac{\zeta'(q)}{\zeta(q)} \right] < 0.$$

Since  $p'$  and  $\sin\left(\frac{\pi}{p}\right)$  are decreasing functions of  $p > 2$ , then

$$\frac{\pi}{\sin\left(\frac{\pi}{p}\right)^2} \left[ \left(1 - \frac{1}{2^{p'}}\right) \zeta(p') - 1 \right]$$

is an increasing function of  $p > 2$ .

As

$$\frac{d}{dp} \left[ \frac{1}{p^2} \left(2 + \frac{\pi^2}{2}(p-2)\right) \right] = \frac{1}{p^3} \left(-\frac{\pi^2}{2}p + 2\pi^2 - 4\right) > 0 \quad \forall p \in [2, 3],$$

then  $h(p)$  is increasing for  $p \in [2, 3]$  and so indeed

$$h(p) < h(p_1) = 1 \quad \forall p \in [2, p_1].$$

Let us now show that  $p_1 > \frac{11}{5}$ . Let  $c_1$  denote the right hand side of the estimate (18) in Lemma 10. Since  $\zeta(q)$  is convex in the segment  $[\frac{3}{2}, 2]$ , then

$$\zeta(q) \leq \left(\frac{\pi^2}{3} - 2c_1\right) (q-2) + \frac{\pi^2}{6}.$$

That is, the straight line joining the points  $(\frac{3}{2}, c_1)$  and  $(2, \frac{\pi^2}{6})$  is above the curve  $\zeta(q)$  for all  $q \in [\frac{3}{2}, 2]$ . Then

$$(21) \quad \zeta\left(\frac{11}{6}\right) \leq \frac{\pi^2}{9} + \frac{c_1}{3}.$$

Note that for  $p = \frac{11}{5}$ ,  $p' = \frac{11}{6}$ . Now,  $\sin(\pi y)$  is concave for  $y \in [\frac{5}{12}, \frac{1}{2}]$ . Then it is above the straight line joining the points  $(\frac{5}{12}, \sin \frac{5\pi}{12})$  and  $(\frac{1}{2}, 1)$ . That is

$$\sin(\pi y) \geq \left(12 - 12 \sin \frac{5\pi}{12}\right) \left(y - \frac{1}{2}\right) + 1 \quad \forall y \in \left[\frac{5}{12}, \frac{1}{2}\right].$$

Then

$$(22) \quad \sin \frac{5\pi}{11} > \frac{\sqrt{6}}{22}(\sqrt{3} + 3) + \frac{5}{11}.$$

Denote by  $c_2$  the right hand side of the latter inequality. From (21) and (22), it follows that

$$\begin{aligned} h\left(\frac{11}{5}\right) &= \frac{\pi}{\left(\frac{11}{5}\right)^2 \sin\left(\frac{5\pi}{11}\right)^2} \left[2 + \frac{\pi^2}{2} \left(\frac{11}{5} - 2\right)\right] \left[\left(1 - \frac{1}{2^{11/6}}\right) \zeta\left(\frac{11}{6}\right) - 1\right] \\ &< \frac{\pi}{\frac{121}{25} c_2^2} \left(2 + \frac{\pi^2}{10}\right) \left[\left(1 - \frac{1}{2^{11/6}}\right) \left(\frac{\pi^2}{9} + \frac{c_1}{3}\right) - 1\right] < 1. \end{aligned}$$

As  $h(p)$  is increasing, then indeed  $p_1 > \frac{11}{5}$ .  $\square$

**Corollary 2.** *Let  $p_1 > 2$  be such that (19) holds true for  $p = p_1$ . The family  $\{\cos_p(n\pi_p)\}_{n=0}^\infty$  forms a Schauder basis of  $L_s$  for all  $s > 1$  and  $2 \leq p \leq p_1$ .*

*Proof.* From Lemma 7 and (16), we have

$$(23) \quad \sum_{\substack{j=3 \\ j=2^1}}^{\infty} |b_j| < \frac{2\pi_{p'}}{\pi^2(p-1)} \left[2 + \frac{\pi^2}{2}(p-2)\right] \left[\left(1 - \frac{1}{2^{p'}}\right) \zeta(p') - 1\right].$$

According to part “b” of Lemma 1,  $\sin_p(\pi_p x)$  is strictly concave on  $(0, \frac{1}{2})$ . Then

$$\begin{aligned} a_1 &= 2 \int_0^1 \sin_p(\pi_p x) \sin(\pi x) dx = 4 \int_0^{\frac{1}{2}} \sin_p(\pi_p x) \sin(\pi x) dx \\ &> 4 \int_0^{\frac{1}{2}} (2x) \sin(\pi x) dx = \frac{8}{\pi^2}. \end{aligned}$$

Hence, in view of Lemma 5, we get

$$(24) \quad b_1 = \frac{\pi}{\pi_p} a_1 > \frac{8}{\pi \pi_p}.$$

From Lemma 11, it then follows that

$$\sum_{\substack{j=3 \\ j=2^1}}^{\infty} |b_j(p)| < b_1(p) \quad \forall p \in [2, p_1].$$

By virtue of (12) this implies the claimed conclusion.  $\square$

**Remark 2.** *An approximation of the solution of (19) via the Newton Method gives  $p_1 \approx 2.42865$  with all digits correct.*

## 6 Connections with other work

In this final section we describe various connections between the statements established above and those reported in the literature.

### The $p$ -exponential functions

Let

$$\exp_p(iy) = \cos_p(y) + i \sin_p(y) \quad \forall y \in \mathbb{R}.$$

By combining Theorem 1 with [2, Theorem 1] or [5, Theorem 4.5], it immediately follows that the family  $\tilde{\mathcal{F}} = \{\exp_p(in\pi_p \cdot)\}_{n=-\infty}^{\infty}$  is a Schauder basis of the Banach space  $L_s(-1, 1)$  for all  $p \in [p_0, p_1]$ .

Indeed, recall that every  $f \in L^s(-1, 1)$  decomposes as  $f = f_e + f_o$  for

$$f_e(x) = \frac{f(x) + f(-x)}{2} \quad \text{and} \quad f_o(x) = \frac{f(x) - f(-x)}{2},$$

the even and odd parts of  $f$ , respectively. The family  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^{\infty}$  comprises only even functions, the family  $\{\sin_p(n\pi_p \cdot)\}_{n=1}^{\infty}$  comprises only odd functions and they are Schauder bases of the corresponding subspaces of  $L_s(-1, 1)$  for  $p \in [p_0, p_1]$ . This implies that there exist two unique scalar sequences  $(\alpha_k)_{k=0}^{\infty}$  and  $(\beta_k)_{k=1}^{\infty}$ , such that

$$f(\cdot) = \alpha_0 + \sum_{k=1}^{\infty} \alpha_k \cos_p(k\pi_p \cdot) + i\beta_k \sin_p(k\pi_p \cdot)$$

in  $L_s(-1, 1)$ . In order to see this, one expands  $f_e$  in  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^{\infty}$  and  $f_o$  in  $\{\sin_p(n\pi_p \cdot)\}_{n=1}^{\infty}$ , in the corresponding even and odd subspaces.

By letting  $c_0 = \alpha_0$ ,

$$c_k = \frac{\alpha_k + \beta_k}{2} \quad \text{and} \quad c_{-k} = \frac{\alpha_k - \beta_k}{2} \quad \forall k \in \mathbb{N},$$

we get

$$f(\cdot) = \sum_{k=-\infty}^{\infty} c_k \exp_p(ik\pi_p \cdot)$$

in  $L_s(-1, 1)$ . Since there is a 1:1 correspondence between the scalar sequences via

$$\alpha_k = c_k + c_{-k} \quad \text{and} \quad \beta_k = c_k - c_{-k},$$

then in fact  $(c_k)_{k=-\infty}^{\infty}$  is unique for the given  $f$ . Thus,  $\tilde{\mathcal{F}}$  satisfies the definition of a Schauder basis for the Banach space  $L_s(-1, 1)$ .

### The regularity of the $p$ -sine functions

Let  $r > 0$  and denote by  $H^r \equiv H^r(0, 1)$  the (Hilbert) Sobolev space of order  $r$ . Let  $1 < p < 2$ . According to the formula [5, (4.4)], it follows that the Fourier coefficients of the  $p$ -sine function are such that

$$|a_j(p)| \leq \frac{16\pi_p^2 c_p}{\pi^3} j^{-3} \quad \forall j \in \mathbb{N}.$$

Then,  $\sin_p(\pi_p \cdot) \in H^\rho$  for all  $\rho < \frac{5}{2}$ .

Numerical estimates for the Sobolev regularity of  $\sin_p(\pi_p \cdot)$  for  $2 < p < 100$  were reported in [3, Figure 2]. From that picture, one may conjecture that for  $p > 3$ ,  $\sin_p(\pi_p \cdot) \notin H^2$ . Moreover, the regularity appears to drop asymptotically to  $\frac{3}{2}$  for  $p$  large. By contrast, it appears that  $\sin_p(\pi_p \cdot) \in H^2$  for  $2 < p < 3$ . The following statement, which is a consequence of Lemma 7, settles this conjecture.

**Corollary 3.** *For  $p > 2$  set  $r(p) = p' + \frac{1}{2}$ . Then  $\sin_p(\pi_p \cdot) \in H^\rho$  for all  $0 \leq \rho < r(p)$ .*

*Proof.* According to Lemma 5,

$$|a_j(p)| = \frac{\pi_p}{j\pi} |b_j(p)|.$$

Then, by virtue of Lemma 7,

$$|a_j(p)| \leq \frac{2\pi_p \pi_{p'}}{\pi^3(p-1)} \left[ 2 + \frac{\pi^2}{2}(p-2) \right] j^{-(p'+1)} \quad \forall j \geq 3.$$

Let  $\langle j \rangle^2 = 1 + j^2$ . For  $\rho < p' + \frac{1}{2}$ ,

$$\sum_{j=1}^{\infty} \langle j \rangle^{2\rho} |a_j(p)|^2 \leq 2^\rho a_1(p)^2 + c(p) \sum_{\substack{j=3 \\ j \equiv 2 \pmod{1}}}^{\infty} \frac{1}{j^{1+\epsilon(p)}} < \infty$$

where

$$c(p) = \frac{2\pi_p \pi_{p'}}{\pi^3(p-1)} \left[ 2 + \frac{\pi^2}{2}(p-2) \right] \quad \text{and} \quad \epsilon(p) = 1 - 2\rho + 2p' > 0.$$

Hence  $\sin_p(\pi_p \cdot) \in H^\rho$  as claimed.  $\square$

The recent paper [8] includes various intriguing results connected to Corollary 3.

### The paper [7]

The recent paper [7] seems to be the only one in the existing literature which conducts an analysis of the basis properties of the  $p$ -cosine functions. In the notation of [7] we fix  $\alpha = 1$  and  $p = q > 1$ . The Fourier coefficients of the  $p$ -cosine functions are

$$\tau_j(p, p, 1) = b_j(p) \quad \forall j \in \mathbb{N} \cup \{0\}.$$

The condition [7, (2.2)] as well as the criterion for determining whether  $\{\cos_p(n\pi_p)\}_{n=0}^\infty$  is a Schauder basis of  $L^s$  are exactly the same as (12). Let us compare some of the results of [7] with those of the present work.

In [7, Proposition 2.5], the estimate [7, (2.20)] is equivalent to the following. There exists  $p_0^* = \frac{72(\pi-2)-2\pi^3}{96(\pi-2)-3\pi^3}$ , such that

$$(25) \quad \tau_1(p, p, 1) \geq \begin{cases} \frac{\pi(p-1)}{2p-1} - \frac{(\pi-2)(p-1)}{3p-2} & 1 < p < p_0^* \\ \frac{\pi(p-1)}{2p-1} - \frac{\pi^3(p-1)}{24(4p-3)} & p_0^* < p < \infty. \end{cases}$$

Here  $p_0^*$  satisfies the identity

$$\frac{4p-3}{3p-2} = \frac{\pi^3}{24(\pi-2)}.$$

Note that  $p_0^* \approx 1.22$ .

Let us consider firstly the regime  $1 < p < 2$ . From [7, Proposition 2.2] it follows that

$$(26) \quad \sum_{k=1}^{\infty} |\tau_{2k+1}(p, p, 1)| \leq \frac{\pi_p(\pi^2 - 8)}{\pi^2} \quad \forall p \in (1, 2).$$

As  $c_p < 1$  whenever  $1 < p < 2$  in (6), then (14) is sharper than (26) in this regime.

If  $1 < p < p_0^*$ , then

$$\frac{\pi_p(\pi^2 - 8)}{\pi^2} > \frac{\pi(p-1)}{2p-1} - \frac{(\pi-2)(p-1)}{3p-2},$$

and no conclusion about the validity of (12) can be derived in this case from (25) and (26). For  $p_0^* < p < 2$ , on the other hand,

$$\frac{\pi_p(\pi^2 - 8)}{\pi^3} < \frac{p-1}{2p-1} - \frac{\pi^2(p-1)}{24(4p-3)} \iff p \in (p_0^\dagger, 2),$$

where  $p_0^\dagger \approx 1.75$ . In order to see this, note that  $\pi_p$  is decreasing and  $\lim_{p \rightarrow 1^+} \pi_p = \infty$ , while the right hand side of this identity is increasing



for  $1 < p < 2$ . Thus, a combination of [7, Proposition 2.2] and [7, Proposition 2.5], only guarantees that  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^\infty$  is a Schauder basis of  $L^s$  for  $p \in [p_0^\dagger, 2)$  where  $p_0^\dagger > \frac{3}{2} > p_0$ .

As it turns, it is not possible to deduce from the results of [7] any basis property of the family  $\{\cos_p(n\pi_p \cdot)\}_{n=0}^\infty$  in the complementary regime  $p > 2$ . Here is how the different estimates on the Fourier coefficients compare in this case.

From [7, Proposition 2.4], we gather that

$$(27) \quad \sum_{k=1}^{\infty} |\tau_{2k+1}(p, p, 1)| \leq \frac{2\pi_{p'}}{\pi^2(p-1)} [4 + \pi(p-1)] \left[ \left(1 - \frac{1}{2^{p'}}\right) \zeta(p') - 1 \right].$$

Since

$$4 + \pi(p-1) \geq 2 + \frac{\pi^2}{2}(p-2) \quad \forall p \leq \frac{4 + 2\pi^2 - 2\pi}{\pi^2 - 2\pi},$$

the upper bound (23) is sharper than (27) for  $2 \leq p \leq 3$ . The latter is the relevant regime in the proof of Theorem 1.

Since  $\pi_p < \pi$  for  $p > 2$ , the lower bound (24) is sharper than [7, (2.19)]. Moreover,

$$\frac{8}{\pi\pi_p} > \frac{\pi(p-1)}{2p-1} - \frac{\pi^3(p-1)}{24(4p-3)} \quad \forall p > 2.$$

Hence the estimate (25), which is [7, (2.20)], is also superseded by (24) for  $p > 2$ .

## Acknowledgements

HM was supported by Heriot-Watt University under the Global Platform Scholarships Scheme.

## 2010 Mathematics Subject Classification

Primary: 42A16. Secondary: 42A65.

## Keywords

Generalised cosine functions, Schauder basis properties,  $p$ -Laplacian.

## References

- [1] A. BEURLING, *The collected works of Arne Beurling. Vol. 1*, Contemporary Mathematicians, Birkhäuser Boston, Inc., Boston, MA, 1989. Complex analysis, Edited by L. Carleson, P. Malliavin, J. Neuberger and J. Wermer. Pages 378–380.

- [2] P. BINDING, L. BOULTON, J. ČEPIČKA, P. DRÁBEK, AND P. GIRG, *Basis properties of eigenfunctions of the  $p$ -Laplacian*, Proc. Amer. Math. Soc., 134 (2006), pp. 3487–3494.
- [3] L. BOULTON AND G. J. LORD, *Approximation properties of the  $q$ -sine bases*, Proc. R. Soc. A, 467 (2011), pp. 2690–2711.
- [4] ———, *Basis properties of the  $p, q$ -sine functions*, Proc. R. Soc. A, 471 (2015).
- [5] P. J. BUSHELL AND D. E. EDMUNDS, *Remarks on generalized trigonometric functions*, Rocky Mountain J. Math., 42 (2012), pp. 25–57.
- [6] D. E. EDMUNDS, P. GURKA, AND J. LANG, *Properties of generalized trigonometric functions*, J. Approx. Theory, 164 (2012), pp. 47–56.
- [7] ———, *Basis properties of generalized trigonometric functions*, J. Math. Anal. Appl., 420 (2014), pp. 1680–1692.
- [8] ———, *Decay of  $(p, q)$ -Fourier coefficients*, Proc. R. Soc. A, 470 (2014), pp. 1–9.
- [9] D. E. EDMUNDS AND J. LANG, *Eigenvalues Embeddings and Generalised Trigonometric Functions*, Springer Verlag, Berlin, 2011.
- [10] Á. ELBERT, *A half-linear second order differential equation*, in Qualitative theory of differential equations, Vol. I, II (Szeged, 1979), vol. 30 of Colloq. Math. Soc. János Bolyai, North-Holland, Amsterdam-New York, 1981, pp. 153–180.
- [11] I. S. GRADSHTEYN AND I. M. RYZHIK, *Table of Integrals, Series and Products*, Elsevier/Academic Press, Amsterdam, seventh ed., 2007.
- [12] J. R. HIGGINS, *Completeness and Basis Properties of Sets of Special Functions*, Cambridge University Press, Cambridge, 1977.
- [13] T. KATO, *Perturbation Theory for Linear Operators*, Springer Verlag, Berlin, 1995.
- [14] P. LINDQVIST AND J. PEETRE, *Erik Lundberg and the hypergoniometric functions*, Normat, 45 (1997), pp. 1–24, 48.
- [15] M. ÔTANI, *A remark on certain nonlinear elliptic equations*, Proc. Fac. Sci. Tokai Univ., 19 (1984), pp. 23–28.
- [16] I. SINGER, *Bases in Banach Spaces I*, Springer Verlag, Berlin, 1970.