



Heriot-Watt University
Research Gateway

Statistics modeling of random polarization in K-distributed sea clutter

Citation for published version:

Pan, Z, Luo, Z, Wang, Y & Wang, W 2024, Statistics modeling of random polarization in K-distributed sea clutter. in OV Angelsky & CY Zenkova (eds), *Sixteenth International Conference on Correlation Optics.*, 129381E, Proceedings of SPIE, vol. 12938, SPIE, 16th International Conference on Correlation Optics 2023, Chernivtsi, Ukraine, 18/09/23. <https://doi.org/10.1117/12.3013059>

Digital Object Identifier (DOI):

[10.1117/12.3013059](https://doi.org/10.1117/12.3013059)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Sixteenth International Conference on Correlation Optics

Publisher Rights Statement:

© 2024 SPIE.

Proceedings Volume 12938, Sixteenth International Conference on Correlation Optics; 129381E (2024)
<https://doi.org/10.1117/12.3013059>

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Statistics modeling of random polarization in K-distributed sea clutter

Zhongbao Pan,^a Zuishuang Luo,^a Ying Wang,^a and Wei Wang^{b,*}

^a Department of Physics, Jiangsu University of Science and Technology, Zhenjiang, China 212003;

^b School of Engineering and Physical Sciences, Heriot-Watt University, Edinburgh, UK EH14 4AS

Corresponding Author: w.wang@hw.ac.uk

ABSTRACT

K-distribution plays a significant role in describing the statistics of the electromagnetic scattering. The influence of shape factor on the K-distribution is very significant because different shape factors in K-distributed sea clutter represent different actual scenarios. In this paper, we investigate the impact of different shape factors in the K-distributed sea clutter and provide the statistics of the Stokes parameters in electromagnetic scattering of the K-distribution under different shape factors.

Keywords: K-distribution, polarization phasor sum, shape factor

1. INTRODUCTION

The K-distribution occupies an important position in the sea clutter model, which can describe the high-resolution low incidence residual angle well [1]. Up to now, speckle fields involving the use of the K-distribution have been treated as scalar fields ignoring the polarization properties arising from the vector nature of the electromagnetic fields [2]. We investigate the statistics of the random polarization of sea clutter waves with K-distribution and provide the probability density function of the Stokes parameter characterizing the fluctuating polarization [3]. The shape parameter is an important variable in the statistical model of sea clutter that indirectly reflects radar and environmental parameters. We investigate the impact of different shape factors in the K-distributed sea clutter and provide the statistics of the Stokes parameters in electric scattering of the K-distribution under different shape factors [4].

2. STATISTICAL PROPERTIES OF STOKES PARAMETERS FOR RANDOM PHASE SUMS

Usually, the polarization state of the light can be described by the Stokes parameters, which can be given as [5]:

$$\begin{aligned} S_0 &= A_x^2 + A_y^2 = I_x + I_y \\ S_1 &= A_x^2 - A_y^2 = I_x - I_y \\ S_2 &= 2A_x A_y \cos(\theta_y - \theta_x) = 2A_x A_y \cos \delta \\ S_3 &= 2A_x A_y \sin(\theta_y - \theta_x) = 2A_x A_y \sin \delta \end{aligned} \quad (1)$$

where I_x and I_y are the intensities of two polarization components and δ is the random phase difference between two

polarization components.

On the other hand, the K-distribution is a family of continuous probability distributions with parameters, so the probability density function whose amplitude follows the K-distribution is therefore expressed as follows [6]:

$$p_A(A) = \frac{2\beta}{\Gamma(1+\nu)} \left(\frac{\beta A}{2}\right)^{\nu+1} K_\nu(\beta A) \quad (2)$$

valid for $\nu > -1$ and $A \geq 0$. Here, ν and β are two parameters used to fit the density function to experimental data, where ν is the shape factor and $K_\nu(\dots)$ is the modified Bessel function of the second kind.

2.1 Statistics of S_0 and S_1

Finding the statistics of S_0 and S_1 are complicated that the probability density function of the intensity can be found with the transformation rule:

$$p_I(I) = \frac{2\beta^{(\nu+1)/2} I^{(\nu-1)/2}}{\Gamma(\nu)} K_{\nu-1}(2\sqrt{\beta I}) \quad (3)$$

The degree of polarization \mathcal{P} is introduced and the relationship between the intensity statistics of light polarized in different directions, we can obtain the respective probability density functions of the two components I_x and I_y :

$$p_{I_m}(I_m) = \frac{2}{I_m \Gamma(\nu)} (2\nu I_m / [(1 \pm \mathcal{P}) \overline{S_0}])^{(\nu+1)/2} K_{\nu-1}(2\sqrt{2\nu I_m / [(1 \pm \mathcal{P}) \overline{S_0}]}), m = x, y \quad (4)$$

The Stokes parameter S_0 is the total intensity in the speckle field. Its probability density function can be obtained by the convolution of two components of the probability density function p_{I_x} and p_{I_y} . That is:

$$p_{S_0}(S_0) = \int_0^{S_0} p_{I_x}(u) p_{I_y}(S_0 - u) du \quad (5)$$

In order to study the influence of form factor ν on the K-distribution, substitute Eq. (4) into Eq. (5). Figure 1 shows the probability density function of S_0 after normalization at the different number of \mathcal{P} for $\nu = 1$.




Figure1. For $\nu = 1$, the intensity probability density function of S_0 with the different number of \mathcal{P} .

The Stokes parameter S_1 characterizes whether the light is more closely polarized in the x-direction or the y-direction. For S_1 , we have the following equation:

$$p_{S_1}(S_1) = \begin{cases} \int_{S_1}^{\infty} p_{I_x}(u)p_{I_y}(u-S_1)du, & S_1 \geq 0 \\ \int_0^{\infty} p_{I_x}(u)p_{I_y}(u-S_1)du, & S_1 < 0 \end{cases} \quad (6)$$

Substitute Eq. (4) into Eq. (6), after the integration operation, we can obtain the probability density function of S_1 with the different number of \mathcal{P} for $\nu=1$:




Figure2. For $\nu=1$, the intensity probability density function of S_1 with the different number of \mathcal{P} .

2.2 Statistics of S_2 and S_3

Next we perform the statistical calculation for S_2 and S_3 . Here we use the monotone transformation $A = \sqrt{I}$ and bring it into Eq. (5) to obtain the probability density function of the amplitude component:

$$p_{A_m}(A_m) = \frac{4}{A_m \Gamma(\nu)} (\sqrt{2\nu} A_m / \sqrt{(1-\mathcal{P})\bar{S}_0})^{(\nu+1)} K_{\nu-1}(2\sqrt{2\nu} A_x / \sqrt{(1-\mathcal{P})\bar{S}_0}), m=x, y \quad (7)$$

By an integral transformation $\gamma = A_x A_y$, The probability density function of $p_\gamma(\gamma)$ can be expressed as:

$$p_\gamma(\gamma) = \int_0^{\infty} p_{A_x}(A_x) p_{A_y}(\gamma/A_x) A_x^{-1} dA_x \quad (8)$$

Then we can get the probability density function of S_2 by the following formula:

$$p_{S_2}(S_2) = \int_{-1}^1 p_\gamma(S_2/2\xi) (2\pi|\xi|\sqrt{1-\xi^2})^{-1} d\xi \quad (9)$$

S_2 characterizes whether the light is closer to the positive 45 degree polarization or the negative 45 degree polarization; S_3 characterizes whether the light is closer to the right-handed circularly polarized light or the left-handed circularly polarized light, so they have the same probability density function distribution. As shown in the following figure:




Figure3. For $\nu=1$, the intensity probability density function of S_2 or S_3 with the different number of \mathcal{P} .

3. CONCLUSION

The statistical properties of the random light field under the K-distribution are analyzed, and the accurate expression of the Stokes parameter of the random polarization phasor sum is obtained. When the degree of shape factor is 1, the probability distributions of intensity and amplitude of polarization in different directions are compared. In a certain class of applications, including synthetic-aperture radar imagery and wireless communications, the distribution plays a significant role in describing the statistical properties of the electromagnetic scattering. In this paper, we have investigated the development of the polarization speckle when the amplitudes of the contributing polarization phasors are distributed and have provided the probability density functions of the Stokes parameters characterizing the fluctuating polarization. These results can be regarded as an extension of previous study on the statistical or probabilistic model with distribution for scalar fields and will provide a deeper insight into the evolution of random polarization for stochastic electromagnetic fields.

REFERENCE

- [1] E. Conte, A. De Maio, and C. Galdi, "Statistical analysis of real clutter at different range resolutions," *IEEE Transactions on Aerospace and Electronic Systems* 40(3), 903-918 (2004).
- [2] S. H. Yueh, J. A. Kong, J. K. Jao, R. T. Shin, and L. M. Novak, "K-distribution and polarimetric terrain radar clutter," *J. Electromagn. Waves Appl* 3(8), 747-768(1989).
- [3] V. Anastassopoulos, G. A. Lampropoulos, A. Drosopoulos, and M. Ray, "High resolution radar clutter statistics," *IEEE Trans. Aerosp. Electron. Syst* 35(1), 43-60 (1999).
- [4] E. Jakeman and P. N. Pusey, "Significance of K distributions in scattering experiments," *Phys. Rev. Lett.* 40, 546-550 (1978).
- [5] Goodman, J. W. (2008) Speckle with a finite number of steps. *Applied optics*, 47(4): A111-A118(2008).
- [6] K. D. Ward, R. J. A. Tough, and S. Watts, "Sea clutter: Scattering, the K distribution and radar performance," *Waves in Random and Complex Media* 17(2), 233-234 (2007).