



Heriot-Watt University
Research Gateway

Fibonacci type presentations and 3-manifolds

Citation for published version:

Howie, J & Williams, G 2017, 'Fibonacci type presentations and 3-manifolds', *Topology and its Applications*, vol. 215, pp. 24–34. <https://doi.org/10.1016/j.topol.2016.10.012>

Digital Object Identifier (DOI):

[10.1016/j.topol.2016.10.012](https://doi.org/10.1016/j.topol.2016.10.012)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Peer reviewed version

Published In:

Topology and its Applications

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Fibonacci type presentations and 3-manifolds

James Howie, Gerald Williams

PII: S0166-8641(16)30277-2

DOI: <http://dx.doi.org/10.1016/j.topol.2016.10.012>

Reference: TOPOL 5943


To appear in: *Topology and its Applications*

Received date: 4 August 2016

Accepted date: 31 October 2016

Please cite this article in press as: J. Howie, G. Williams, Fibonacci type presentations and 3-manifolds, *Topol. Appl.* (2016), <http://dx.doi.org/10.1016/j.topol.2016.10.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Fibonacci type presentations and 3-manifolds

James Howie and Gerald Williams*

November 2, 2016

Abstract

We study the cyclic presentations with relators of the form $x_i x_{i+m} x_{i+k}^{-1}$ and the groups they define. These “groups of Fibonacci type” were introduced by Johnson and Mawdesley and they generalize the Fibonacci groups $F(2, n)$ and the Sieradski groups $S(2, n)$. With the exception of two groups, we classify when these groups are fundamental groups of 3-manifolds, and it turns out that only Fibonacci, Sieradski, and cyclic groups arise. Using this classification, we completely classify the presentations that are spines of 3-manifolds, answering a question of Cavicchioli, Hegenbarth, and Repovš. When n is even the groups $F(2, n), S(2, n)$ admit alternative cyclic presentations on $n/2$ generators. We show that these alternative presentations also arise as spines of 3-manifolds.

Keywords: Fibonacci group, Sieradski group, cyclically presented group, 3-manifold group, spine of a manifold.

MSCs: 20F05, 57M05 (primary), 57M50 (secondary).

1 Introduction

Let F_n be the free group of rank $n \geq 1$ with generators x_0, \dots, x_{n-1} and let $w = w(x_0, \dots, x_{n-1})$ be a word in F_n and let $\theta : F_n \rightarrow F_n$ be an automorphism given by $x_i \mapsto x_{i+1}$ (subscripts mod n). The presentation

$$\mathcal{G}_n(w) = \langle x_0, \dots, x_{n-1} \mid \theta^i(w) \ (0 \leq i \leq n-1) \rangle \quad (1)$$

is called a *cyclic presentation*, and the group $G_n(w)$ that it defines is a *cyclically presented group*. Since cyclic presentations are balanced (that is, they have an equal number of generators and relations) the questions as to which cyclic presentations are spines of 3-manifolds and which cyclically presented groups are

*Part of this research was carried out during a visit by the second named author to the Department of Mathematics at Heriot Watt University in January and February 2016. That visit was financed by the Edinburgh Mathematical Society Research Support Fund. The authors would like to thank the EMS for its support and the second named author would like to thank the Department of Mathematics at Heriot Watt University for its hospitality during that visit.

fundamental groups of 3-manifolds arise and have been considered by many authors (see, for example, [5]). For instance, in [11], Dunwoody provides an algorithm to determine if a cyclic presentation is the spine of a 3-manifold. There is a large body of literature showing that certain cyclic presentations arise as spines of 3-manifolds, which hence define fundamental groups of 3-manifolds (see, for example, [6],[27] and the references therein). Conversely, many theorems show that certain cyclic presentations do not define the fundamental groups of hyperbolic 3-orbifolds (in particular 3-manifolds) of finite volume (see, for example, Theorem 3.1 of any of [31],[1],[8],[6], or see [41]). We contribute to this theory by studying one three-parameter family in detail, namely the cyclic presentations $\mathcal{G}_n(m, k) = \mathcal{G}_n(x_0 x_m x_k^{-1})$ (where $n \geq 1$, $0 \leq m, k \leq n-1$) and the groups $G_n(m, k)$ they define.

The groups $G_n(m, k)$ are known as the *groups of Fibonacci type* $G_n(m, k)$ and were introduced in [26]. Their presentations $\mathcal{G}_n(m, k)$ generalize the presentations $\mathcal{F}(2, n) = \mathcal{G}_n(1, 2)$ of the *Fibonacci groups* $F(2, n)$ and the presentations $\mathcal{S}(2, n) = \mathcal{G}_n(2, 1)$ of the *Sieradski groups* $S(2, n)$ of [40] and the presentations $\mathcal{H}(n, m) = \mathcal{G}_n(m, 1)$ of the groups $H(n, m)$ studied in [15]. Together with the cyclic presentations $\mathcal{G}_n(x_0 x_k x_l)$ studied in [8],[12] they form the triangular cyclic presentations – that is, cyclic presentations where the relators have length three. The groups $G_n(m, k)$ have been studied for their algebraic properties in [1],[7],[15],[24],[26],[43],[44] – see [45] for a survey of such results. In this article we study geometric and topological aspects of the groups $G_n(m, k)$ and their presentations.

Our starting point is two results concerning the Fibonacci and Sieradski cases. The first is that if $n \geq 2$ is even then $\mathcal{F}(2, n)$ is the spine of a closed 3-manifold and hence $F(2, n)$ is the fundamental group of a closed 3-manifold [18],[20],[21],[9] – see Theorem 1. The second is that if $n \geq 2$ then the presentation $\mathcal{S}(2, n)$ is the spine of a closed 3-manifold and hence $S(2, n)$ is the fundamental group of a closed 3-manifold [40],[4] – see Theorem 2. Motivated by these results, in [5, Problem 6] Cavicchioli, Hegenbarth and Repovš asked which presentations $\mathcal{G}_n(m, k)$ are spines of closed 3-manifolds. Related to this is the question as to which groups $G_n(m, k)$ are fundamental groups of closed 3-manifolds. With the exception of the (known to be challenging) groups $H(9, 4)$, $H(9, 7)$ we answer the second question in Theorem A, and this allows us to completely answer the first question in Theorem B.

These theorems show that the only cases where spines are possible are in the cases of the Fibonacci and Sieradski presentations of Theorems 1,2, and the only cases where the group is the fundamental group of a 3-manifold are the cases of the corresponding Fibonacci and Sieradski groups and when it is a finite cyclic group. Contained within this classification is the result (Theorem 3) that for odd $n \geq 3$ the group $F(2, n)$ is the fundamental group of a 3-manifold only when $n = 3, 5$, or 7 (in which case it is cyclic). It has previously been observed ([9, page 55]) that the presentation $\mathcal{F}(2, 3)$ is the spine of a closed 3-manifold and the presentations $\mathcal{F}(2, 5)$, $\mathcal{F}(2, 7)$ are not spines of 3-manifolds and so we can conclude that $\mathcal{F}(2, n)$ is the spine of a manifold if and only if $n = 3$ or n is even, confirming the expectation expressed in [9, page 55].

An alternative presentation of the Fibonacci group $F(2, 2m)$ was obtained in [28, Example 1.2], namely the presentation $\mathcal{G}_m(x_0^{-1}x_1^2x_2^{-1}x_1)$; in the same way we can obtain the alternative presentation $\mathcal{G}_m(x_0x_1^2x_2x_1^{-1})$ of the Sieradski group $S(2, 2m)$. In Theorem C we prove that these alternative presentations are also spines of 3-manifolds.

2 3-manifold groups

By a *3-manifold group* we mean the fundamental group of a (not necessarily closed, compact, or orientable) 3-manifold. A 2-dimensional subpolyhedron P of a compact, connected, 3-manifold with boundary M is called a *spine* of M if M collapses to P . A 2-dimensional subpolyhedron P of a closed, connected, 3-manifold M is called a *spine* of M if $M \setminus \text{Int}B^3$ collapses to P , where B^3 is a 3-ball in M . (See, for example, [32].) We shall say that a finite presentation \mathcal{P} is a *3-manifold spine* if its presentation complex $K_{\mathcal{P}}$ is a spine of a (not necessarily closed, compact, or orientable) 3-manifold. Clearly if \mathcal{P} is a 3-manifold spine then the group it defines is a 3-manifold group. In this section we classify which groups $G_n(m, k)$ are 3-manifold groups.

The following theorems assert that if $n \geq 2$ is even then the Fibonacci group $F(2, n)$ is a 3-manifold group and its presentation $\mathcal{F}(2, n)$ is a 3-manifold spine and that if $n \geq 2$ then the Sieradski group $S(2, n)$ is a 3-manifold group and its presentation $\mathcal{S}(2, n)$ is a 3-manifold spine. Using new methods these theorems have recently been reproved in [3, Theorem 5.4], where some history of these results is also described.

Theorem 1 (([18],[20],[21],[9])). *For each even $n \geq 2$ the presentation complex of $\mathcal{F}(2, n)$ is the spine of a closed 3-manifold. Moreover, it is the spine of the $n/2$ -fold cyclic cover of S^3 branched over the figure eight knot, and this is the unique closed prime 3-manifold for which $\mathcal{F}(2, n)$ is a spine.*

Theorem 2 (([40],[4])). *For each $n \geq 2$ the presentation complex of $\mathcal{S}(2, n)$ is the spine of a closed 3-manifold. Moreover, it is the spine of the n -fold cyclic cover of S^3 branched over the trefoil knot.*

By considering covers and connected sums we have that a free product $G_1 * G_2$ is the fundamental group of a 3-manifold if and only if G_1 and G_2 are fundamental groups of 3-manifolds. Further, using [17, Theorem 1], we have that $G_1 * G_2$ is the fundamental group of a closed 3-manifold if and only if G_1 and G_2 are fundamental groups of closed 3-manifolds. In the case of finitely generated groups, by [38], we have that $G_1 * G_2$ is the fundamental group of a compact 3-manifold if and only if G_1 and G_2 are fundamental groups of compact 3-manifolds.

By [1, Lemma 1.2(1)] we have that $G_n(m, k)$ is isomorphic to the free product of $d = (n, m, k)$ copies of $G_{n/d}(m/d, k/d)$ and so we may assume that $d = 1$. If $(n, k) = 1$ or $(n, m - k) = 1$ then $G_n(m, k) \cong H(n, m')$ for some m' (see [1, Lemma 1.3]). For this reason we can divide our theorem into a statement about the groups $H(n, m)$ and a statement about the groups $G_n(m, k)$ where the parameters

satisfy $(n, m, k) = 1$, $(n, k) > 1$ and $(n, m - k) > 1$. Note that these latter conditions imply that $n \geq 6$, $m \neq k$, $m \neq 0$ and $k \neq 0$. We prove

Theorem A. (a) Suppose $n \geq 1$, $0 \leq m, k \leq n - 1$, $(n, m, k) = 1$, $(n, k) > 1$, $(n, m - k) > 1$. Then $G_n(m, k)$ is a 3-manifold group if and only if $(m, k) = 1$ and either $2k \equiv 0 \pmod{n}$ or $2(m - k) \equiv 0 \pmod{n}$, in which case $G_n(m, k) \cong \mathbb{Z}_s$, where $s = 2^{n/2} - (-1)^{m+n/2}$.

(b) Suppose $n \geq 1$, $(n, m) \neq (9, 4), (9, 7)$. Then $H(n, m)$ is a 3-manifold group if and only if the pair (n, m) is one of the following:

- (i) $(n, 0)$ in which case $H(n, m) \cong \mathbb{Z}_{2^{n-1}}$;
- (ii) $(n, 1)$ in which case $H(n, m) = 1$;
- (iii) $(n, 2)$, in which case $H(n, m) \cong S(2, n)$;
- (iv) $(n, n - 1)$ ($n \geq 4$ is even or $n = 3, 5$, or 7), in which case $H(n, m) \cong F(2, n)$;
- (v) $(5, 3)$ or $(7, 4)$, in which case $H(n, m) \cong F(2, 5) \cong \mathbb{Z}_{11}$ or $H(n, m) \cong F(2, 7) \cong \mathbb{Z}_{29}$, respectively;
- (vi) $(2t - 2, t)$ ($t \geq 3$) in which case $H(n, m) \cong \mathbb{Z}_{2^{t-1}+1}$.

While we are unable to deal with the groups $H(9, 4), H(9, 7)$, Theorem 3.1 of [1] gives that they are not the fundamental groups of hyperbolic 3-orbifolds (in particular, 3-manifolds) of finite volume. Further, in Lemma 6 we shall show that the presentations $\mathcal{H}(9, 4), \mathcal{H}(9, 7)$ are not 3-manifold spines. It therefore seems unlikely that either $H(9, 4)$ or $H(9, 7)$ is a 3-manifold group. It is not known whether the groups $H(9, 4), H(9, 7)$ are finite or infinite, or if their presentations $\mathcal{H}(9, 4), \mathcal{H}(9, 7)$ are aspherical (that is, if their presentation complexes are topologically aspherical).

A group G is of *type FK* if there is a finite aspherical cell complex $K(G, 1)$ whose fundamental group is isomorphic to G . If G is of type FK then its Euler characteristic $\chi(G)$ is defined to be the Euler characteristic of a $K(G, 1)$. An immediate corollary of [37, Theorem 2] is that if G is of type FK and $\chi(G) \geq 1$ then G is a (virtual) 3-manifold group if and only if $G = 1$. In particular, if $G = \pi_1(K_{\mathcal{P}}) \neq 1$ where $K_{\mathcal{P}}$ is an aspherical presentation complex of a finite balanced presentation \mathcal{P} , then G is not a 3-manifold group. (Moreover, since groups defined by aspherical presentations are torsion-free it also follows that G is not the fundamental group of any 3-orbifold.) This is a useful tool for us as, with the exception of the presentations $\mathcal{H}(9, 4), \mathcal{H}(9, 7)$, the aspherical presentations $\mathcal{G}_n(m, k)$ have been classified in [15],[43].

The presentations $\mathcal{F}(2, n)$ are not aspherical. We now show that, nevertheless, if n is odd then $F(2, n)$ is rarely a 3-manifold group.

Theorem 3. *If $n \geq 3$ is odd then $F(2, n)$ is a 3-manifold group if and only if $n = 3, 5$, or 7 .*

Proof. If $n = 3, 5$, or 7 then $G = F(2, n)$ is a finite cyclic group so assume that $n \geq 9$ is odd. Let $g = x_0 x_1 \dots x_{n-1}$. We have that $g^2 = 1$ ([25, page 83]),

and by (the proof of) [1, Proposition 3.1] we have that $G/\langle\langle g \rangle\rangle \cong G^{\text{ab}}$. Now, as argued in [1, Proposition 3.1], G^{ab} is finite and for $n \geq 9$ the group G is infinite (see [34],[23],[10],[42]) so $G \not\cong G^{\text{ab}}$ and hence g is an element of G of order 2.

Suppose that G is the fundamental group of a 3-manifold M . Since $G/\langle\langle g \rangle\rangle \cong G^{\text{ab}}$ we have that $g \in [G, G]$ and hence g is orientation preserving and so $\langle g \rangle \cong \mathbb{Z}_2$ is an orientation preserving subgroup of $G = \pi_1(M)$ of finite order. Then by [13, Theorem 8.2] (see also [19, Theorem 9.8]) we have that $M = R \# M_1$, where R is closed and orientable, $\pi_1(R)$ is finite, and $\langle g \rangle$ is conjugate to a subgroup of $\pi_1(R)$. Since G is infinite we have that $\pi_1(M_1) \neq 1$ and since it is a 2-generator group, Gruško's Theorem implies that $\pi_1(R)$ and $\pi_1(M_1)$ are each cyclic. But the derived subgroup of a free product of cyclic groups is free contradicting our deduction that $g \in [G, G]$ is an element of order 2. \square

The classification in [15] gives that the presentations $\mathcal{H}(6, 3)$, $\mathcal{H}(7, 3)$, $\mathcal{H}(7, 5)$, $\mathcal{H}(8, 3)$, $\mathcal{H}(9, 3)$, $\mathcal{H}(9, 6)$ are not aspherical. Again, we can show that the corresponding groups are not 3-manifold groups.

Lemma 4. *The following groups are not 3-manifold groups: $H(6, 3)$, $H(7, 3) \cong H(7, 5)$, $H(8, 3)$, $H(9, 3) \cong H(9, 6)$.*

Proof. For $H(6, 3)$, $H(8, 3)$ and $H(7, 3) \cong H(7, 5)$ we use the classification of abelian subgroups of 3-manifold groups given in [13, Theorem 9.1] (see also, for example [19, Theorem 9.13]). We have that $H(6, 3) \cong \mathbb{Z}_2^3 \times \mathbb{Z}_7$ ([26]) and so contains \mathbb{Z}_2^3 , and so $H(6, 3)$ is not a 3-manifold group. A calculation using GAP [14] shows that the third derived subgroup of $H(8, 3)$ is isomorphic to \mathbb{Z}_3^6 , and so $H(8, 3)$ is not a 3-manifold group. Using quoticpic R.M.Thomas showed (see [15, Theorem 7.3]) that the second derived subgroup of $H(7, 3) \cong H(7, 5)$ is isomorphic to \mathbb{Z}^8 , and so $H(7, 3) \cong H(7, 5)$ is not a 3-manifold group.

Consider now the group $H = H(9, 3)$ (which, as noted in [15], is isomorphic to $H(9, 6)$). Using the van Kampen diagram given in [45, Figure 2] we may see that, setting $g = x_0 x_4 x_8 x_3 x_7 x_2 x_6 x_1 x_5$ the relation $g^2 = 1$ holds in $H = H(9, 3)$. A calculation in GAP gives that $H(9, 3)/\langle\langle g \rangle\rangle$ is a finite group of order $2^{15} \cdot 7$, and since $H(9, 3)$ is infinite [7, Lemma 15] we deduce that g is an element of order exactly 2 in H .

Suppose for contradiction that H is a 3-manifold group. Since H is finitely presentable there is a compact 3-manifold M with $\pi_1(M) \cong H$ ([38]). We have that $H^{\text{ab}} \cong \mathbb{Z}_7$ and hence there is no epimorphism of H onto \mathbb{Z}_2 , so M is orientable. Let $M = M_1 \# \dots \# M_l$ be the unique prime decomposition of M corresponding to a maximal free product decomposition $H_1 * \dots * H_l$ of H . Since M is orientable no M_j contains a 2-sided projective plane so, by [19, Corollary 9.9], each $H_j = \pi_1(M_j)$ is either finite or torsion-free. Since H is infinite and has non-trivial torsion it must therefore be a non-trivial free product. Another calculation in GAP shows that H has a 3-generator presentation so by Gruško's Theorem at least one of the free factors must be cyclic. But $H^{\text{ab}} \cong \mathbb{Z}_7$ so we have that $H \cong \mathbb{Z}_7 * Q$, where Q is perfect. Then the derived subgroup H' , being the free product of 7 copies of Q , must also be perfect, but a calculation in GAP shows that $(H')^{\text{ab}} \cong \mathbb{Z}_2^6$ so H' is not perfect, a contradiction. \square

The above results, together with the asphericity classifications mentioned earlier, combine to prove Theorem A.

Proof of Theorem A. (a) As described above, by [37, Theorem 2] we may assume that the presentation $\mathcal{G}_n(m, k)$ is not aspherical. Theorem 2 of [43] gives that $(m, k) = 1$ and either $2k \equiv 0 \pmod n$ or $2(m - k) \equiv 0 \pmod n$, in which case (by [43, Lemma 3]) $G_n(m, k) \cong \mathbb{Z}_s$, where $s = 2^{n/2} - (-1)^{m+n/2}$, and so is a 3-manifold group.

(b) If $m = 1$ then $H(n, m) = 1$ so we may assume $m \neq 1$. By [37, Theorem 2] we may assume that the presentation $\mathcal{H}(n, m)$ is not aspherical. Then Theorem 3.2 of [15] gives that (n, m) is one of the following: $(n, 0)$, $(n, 2)$ ($n \geq 2$), $(n, n - 1)$ ($n \geq 3$), $(2t - 1, t)$ ($t \geq 3$), $(2t - 2, t)$ ($t \geq 3$), $(6, 3)$, $(7, 3)$, $(7, 5)$, $(8, 3)$, $(9, 3)$, $(9, 6)$. If $(n, m) = (n, 0)$ then $H(n, m) \cong \mathbb{Z}_{2^n - 1}$ (by [15, Proposition 2.2(b)]) so it is a 3-manifold group. If $(n, m) = (n, 2)$ we have $H(n, m) = S(2, n)$ so it is a 3-manifold group by Theorem 2. If $(n, m) = (n, n - 1)$ then $H(n, m) \cong F(2, n)$ so it is a 3-manifold group if and only if $n = 3, 5$, or 7 or n is even by Theorem 1 and Theorem 3. If $(n, m) = (2t - 1, t)$ ($t \geq 3$) then $H(n, m) \cong F(2, 2t - 1)$ so is a 3-manifold group if and only if $t = 3$ or 4 by Theorem 3. If $(n, m) = (2t - 2, t)$ then $H(n, m) \cong \mathbb{Z}_{2^{t-1} + 1}$ by [15, Proposition 2.2(c)], so is a 3-manifold group. In the remaining cases $H(n, m)$ is not a 3-manifold group by Lemma 4. \square

The groups $G_n(x_0 x_m x_k^{-1})$ belong to the class of groups $G_n^\epsilon(m, k, h)$ considered in [8]. Problem 3.4 of that paper asks for which values of the parameters m, k, h, ϵ the group $G_n^\epsilon(m, k, h)$ is the fundamental group of a closed, connected, orientable 3-manifold for infinitely many n . Theorem A can therefore be viewed as a starting point for answering this question. The cyclically presented groups $G_n(x_0 x_k x_l)$ considered in [8], [12] also belong to the class $G_n^\epsilon(m, k, h)$ and so we record without proof the following related result. This can be obtained from the results of [12] together with arguments similar to those used above, and with the fact that non-cyclic metacyclic groups of odd order are not 3-manifold groups (see, for example, [35, page 111]).

Theorem 5. *Suppose $(n, k, l) = 1$. The cyclically presented group $G_n(x_0 x_k x_l)$ is a 3-manifold group if and only if it is isomorphic to one of the following groups: \mathbb{Z}_3 , $\mathbb{Z}_{2^n - (-1)^n}$, $\mathbb{Z} * \mathbb{Z} * \mathbb{Z}_{19}$, $\mathbb{Z} * \mathbb{Z} * \mathbb{Z}_\gamma$ ($\gamma = (2^{n/3} - (-1)^{n/3})/3$), $\mathbb{Z} * \mathbb{Z}$.*

(The conditions on the parameters n, k, l that identify when each group occurs can be extracted from [12, Table 1].)

3 Presentations as spines of manifolds

A wedge $P_1 \vee P_2$ of two 2-dimensional subpolyhedra of a 3-manifold M is a spine of M if and only if P_1 and P_2 are the spines of 3-manifolds M_1 and M_2 where $M = M_1 \# M_2$ (see [29, Lemma 1]). The argument of [1, Lemma 1.2(1)] shows that the presentation complex of $\mathcal{G}_n(m, k)$ is the wedge of $d = (n, m, k)$ copies of the presentation complex of $\mathcal{G}_{n/d}(m/d, k/d)$ so again we may assume that $d = 1$. Similarly, the proof of [1, Lemma 1.3] gives that if $(n, k) = 1$ or $(n, m - k) = 1$ then

the presentation complex of $\mathcal{G}_n(m, k)$ is homeomorphic to the presentation complex of $\mathcal{H}(n, m')$ for some m' so, as with Theorem A, we may divide the statement into two parts. Note that in part (b) we no longer include the hypotheses that $(n, m) \neq (9, 4), (9, 7)$. We use the symbol \simeq to denote homeomorphism of (the presentation complexes) of presentations.

Theorem B. (a) Suppose $n \geq 1$, $0 \leq m, k \leq n - 1$, $(n, m, k) = 1$, $(n, k) > 1$, $(n, m - k) > 1$. Then $\mathcal{G}_n(m, k)$ is not a 3-manifold spine.

(b) Suppose $n \geq 1$. Then $\mathcal{H}(n, m)$ is a 3-manifold spine if and only if one of the following holds:

- (i) $m = 1$, in which case $\mathcal{H}(n, m)$ is a spine of S^3 ;
- (ii) $m = 2$, in which case $\mathcal{H}(n, m) \simeq \mathcal{S}(2, n)$;
- (iii) $m = n - 1$ and either $n = 3$ or n is even, in which case $\mathcal{H}(n, m) \simeq \mathcal{F}(2, n)$.

In Lemmas 6,7,8 we shall show that certain presentations are not 3-manifold spines. In Lemma 9 we show that the presentation $\mathcal{H}(n, 1)$ is a 3-manifold spine. The remaining presentations are either 3-manifold spines by Theorems 1,2, or do not define 3-manifold groups, by Theorem A, so are not 3-manifold spines.

Recall that the *Whitehead graph* of a presentation $\mathcal{P} = \langle X \mid R \rangle$ is the graph with vertices v_x, v'_x ($x \in X$) and an edge (v_x, v_y) (resp. (v'_x, v'_y) , (v_x, v'_y)) for each occurrence of a cyclic subword xy^{-1} (resp. $x^{-1}y$, $(xy)^{\pm 1}$) in a relator $r \in R$. Writing v_i, v'_i in place of v_{x_i}, v'_{x_i} the Whitehead graph of the cyclic presentation $\mathcal{G}_n(m, k)$ therefore has $2n$ vertices v_i, v'_i and $3n$ edges (v_i, v'_{i+m}) , (v_i, v'_{i-m-k}) , (v'_i, v'_{i+k}) ($0 \leq i \leq n - 1$, subscripts mod n).

The following information can be found in [33],[39, Chapter 9]. Suppose that \mathcal{P} is a presentation such that its presentation complex $K_{\mathcal{P}}$ is the spine of a closed 3-manifold M . Then there is a 3-complex C with face pairing such that M is obtained from C by identifying the faces. Moreover, the resulting cell structure on M has a single vertex, a single 3-cell, and 2-skeleton homeomorphic to $K_{\mathcal{P}}$. The Whitehead graph Γ is the link of the single vertex of $K_{\mathcal{P}}$ and so embeds in the link of the single vertex of M . The link of the vertex of M is a 2-sphere and so Γ has a planar embedding on this sphere (see also [22, page 33]). In the case when Γ is connected the 3-complex C is a polyhedron π bounding a 3-ball. Given a planar embedding of Γ there is a one-one correspondence between the faces F of this embedding and the vertices u_F of π . This correspondence maps a face F of degree d of Γ to a vertex of degree d of π . Moreover, if the vertices in the face F read cyclically around the face are w_0, \dots, w_d where $w_i \in \{v_i, v'_i\}$ then the directed edges e_0, \dots, e_d incident to u_F , read cyclically are labelled x_0, \dots, x_d and directed towards u_F if $w_i = v_i$ and away from u_F if $w_i = v'_i$. The faces of π correspond to the relators of the presentation \mathcal{P} : for each relator r there are two faces whose boundaries, read either clockwise or anticlockwise, spell r .

The groups defined by the presentations considered in Lemmas 6,7,8 have finite abelianisation of odd order. In this situation it follows from [36, Lemma 1] that if the presentation complex is the spine of a 3-manifold then it is the spine of a

closed 3-manifold. In these lemmas we either show that the Whitehead graph is non-planar or we show that the Whitehead graph is connected and has a unique planar embedding and that this planar embedding can be used to show that the polyhedron π does not exist. Either way, the presentation complex is not a 3-manifold spine.

Lemma 6. *The presentations $\mathcal{H}(9, 4)$, $\mathcal{H}(9, 7)$ are not 3-manifold spines.*

Proof. We argue that for $m = 4$ or 7 , the Whitehead graph Γ of $\mathcal{H}(9, m)$ is non-planar. We have that Γ is the graph with vertices $v_0, \dots, v_8, v'_0, \dots, v'_8$ and edges $(v_i, v_{i+3}), (v'_i, v'_{i+1}), (v_i, v'_{i+m})$ ($0 \leq i \leq 8$, subscripts mod 9). Removing the edges $(v'_0, v'_1), (v'_3, v'_4), (v'_6, v'_7)$ then contracting the edges (v_i, v_{i+3}) ($0 \leq i \leq 8$), (v'_i, v'_{i+1}) ($i = 1, 2, 4, 5, 7, 8$) and removing loops leaves the non-planar graph $K_{3,3}$ so Γ is non-planar. \square

Lemma 7. *If $n \geq 3$ then the presentation $\mathcal{H}(n, 0) = \mathcal{G}_n(x_0^2 x_1^{-1})$ is not a 3-manifold spine.*

Proof. The Whitehead graph for $\mathcal{H}(n, 0)$ has vertices $v_0, \dots, v_{n-1}, v'_0, \dots, v'_{n-1}$ and edges $v_i - v'_i, v_i - v_{i+1}, v'_i - v'_{i+1}$ for each $0 \leq i \leq n-1$ so there is a unique planar embedding with two n -gons, $v_0 - v_1 - \dots - v_{n-1} - v_0$ and $v'_0 - v'_1 - \dots - v'_{n-1} - v'_0$, and n 4-gons $v_i - v_{i+1} - v'_{i+1} - v'_i - v_i$ ($0 \leq i \leq n-1$). If $\mathcal{H}(n, 0)$ is a 3-manifold spine then in the face pairing polyhedron π there is a degree 4 vertex u corresponding to the 4-gon $v_0 - v_1 - v'_1 - v'_0 - v_0$. The 2-cells incident to u correspond to the relators $x_0^2 x_1^{-1}, x_1^2 x_2^{-1}, x_2^2 x_1^{-1}, x_0^2 x_1^{-1}$ and so there are more than two 2-cells for the relator $x_0^2 x_1^{-1}$, a contradiction. \square

Lemma 8. *Suppose $n \geq 1$, $(m, k) = 1$, $m \neq k$, $k \neq 0$ and either $2k \equiv 0 \pmod n$ or $2(m-k) \equiv 0 \pmod n$. Then $\mathcal{G}_n(m, k)$ is not a 3-manifold spine.*

Proof. If $2(m-k) \equiv 0 \pmod n$ then $\mathcal{G}_n(m, k)$ has relators $x_i x_{i+m} x_{i+m+n/2}^{-1}$. Inverting these and replacing each x_i by its inverse, then cyclically permuting gives the relators $x_{i+m} x_i x_{i+m+n/2}^{-1}$. Subtracting m from the subscripts gives $x_i x_{i-m} x_{i+n/2}^{-1}$; then negating the subscripts and setting $j = -i$ gives $x_j x_{j+m} x_{j+n/2}^{-1}$ which are the relators for the case $2k \equiv 0 \pmod n$. Thus, without loss of generality, we may assume $k = n/2$ and $(m, n/2) = 1$. Then the Whitehead graph Γ has vertices $v_0, \dots, v_{n-1}, v'_0, \dots, v'_{n-1}$ and edges $v_i - v_{i+m+n/2}, v'_i - v'_{i+n/2}, v_i - v'_{i+m}$ ($0 \leq i \leq n-1$). The hypothesis $(m, k) = 1$ implies that $(m + n/2, n) = 1$ or 2 .

Suppose that $(m + n/2, n) = 1$. Then there exist integers α, β such that $\alpha(m + n/2) + \beta n = 1$, so $\alpha(m + n/2) \equiv 1 \pmod n$ and α is odd. For each i ($0 \leq i \leq n-1$) set $w_i = v_{(m+n/2)i}, w'_i = v'_{(m+n/2)i}$ so $v_i = w_{\alpha i}$ and $v'_i = w'_{\alpha i}$. For each i , writing $j = \alpha i \pmod n$ gives that Γ has vertices w_j, w'_j ($0 \leq j \leq n-1$) and edges $(w_j, w_{j+1}), (w'_j, w'_{j+n/2}), (w_j, w'_{j+\alpha m})$. Contracting the edges $(w_j, w'_{j+\alpha m})$ gives the graph with vertices w_j ($0 \leq j \leq n-1$) and edges $(w_j, w_{j+1}), (w_j, w_{j+n/2})$, which is non-planar for $n \geq 6$. Hence Γ is non-planar and so $\mathcal{G}_n(m, k)$ is not a 3-manifold spine.

Suppose then that $(m + n/2, n) = 2$. Then there is a planar embedding of Γ with two $n/2$ -gons $(v_0 - v_{m+n/2} - v_{2m} - \dots - v_{-2m} - v_{-m+n/2} - v_0)$ and $(v_{n/2} - v_m -$

$v_{2m+n/2} \cdots v_{-2m+n/2} v_{-m} v_{n/2}$), $n/2$ 2-gons ($v'_i - v'_{i+n/2} - v'_i$, $0 \leq i \leq n/2 - 1$), and $n/2$ octagons ($v_i - v_{i+m+n/2} - v'_{i+2m+n/2} - v'_{i+2m} - v_{i+m} - v_{i+n/2} - v'_{i+m+n/2} - v'_{i+m} - v_i$, $0 \leq i \leq n/2 - 1$). Moreover, this planar embedding is unique up to self homeomorphism of S^2 . If $\mathcal{G}_n(m, k)$ is a 3-manifold spine then in the face pairing polyhedron π there are two sink vertices of degree $n/2$ and $n/2$ source vertices of degree 2, and the remaining $n/2$ vertices are of degree 8 and are neither sources nor sinks.

At the i 'th source vertex, u_i , the two outgoing edges have labels x_i and $x_{i+n/2}$. The union of the two incident 2-cells is a 2-gon C_i whose boundary label is the positive word $x_{i+m} x_{i+m+n/2}$. In particular, neither of the vertices of C_i is a sink or a source vertex. The incoming edges at the two sink vertices, $v_{\text{even}}, v_{\text{odd}}$, are, in cyclic order $x_0, x_{m+n/2}, x_{2m}, \dots, x_{-m+n/2}$ and $x_1, x_{1+m+n/2}, x_{1+2m}, \dots, x_{1-m+n/2}$, respectively. The union of the $n/2$ 2-cells incident at v_{even} (resp. v_{odd}) is an $n/2$ -gon D_{even} (resp. D_{odd}) with boundary label $W_{\text{even}} = x_m x_{2m+n/2} x_{3m} \cdots x_{-m} x_{n/2}$ (resp. $W_{\text{odd}} = x_{1+m} x_{1+2m+n/2} x_{1+3m} \cdots x_{1-m} x_{1+n/2}$). Now $D_{\text{even}} \cup D_{\text{odd}}$ contains precisely one 2-cell corresponding to each of the relators, as does $\bigcup_i C_i$, so the 2-sphere $\partial\pi$ must be formed from $D_{\text{even}}, D_{\text{odd}}$ and the C_i ($0 \leq i \leq n/2 - 1$) by identifying similarly labelled 1-cells in their boundaries. However, identifying the 1-cells of ∂D_{even} with the corresponding 1-cells of the ∂C_i makes $D_{\text{even}} \cup (\bigcup_i C_i)$ into an $n/2$ -gon whose boundary label is the *reverse* of W_{odd} , that is $x_{1+n/2} x_{1-m} \cdots x_{1+3m} x_{1+2m+n/2} x_{1+m}$. Therefore it is not possible to form the 2-sphere $\partial\pi$ from this $n/2$ -gon and D_{odd} by identifying the 1-cells in their boundaries, and hence $\mathcal{G}_n(m, k)$ is not a 3-manifold spine. \square

Lemma 9. *Let $n \geq 1$. Then the presentation $\mathcal{H}(n, 1) = \mathcal{G}_n(x_0 x_1 x_1^{-1})$ is a spine of S^3 .*

Proof. If $n = 1$ then $\mathcal{H} = \mathcal{H}(n, 1) = \langle x_0 \mid x_0 x_0 x_0^{-1} \rangle$ which is a spine of S^3 by [46], so assume $n \geq 2$. Consider the polyhedron in Figure 1 with face pairings as shown. We shall show that the identification of faces results in a 3-complex M whose 2-skeleton is the presentation complex of \mathcal{H} , so M has one 0-cell, n 1-cells, n 2-cells and one 3-cell, so is a manifold by [39, Theorem I, Section 60]. To this end note that all edges labelled x_1 , namely $[S, S]_1, [N, S]_1, [S, v_1]$ are identified by the following cycle of faces identifications:

$$[S, v_1] \xleftarrow{F_0} [S, S]_1 \xleftarrow{F_1} [N, S]_1 \xleftarrow{F_0} [S, v_1].$$

By symmetry, for each $0 \leq i \leq n - 1$, all edges labelled x_i , namely $[S, S]_i, [N, S]_i, [S, v_i]$ are similarly identified. The initial vertices of the edges labelled x_i , namely N, S are identified and the terminal vertices of the edges labelled x_i , namely S, v_i are identified. Therefore the resulting complex has one 0-cell, n 1-cells, n 2-cells, and one 3-cell, and so its 2-skeleton is the presentation complex of \mathcal{H} , and the resulting manifold is S^3 . \square

Proof of Theorem B. (a) By Theorem A we may assume that $(m, k) = 1$ and either $2k \equiv 0 \pmod n$ or $2(m - k) \equiv 0 \pmod n$, in which case $\mathcal{G}_n(m, k)$ is not a 3-manifold spine by Lemma 8.




Figure 1: Face pairing polyhedron for the presentation $\mathcal{H}(n, 1) = \mathcal{G}_n(x_0 x_1 x_1^{-1})$.

(b) If $(n, m) = (9, 4)$ or $(9, 7)$ then $\mathcal{H}(n, m)$ is not a 3-manifold spine by Lemma 6, so assume $(n, m) \neq (9, 4), (9, 7)$. By Theorem A we may assume that $m = 0$ or $m = 1$ or $m = 2$ or $(m = n - 1$ and $n = 3, 5, 7$ or $n \geq 4$ is even) or $(n, m) = (5, 3)$ or $(7, 4)$ or $m = n/2 + 1$ (where $n \geq 4$ is even). In the case $m = 0$ the presentation $\mathcal{H}(n, m)$ is not a 3-manifold spine by Lemma 7. In the cases $m = 1$ and $m = 2$ the presentation $\mathcal{H}(n, m)$ is a 3-manifold spine by Lemma 9 and Theorem 2, respectively. In the case $m = n - 1$ we have that $\mathcal{H}(n, m)$ is homeomorphic to $\mathcal{F}(2, n)$ so when $n \geq 4$ is even then it is a 3-manifold spine by Theorem 1 and if $n = 3, 5$ or 7 then it is a 3-manifold spine if and only if $n = 3$ by [9, page 55]. In the cases $(n, m) = (5, 3)$ or $(7, 4)$ we have that $\mathcal{H}(n, m)$ is homeomorphic to $\mathcal{F}(2, 5)$ or $\mathcal{F}(2, 7)$, so again is not a 3-manifold spine. In the case $m = n/2 + 1$ we have that the presentation $\mathcal{H}(n, m)$ is not a 3-manifold spine by Lemma 8. \square

4 Fibonacci and Sieradski manifolds revisited

In [28, Example 1.2] the presentation $\mathcal{G}_m(x_0^{-1} x_1^2 x_2^{-1} x_1)$ of the Fibonacci group $F(2, 2m)$ was obtained. In the same way we can obtain an alternative presentation of the Sieradski group $S(2, 2m)$:

$$\begin{aligned}
 S(2, 2m) &= \langle x_i \mid x_i x_{i+2} = x_{i+1} \ (0 \leq i \leq 2m - 1) \rangle \\
 &= \langle x_{2j}, x_{2j+1} \mid x_{2j} x_{2j+2} = x_{2j+1}, x_{2j+1} x_{2j+3} = x_{2j+2} \ (0 \leq j \leq m - 1) \rangle \\
 &= \langle x_{2j} \mid (x_{2j} x_{2j+2})(x_{2j+2} x_{2j+4}) = x_{2j+2} \ (0 \leq j \leq m - 1) \rangle \\
 &= \langle y_j \mid (y_j y_{j+1})(y_{j+1} y_{j+2}) = y_{j+1} \ (0 \leq j \leq m - 1) \rangle \\
 &= \mathcal{G}_m(y_0 y_1^2 y_2 y_1^{-1}).
 \end{aligned}$$

Theorem C. For each $m \geq 3$ the cyclic presentation $\mathcal{G}_m(x_0^{-1}x_1^2x_2^{-1}x_1)$ of the Fibonacci group $F(2, 2m)$ and the cyclic presentation $\mathcal{G}_m(x_0x_1^2x_2x_1^{-1})$ of the Sieradski group $S(2, 2m)$ are spines of closed 3-manifolds.

Proof of Theorem C. Consider first the presentation $\mathcal{G} = \mathcal{G}_m(x_0^{-1}x_1^2x_2^{-1}x_1)$ of the Fibonacci group $F(2, 2m)$ and consider the polyhedron in Figure 2 with face pairings as shown. We shall show that the identification of faces results in a 3-complex M whose 2-skeleton is the presentation complex of \mathcal{G} , so M has one 0-cell, m 1-cells, m 2-cells and one 3-cell, so is a manifold by [39, Theorem I, Section 60]. To this end note that all edges labelled x_2 are identified by the following cyclic of faces identifications:

$$[N, v_1] \xleftarrow{F_1} [u_2, S] \xleftarrow{F_2} [w_2, v_2] \xleftarrow{F_1} [w_1, w_2] \xleftarrow{F_1} [u_1, w_1] \xleftarrow{F_0} [N, v_1].$$

By symmetry, for each $0 \leq i \leq m-1$, all edges labelled x_i are similarly identified. The initial vertices of the edges labelled x_i , namely $u_i, u_{i-1}, w_i, w_{i-1}, N$ are identified and the terminal vertices of the edges labelled x_i , namely $v_i, v_{i-1}, w_i, w_{i-1}, S$ are identified. Therefore the resulting complex has one 0-cell, m 1-cells, m 2-cells, and one 3-cell, and so its 2-skeleton is the presentation complex of \mathcal{G} , as required.

Consider now the presentation $\mathcal{G}_n(x_0x_1^2x_2x_1^{-1})$ of the Sieradski group $S(2, 2m)$ and consider the polyhedron in Figure 3. We argue as above. The edges labelled x_3 are identified by the following cycle of face identifications:

$$[N, u_4] \xleftarrow{F_2} [v_2, S] \xleftarrow{F_1} [w_2, u_3] \xleftarrow{F_2} [u_3, v_3] \xleftarrow{F_2} [v_3, w_3] \xleftarrow{F_3} [N, u_4].$$

By symmetry, for each $0 \leq i \leq m-1$, all edges labelled x_i are similarly identified. The initial vertices of the edges labelled x_i , namely $v_{i-1}, w_{i-1}, u_i, v_i, N$ are identified and the terminal vertices of the edges labelled x_i , namely $u_{i+1}, u_i, v_i, w_i, S$ are identified. \square

Observe that the polyhedron given in Figure 2 for the presentation $\mathcal{G}_m(x_0^{-1}x_1^2x_2^{-1}x_1)$ of $F(2, 2m)$ appears to be of a fundamentally different nature to the polyhedron for the presentation $\mathcal{F}(2, 2m) = \mathcal{G}_m(x_0x_1x_2^{-1})$ given in [18, Figure 1]: the former consists of $2m$ pentagons whereas the latter consists of $4m$ triangles. Similarly, the polyhedron given in Figure 3 for the presentation $\mathcal{G}_n(x_0x_1^2x_2x_1^{-1})$ of $S(2, 2m)$ appears to be of a fundamentally different nature to the polyhedron for the presentation $\mathcal{S}(2, 2m) = \mathcal{G}_{2m}(x_0x_2x_1^{-1})$ given in [40, Figure 8] or [4, Figure 1]: again, the former consists of $2m$ pentagons, while the latter consists of $4m$ triangles.

The alternative Fibonacci presentation $\mathcal{G}_n(x_0^{-1}x_1^2x_2^{-1}x_1)$ is already known to be a 3-manifold spine, as it appears in line 6 of [11, Table 1]; however, our proof gives an explicit construction of the face-pairing polyhedron. As observed in [28, Example 1.2] the polynomial $1 + t - t^2$ associated to the original Fibonacci presentation $\mathcal{G}_{2m}(x_0x_1x_2^{-1})$ is not an Alexander polynomial, but the polynomial $1 - 3t + t^2$ associated to the alternative Fibonacci presentation $\mathcal{G}_m(x_0^{-1}x_1^2x_2^{-1}x_1)$ is the Alexander polynomial of the figure eight knot. For the Sieradski presentations, the polynomial $1 - t + t^2$ for the presentation $\mathcal{G}_{2m}(x_0x_2x_1^{-1})$ is the Alexander polynomial of the trefoil knot, but the polynomial $1 + t + t^2$ for the presentation $\mathcal{G}_m(x_0x_1^2x_2x_1^{-1})$ is not an Alexander polynomial [2].




Figure 2: Face pairing polyhedron for the presentation $\mathcal{G}_m(x_0^{-1}x_1^2x_2^{-1}x_1)$ of the Fibonacci group $F(2, 2m)$.




Figure 3: Face pairing polyhedron for the presentation $\mathcal{G}_n(x_0x_1^2x_2x_1^{-1})$ of the Sieradski group $S(2, 2m)$.

Acknowledgements

We thank Bill Bogley for helpful comments on a draft of this article and the referee(s).

References

- [1] V.G. Bardakov and A.Yu. Vesnin. A generalization of Fibonacci groups. *Alg. and Logic*, 42:131–160, 2003.
- [2] Gerhard Burde and Heiner Zieschang. *Knots*. Berlin: Walter de Gruyter, 2nd revised and extended edition, 2003.
- [3] J.W. Cannon, W.J. Floyd, L. Lambert, W.R. Parry, and J.S. Purcell. Bi-twist manifolds and two-bridge knots. *Pacific J. Math.*, 284(1):1–39, 2016.
- [4] A. Cavicchioli, F. Hegenbarth, and A.C. Kim. A geometric study of Sieradski groups. *Algebra Colloq.*, 5(2):203–217, 1998.
- [5] Alberto Cavicchioli, Friedrich Hegenbarth, and Dušan Repovš. On manifold spines and cyclic presentations of groups. In *Knot theory*, volume 42, pages 49–56. Warszawa: Polish Academy of Sciences, Institute of Mathematics, Banach Cent. Publ., 1998.
- [6] Alberto Cavicchioli, Dušan Repovš, and Fulvia Spaggiari. Topological properties of cyclically presented groups. *J. Knot Theory Ramifications* 12(2):243–268, 2003.
- [7] Alberto Cavicchioli, E.A. O’Brien, and Fulvia Spaggiari. On some questions about a family of cyclically presented groups. *J. Algebra*, 320(11):4063–4072, 2008.
- [8] Alberto Cavicchioli, Dušan Repovš, and Fulvia Spaggiari. Families of group presentations related to topology. *J. Algebra*, 286(1):41–56, 2005.
- [9] Alberto Cavicchioli and Fulvia Spaggiari. The classification of 3-manifolds with spines related to Fibonacci groups. In *Algebraic Topology, Proc. Conf., S. Feliu de Guíxols/Spain 1990*, Lect. Notes Math., pages 50–78, 1992.
- [10] Christopher P. Chalk. Fibonacci groups with aspherical presentations. *Commun. Algebra*, 26(5):1511–1546, 1998.
- [11] M.J. Dunwoody. Cyclic presentations and 3-manifolds. In *Groups - Korea '94. Proceedings of the international conference, Pusan, Korea, August 18-25, 1994*. Berlin: Walter de Gruyter, 1995.
- [12] Martin Edjvet and Gerald Williams. The cyclically presented groups with relators $x_i x_{i+k} x_{i+l}$. *Groups, Geom., Dyn.*, 4(4):759–775, 2010.

- [13] D. B. A. Epstein. Projective planes in 3-manifolds. *Proc. London Math. Soc.* (3), 11:469–484, 1961.
- [14] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.7.9*, 2015.
- [15] N.D. Gilbert and J. Howie. LOG groups and cyclically presented groups. *J. Algebra*, 174(1):118–131, 1995.
- [16] Jens Harlander. Some aspects of efficiency. Baik, Young Gheel (ed.) et al., Groups - Korea '98. Proceedings of the 4th international conference, Pusan, Korea, August 10-16, 1998. Berlin: Walter de Gruyter. 165-180, 2000.
- [17] W. Heil. On Kneser's conjecture for bounded 3-manifolds. *Proc. Camb. Philos. Soc.*, 71:243–246, 1972.
- [18] H. Helling, A.C. Kim, and J.L. Mennicke. A geometric study of Fibonacci groups. *J. Lie Theory*, 8(1):1–23, 1998.
- [19] John Hempel. *3-manifolds.*, volume 86 of *Annals of Mathematics Studies*. Princeton, New Jersey: Princeton University Press and University of Tokyo Press, 1976.
- [20] Hugh Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. On a remarkable polyhedron geometrizing the figure eight knot cone manifolds. *J. Math. Sci., Tokyo*, 2(3):501–561, 1995.
- [21] Hugh M. Hilden, María Teresa Lozano, and Jose María Montesinos-Amilibia. The arithmeticity of the figure eight knot orbifolds. In *Topology '90. Contributions from a research semester in low dimensional topology, held at Ohio State University, Columbus, OH (USA), from Feb. through June 1990*, pages 169–183. Berlin: Walter de Gruyter, 1992.
- [22] Cynthia Hog-Angeloni and Wolfgang Metzler. Geometric aspects of two-dimensional complexes. In Cynthia Hog-Angeloni et al., editor, *Two-dimensional homotopy and combinatorial group theory*, number 197 in Lond. Math. Soc. Lect. Note Ser., pages 1–50. Cambridge: Cambridge University Press, 1993.
- [23] Derek F. Holt. An alternative proof that the Fibonacci group $F(2,9)$ is infinite. *Exp. Math.*, 4(2):97–100, 1995.
- [24] James Howie and Gerald Williams. Tadpole Labelled Oriented Graph groups and cyclically presented groups. *J. Algebra*, 371:521–535, 2012.
- [25] D.L. Johnson. *Topics in the theory of group presentations*, volume 42 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 1980.
- [26] D.L. Johnson and H. Mawdesley. Some groups of Fibonacci type. *J. Aust. Math. Soc.*, 20:199–204, 1975.

- [27] Soo Hwan Kim and Yangkok Kim. On the generalized Dunwoody 3-manifolds. *Osaka J. Math.*, 50(2):457-476, 2013.
- [28] Ann Chi Kim and Andrei Vesnin. Cyclically presented groups and Takahashi manifolds. *RIMS Kokyuroku*, 1022:200–212, 1997.
- [29] G.W. Knutson. A characterization of closed 3-manifolds with spines containing no wild arcs. *Proc. Am. Math. Soc.*, 21:310–314, 1969.
- [30] Wolfgang Lück. Survey on aspherical manifolds. In *European congress of mathematics. Proceedings of the 5th ECM congress, Amsterdam, Netherlands, July 14–18, 2008*, pages 53–82. Zürich: European Mathematical Society (EMS), 2010.
- [31] Colin Maclachlan. Generalisations of Fibonacci numbers, groups and manifolds. In *Combinatorial and geometric group theory. Proceedings of a workshop held at Heriot-Watt University, Edinburgh, spring of 1993*. Lond. Math. Soc. Lecture Note Series Vol. 204, pages 233–238 (1995).
- [32] Sergei Matveev. *Algorithmic topology and classification of 3-manifolds*. Berlin: Springer, 2nd edition, 2007.
- [33] L. Neuwirth. An algorithm for the construction of 3-manifolds from 2-complexes. *Proc. Camb. Philos. Soc.*, 64:603–613, 1968.
- [34] M.F. Newman. Proving a group infinite. *Arch. Math.*, 54(3):209–211, 1990.
- [35] Peter Orlik. *Seifert manifolds*. Lecture Notes in Mathematics, Vol. 291. Springer-Verlag, Berlin-New York, 1972.
- [36] R.P. Osborne and R.S. Stevens. Group presentations corresponding to spines of 3-manifolds. III. *Trans. Am. Math. Soc.*, 234:245–251, 1977.
- [37] John G. Ratcliffe. Euler characteristics of 3-manifold groups and discrete subgroups of $SL(2, \mathbb{C})$. *J. Pure and Appl. Algebra*, 44:303–314, 1987.
- [38] G.P. Scott. Compact submanifolds of 3-manifolds. *J. Lond. Math. Soc., II. Ser.*, 7:246–250, 1973.
- [39] H. Seifert and W. Threlfall. *A textbook of topology*, volume 89 of *Pure and Applied Mathematics*. Academic Press, New York, 1980.
- [40] Allan J. Sieradski. Combinatorial squashings, 3-manifolds, and the third homology of groups. *Invent. Math.*, 84:121–139, 1986.
- [41] Andrzej Szczepański and Andrei Vesnin. On generalized Fibonacci groups with an odd number of generators. *Commun. Algebra*, 28(2):959-965 (2000).
- [42] Richard M. Thomas. The Fibonacci groups revisited. Groups, Vol. 2, Proc. Int. Conf., St. Andrews/UK 1989, Lond. Math. Soc. Lect. Note Ser. 160, 445-454, 1991.

- [43] Gerald Williams. The aspherical Cavicchioli-Hegenbarth-Repovš generalized Fibonacci groups. *J. Group Theory*, 12(1):139–149, 2009.
- [44] Gerald Williams. Unimodular integer circulants associated with trinomials. *Int. J. Number Theory*, 6(4):869–876, 2010.
- [45] Gerald Williams. Groups of Fibonacci type revisited. *Int. J. Algebra Comput.*, 22(8):19, 2012.
- [46] E.C. Zeeman. On the dunce hat. *Topology*, 2:341–358, 1963.

DEPARTMENT OF MATHEMATICS AND MAXWELL INSTITUTE FOR MATHEMATICAL SCIENCES, HERIOT-WATT UNIVERSITY, EDINBURGH EH14 4AS, UK.
E-mail address, J.Howie@hw.ac.uk

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ESSEX, WIVENHOE PARK, COLCHESTER, ESSEX CO4 3SQ, UK.
E-mail address, Gerald.Williams@essex.ac.uk