



Heriot-Watt University
Research Gateway

Coherent Perfect Absorption in Metamaterials with Entangled Photons

Citation for published version:

Altuzarra, C, Vezzoli, S, Valente, J, Gao, W, Soci, C, Faccio, DFA & Couteau, C 2017, 'Coherent Perfect Absorption in Metamaterials with Entangled Photons', *ACS Photonics*, vol. 4, no. 9, pp. 2124-2128.
<https://doi.org/10.1021/acsp Photonics.7b00514>

Digital Object Identifier (DOI):

[10.1021/acsp Photonics.7b00514](https://doi.org/10.1021/acsp Photonics.7b00514)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Peer reviewed version

Published In:

ACS Photonics

Publisher Rights Statement:

This document is the Accepted Manuscript version of a Published Work that appeared in final form in ACS Photonics, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <http://pubs.acs.org/doi/abs/10.1021/acsp Photonics.7b00514>

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Coherent Perfect Absorption in Metamaterials with Entangled Photons

Charles Michael Xavier Altuzarra, Stefano Vezzoli, Joao Valente,
WEIBO GAO, Cesare Soci, Daniele Faccio, and Christophe Couteau


ACS Photonics, **Just Accepted Manuscript** • DOI: 10.1021/acsp Photonics.7b00514 • Publication Date (Web): 08 Aug 2017

Downloaded from <http://pubs.acs.org> on August 18, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



(Table of Content Graphic)

125x63mm (96 x 96 DPI)

Coherent Perfect Absorption in Metamaterials with Entangled Photons

Charles Altuzarra^{1,2}, Stefano Vezzoli^{1,4}, Joao Valente³, Weibo Gao¹, Cesare Soci¹, Daniele Faccio⁴, Christophe Couteau^{1,2,5,*}

¹ Centre for Disruptive Photonic Technologies, TPI, Nanyang Technological University, 637371, Singapore

² CINTRA, CNRS-NTU-Thales, CNRS UMI 3288, Singapore

³ Optoelectronics Research Centre and Centre for Photonic Metamaterials, University of Southampton, Southampton, UK

⁴ Institute for Photonics and Quantum Sciences and SUPA, Heriot-Watt University, Edinburgh EH14 4AS, UK

⁵ Laboratory for Nanotechnology, Instrumentation and Optics, ICD CNRS UMR 6281, University of Technology of Troyes, Troyes, France

* Corresponding author: christophe.couteau@utt.fr

(Dated: August 3, 2017)

Quantum nonlocality, i.e. the presence of strong correlations in spatially separated systems which are forbidden by local realism, lies at the heart of quantum communications and quantum computing. Here, we use polarization-entangled photon pairs to demonstrate a nonlocal interaction of light with a plasmonic structure. Through the detection of one photon with a polarization-sensitive device, we can prevent or allow absorption of a second, remotely located photon. We demonstrate this with pairs of entangled photons in polarization, one of which is coupled into a plasmon of a thin metamaterial absorber in the path of a standing wave of an interferometer. Thus, we realize a quantum eraser experiment using photons and plasmonic resonances from metamaterials which promises opportunities for probabilistic quantum gating and controlling plasmon-photon conversion and entanglement. Moreover, by using the so-called coherent perfect absorption effect, we can expect near perfect interaction.

Keywords: Plasmonic metamaterials, Coherent Perfect Absorption, Quantum interferometry, Quantum optics

One of the quintessential aspects of quantum mechanics is the existence of entangled states whereby the classical description of a particle being in a well-defined state is replaced with a quantum description based on superposition of states. Moreover, quantum entanglement provides a unique route for nonlocal correlations between remote particles such as photons. Beyond the relevance to the fundamental questions of quantum physics^{1–4}, nonlocality is a resource for a number of applications such as quantum teleportation, quantum erasure, and interaction-free measurements^{5–11}. In addition to demonstrating a new application of nonlocality, our work presented here is also relevant to the field of quantum plasmonics. This rather recent field emerged in 2002 when Alteuischer *et al.*¹² showed that light passing through a metallic nanohole array conserved the quantum state of entangled photons. In parallel, Lukin and co-workers demonstrated that single plasmons can be generated from single photons as a process that can be deterministic^{13,14}. More recently, Hong-Ou-Mandel two-photon quantum interference with plasmons was theoretically studied¹⁵ and experiments were carried out^{16–18}, thereby providing experimental proof that propagating plasmons retained the quantum coherency of the photons that launched them.

In this work, we demonstrate that entangled photons can be used to achieve a remote interaction of photons, in the form of a quantum eraser, with the excitation of plasmonic modes resulting in coherent perfect absorption of light. In order to do so with polarization-entangled

photons, we constructed a polarization-sensitive ‘quantum eraser’ interferometer for which we show that the conditions for interference can be non-locally modified through a polarization-sensitive detection of the entangled photons. We previously demonstrated that the level of coupling to plasmons by using a thin metamaterial placed in a single photon interferometer can be varied from nearly 0% (‘perfect transmission’) to 100% (‘perfect absorption’), depending on the position in the standing wave, whereas only 50% absorption is observed, when the standing wave is not formed^{19,20}.

The concept.— In a simplified Sagnac interferometer, an input beamsplitter creates two optical paths A and B (see Fig.1). To render the device polarization-sensitive we introduce a half-wave plate only in path A and orient its main axis to be 45° to the plane of the interferometer²¹. In what follows we will refer to light linearly polarized in the plane of the interferometer as horizontally polarized (H-polarized) light and light polarized perpendicular to the plane as vertically polarized (V-polarized) light. Such an interferometer creates standing waves only for light polarized along the fast and slow axis of the waveplate. Input linear polarizations of +45° or –45° to the plane of the interferometer will not be affected by the wave-plate and will evolve through both paths A and B as identically polarized traveling waves forming two standing waves in the interferometer with the antinode for +45° corresponding to the node for –45°. Conversely, the half-wave plate converts a vertical polarization into a horizontal one, and vice versa:

so if vertically or horizontally polarized light is launched in the interferometer, optical paths A and B will contain counter-propagating orthogonal waves that do not interfere: a standing wave is not formed in the interferometer.

Now, with polarization-entangled photon pairs (denoted idler i and signal s photons for each pair, see Fig.1) it is possible to nonlocally select the state of the signal photons inside the interferometer, through a measurement on the idler photons outside (and that never enter) the interferometer. This is achieved by adding a polarizer on the idler channel. When the idler polarizer is set to either an angle of $+45^\circ$ or -45° to the plane of the interferometer, the polarization state of the polarization-entangled signal photon will be polarized at -45° or $+45^\circ$ correspondingly. The signal photons path-entangled wavefunction will form a standing wave in the interferometer and strong dissipation in the 'coherent absorption' regime and zero dissipation in the 'coherent transmission' regime can be observed²². Conversely, when the polarizer in the idler channel is set vertically or horizontally, the polarization state of the signal photon will be necessarily projected to the horizontal or vertical polarization, correspondingly. Therefore, due to distinguishability of optical paths^{7,8}, no standing wave will be formed in the interferometer and the coherent absorption process is no longer visible.

Therefore, the scheme described here is a dissipative form of a quantum eraser: the interferometer polarizer rotated to the horizontal or vertical polarizations introduces a "which-path" information and hence removes any standing-wave effects at the metamaterial. By selecting idler photons (i.e., outside the interferometer) at 45° , the "which-path" information is nonlocally erased and coherent control of absorption is observed at the metamaterial.

The experiment.— The experimental setup is shown in Fig.1. We generated pairs of polarization-entangled photons at the wavelength of 810 nm by spontaneous parametric down-conversion (SPDC). A 200-mW laser diode with emission centered at the wavelength of $\lambda_p = 405$ nm was used to pump a 2mm-thick type-II beta-barium borate (BBO) nonlinear crystal producing non-collinear, degenerate photon pairs. Polarization entanglement is achieved by adding 1mm-thick BBO compensation crystals and half wave plates set at 45° . The photon pairs collected from the areas of intersections of phase-matching cones were coupled to single-mode fibres with collimation lenses. A 10-nm bandpass filter centered at 810 nm was used to block the pump radiation and select 'twin' SPDC photons. As detailed in Fig.1, the idler channel was connected to a photon counting avalanche photodiode detector and a coincidence counter (IDQuantique ID800). It was used to control the presence of the signal photon within the interferometer. The signal photons were coupled to the interferometer. A variable retarder was used to compensate for the polarization change in the signal fiber. The photons enter the interferometer through a lossless (50 : 50) non-polarizing beam splitter. The thin metamaterial absorber was placed at the centre

of the interferometer between two x10 microscope objectives producing a spot size of $\approx 10 \mu m$ in diameter. The absorber's position was scanned using a piezoelectrically actuated linear translation stage over a few optical wavelengths. The sum of the photon counts were detected by the two avalanche detectors (PD_A and PD_B) in coincidence with the idler photon PD_i within a 10 ns time window.

The SPDC source creates a quantum superposition of polarized photons of orthogonal basis. The wavefunctions general form for polarization entangled states of this type is²³:

$$|\Psi\rangle = \frac{1}{\sqrt{2}} (|H\rangle_i |V\rangle_s - |V\rangle_i |H\rangle_s) \quad (1)$$

where indices i and s denote the idler and signal photons respectively and $|H\rangle$ and $|V\rangle$ denote the horizontal and vertical polarization states respectively.

The path entanglement wavefunction of a single photon that enables interference has the general form:

$$|\Psi\rangle = \frac{1}{\sqrt{2}} (|1\rangle_A |0\rangle_B - e^{i\phi} |0\rangle_A |1\rangle_B) \quad (2)$$

By integrating the path entanglement wavefunction (Eq.2) into the polarization entanglement wavefunction (Eq.1), in representation of our optical scheme, we obtain:

$$|\Psi\rangle = \frac{1}{2} [|H\rangle_i (|H\rangle_A |0\rangle_B - e^{i\phi} |0\rangle_A |V\rangle_B) - |V\rangle_i (|V\rangle_A |0\rangle_B - e^{i\phi} |0\rangle_A |H\rangle_B)] \quad (3)$$

And by expanding, we arrive at the path entanglement of two polarization wavefunctions:

$$|\Psi\rangle = \frac{1}{2} [(|H\rangle_i |H\rangle_A - |V\rangle_i |V\rangle_A) |0\rangle_B - e^{i\phi} |0\rangle_A (|H\rangle_i |V\rangle_B - |V\rangle_i |H\rangle_B)] \quad (4)$$

We first measured the degree of polarization entanglement of the generated photons. These measurements were performed for two different polarization basis sets, $|H, V\rangle$ and $\pm 45^\circ$ which correspond to 1) horizontal and vertical polarizations and 2) polarizations at $+45^\circ$ and -45° to the plane of the interferometer. The Bell parameter was then found to be $S = \sqrt{2}(V_1 + V_2) = 2.66 \pm 0.01$ where $V_{1,2}$ were visibilities calculated from the correlation curves for the two basis sets. Here, according to the Clauser-Horne-Shimony-Holt inequality²³⁻²⁵, a value of S greater than 2 implies nonlocal quantum correlations. We note that our measured value of the Bell parameter of $S = 2.66$, is close to the maximum value of $S = 2\sqrt{2} \approx 2.88$ that is expected for perfectly entangled states.

The plasmonic metamaterial absorber made of split ring resonators was designed to provide a nearly 50% traveling wave absorption, similarly to work reported




FIG. 1. Nonlocal dissipation management with entangled photons. Polarization-entangled photon pairs are generated by a type II spontaneous parametric down conversion process (laser at 405 nm impinging a BBO nonlinear crystal). One photon of the pair is introduced in the interferometer (the signal photon at the 50/50 beamsplitter) and can take two paths (A and B) with a fiber link and a compensator for synchronization issues. Hwp is a half waveplate in arm A. A thin metamaterial plasmonic absorber is introduced in the polarization-sensitive interferometer. By detecting the idler photon outside the interferometer with the help of a polarizer (ξ), one can non-locally prevent or allow deterministic absorption of the signal entangled photon that dissipates through coupling into a plasmon of the absorber placed in the interferometer. PD_A , PD_B and PD_i are photon detectors.

in Ref.22. We fabricated a freestanding gold film of subwavelength thickness nanostructured to create polarization-independent plasmonic absorption at the operational wavelength of 810 nm.

To demonstrate the nonlocal interaction with polarization-entangled photons, we introduce one photon of the entangled pair (signal) in the polarization sensitive interferometer where it interacts with the plasmonic metamaterial absorber. We placed the absorber on a piezo-driven actuator in the center of the interferometer (see Fig.1). We then detected the level of light intensity at the interferometer output by taking the sum of the photon counts registered by photodetectors PD_A and PD_B that are heralded by the detection events of the idler photons on PD_i . We then normalized these to the total level of photon counts when the absorber is removed from the interferometer and recorded the normalized level as a function of absorbers position along the standing wave. The results of these measurements are presented in Fig.2.

No dependence of photon counts jointly registered by photodetectors PD_A and PD_B on the position of the plasmonic metamaterial were seen if the heralding was performed with an idler polarization set (with polarizer ξ in Fig.1) to vertical or horizontal (curves with black diamonds for V and green squares for H in Fig.2-a). This

measurement and distinguishability of which-path information derived from (Eq.1) can be described by:

$$\langle H|\Psi\rangle = \frac{1}{2} (|H\rangle_A|0\rangle_B - e^{i\phi}|0\rangle_A|V\rangle_B) \quad (5)$$

Thus, the single photon wavefunction does not interfere or form a standing wave in the interferometer: coherent control of absorption is lost and photons entering the metamaterial film suffer probabilistic absorption of approximately 50%. A small difference in the level of absorption between curves for vertical and horizontal polarizations is explainable by residual anisotropy of the plasmonic absorber.

On the contrary, if heralding is performed with the idler polarization set to $+45^\circ$ with respect to the plane of the interferometer, we observe a clear oscillation of absorption as a function of the absorber's position in the standing wave (curve with red circles in Fig.2-a). The which-path information has been erased and the path-entangled single photon wavefunction forms a standing wave in the interferometer:

$$\langle +45|\Psi\rangle = \frac{1}{2} (| -45\rangle_A|0\rangle_B - e^{i\phi}|0\rangle_A| -45\rangle_B) \quad (6)$$

The corresponding modulation of the absorption/transmission is shown in Fig.2-a. We note




FIG. 2. Normalized transmission of the plasmonic thin absorber placed in the interferometer, a) registered for different polarization states of the photons detected in the idler channel with vertical polarization (black diamonds), horizontal polarization (green squares) and with polarization at 45° to the plane of interferometer (red circles). b) presents the photons detected in the idler channel with polarization at 45° to the plane of interferometer for two different levels of polarization entanglement of the idler and signal photons: 'strong' entanglement with $S=2.66$ (red circles) and 'weak' entanglement with $S=2.09$ (blue triangles).

that when absorption is achieved for $+45^\circ$, transmission is correspondingly observed for -45° . Each entangled photon entering the interferometer is therefore deterministically absorbed and converted into a plasmon in the nanostructure.

Finally, in order to underline the role of polarization entanglement, we compared the results from experiments performed with two levels of polarization entanglement. We adjusted our photon source from the regime when it generated entangled states close to maximum Bell parameter $S = 2.657 \pm 0.004$, to a rather low degree of entanglement with $S = 2.087 \pm 0.004$. The source of photons with a lower degree of entanglement yielded absorption modulation of approximately 14% (see Fig.2-b, curve with blue triangles) compared to the 80% (see Fig.2-b, curve with

red circles) attainable with strongly entangled photons.

The reduction of the absorption modulation visibility in the experiments performed with photon pairs with high and low values for the Bell parameter is relevant to the fundamental differences in polarization properties of the generated photons. The regime of high Bell parameter implies that photons traveling in the same direction, but emerging from the intersections of two phase-matching cones of the parametric down conversion crystal can create arbitrary superpositions of vertically and horizontally polarized states defined by the nonlinear down-conversion process. As a result entangled, orthogonally polarized pairs of photons are generated with arbitrary polarization basis. In contrast, in the regime of low values of the Bell parameter, quantum superpositions of vertically and horizontally polarized states are not formed and the parametric device generates orthogonally polarized photons where idler and signal photons can either be horizontally or vertically polarized. This mixed state produced is thus not sufficient to produce interferences, unlike an entangled state. Therefore, this underlines a relationship between coherent perfect absorption and quantum states of light. First of all, it is possible to show that the visibility of modulation of absorption scales with the purity p of the quantum state $V_{CPA} = p$, and that the Bell parameter may be described as $S = \sqrt{2}(1 + p)$. These two relations then combine to give the dependence of the modulation of absorption visibility on the Bell parameter $V_{CPA} = \frac{S}{\sqrt{2}} - 1$. For the case of our highly entangled state with $S=2.66$, we therefore predict $V_{CPA} = 0.88$, which is in very good agreement with the experimentally reported value of $V_{CPA} = 0.8$. This slight difference may be ascribed to a minor decoherence effect caused by the metamaterial.

Conclusion.— In conclusion, we have demonstrated a regime of nonlocal interaction of polarization-heralded photons with plasmonic resonances from a metamaterial via the so-called coherent perfect absorption process with a fairly high efficiency. By selecting the idler photon polarization that never entered the interferometer, we can switch the metamaterial from the regime of travelling-wave absorption to the regime of coherent absorption. This quantum eraser experiment with plasmons can be seen as a new type of quantum gate with 'a posteriori knowledge'^{26,27} for which the output signal can be switched nonlocally in the 'coherent transmission' regime from unitary transmission to a probabilistic 50% transmission. Alternatively, in the 'coherent absorption' configuration, the system can be switched nonlocally from zero transmission to a probabilistic 50% transmission^{28,29}.

It should be noted that in our experiment, we do not nonlocally control the total dissipation of light energy on the absorber. Indeed, photons with other polarization states are always simultaneously present and also enter the energy balance making the total dissipation independent from the heralding or remote interaction. In a broad sense, our results show that plasmonic losses,

whilst certainly detrimental for many applications, can also be harnessed as a resource, in line with other recent work in quantum plasmonics aimed at controlling quantum states via a loss mechanism^{30,31}. The approach shown here can be used not only for the nonlocal coupling of photons to localized plasmons, but can also be exploited for entangling photons to plasmon polaritons in polarization-sensitive schemes. Experiments such as plasmon teleportation⁵ or entanglement swapping³² can thus be envisaged in the future since our method based on coherent perfect absorption can be made quasi-perfect by optimizing the experimental parameters¹⁹. Currently, one cannot 'use' the single or entangled plasmons created due to their fast decaying behaviors but in the close future it would be interesting to study plasmons created at a metamaterial's interface through pump-probe measurements or to convert them back to photons

with a photonic structure, for instance with a waveguide.

All experimental data is available at <http://dx.doi.org/10.17861/39643814-5752-4a30-9db1-6f5d65b5c174>

Acknowledgements. The authors acknowledge the support of the Singapore MOE Grant MOE2011-T3-1-005, EPSRC (U.K.) grants EP/M009122/1 and EP/J00443X/1 and EU Grant ERC GA 306559. C.C. would like to thank the Champagne-Ardenne region, the French LABEX Action and the EU COST Action Nanoscale Quantum Optics. Authors acknowledge N. Zheludev for inspiring and supporting this work and T. Rogers, G. Adamo and Y. D. Chong for useful discussions.

- [1] Einstein, A.; Podolsky, B.; Rosen, N. Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Phys. Rev.* **1935**, *47*, 777-780.
- [2] Hensen, B.; Bernien, H.; Dréau, A. E.; Reiserer, A.; Kalb, N.; Blok, M. S.; Ruitenberg, J.; Vermeulen, R. F. L.; Schouten, R. N.; Abellán, C. et al. Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres. *Nature* **2015**, *526*, 682-686.
- [3] Giustina, M.; Versteegh, M. A. M.; Wengerowsky, S.; Handsteiner, J.; Hochrainer, A.; Phelan, K.; Steinlechner, F.; Kofler, J.; Larsson, J.-Å.; Abellán, C. et al. Significant-Loophole-Free Test of Bells Theorem with Entangled Photons. *Phys. Rev. Lett.* **2015**, *115*, 250401.
- [4] Shalm, L. K.; Meyer-Scott, E.; Christensen, B. G.; Bierhorst, P.; Wayne, M. A.; Stevens, M. J.; Gerrits, T.; Glancy, S.; Hamel, D. R.; Allman, M. S. et al. Strong Loophole-Free Test of Local Realism. *Phys. Rev. Lett.* **2015**, *115*, 250402.
- [5] Bouwmeester, D.; Pan, J.-W.; Mattle, K.; Eibl, M.; Weinfurter, H.; Zeilinger, A. Experimental quantum teleportation. *Nature* **1997**, *390*, 575-579.
- [6] Ma, S. X.; Herbst, T.; Scheidl, T.; Wang, D.; Kropatschek, S.; Naylor, W.; Wittmann, B.; Mech, A.; Kofler, J.; Anisimova, E. et al. Quantum teleportation over 143 kilometres using active feed-forward. *Nature* **2012**, *489*, 269-273.
- [7] Scully, M. O.; Drühl, K. Quantum eraser: A proposed photon correlation experiment concerning observation and "delayed choice" in quantum mechanics. *Phys. Rev. A.* **1982**, *25*, 2208.
- [8] Scully, M.O.; Englert, B-G.; Walther, H. Quantum optical tests of complementarity. *Nature* **1991**, *351*, 111-116.
- [9] Yoon-Ho, K.; Yu, R.; Kulik, S. P.; Shih, Y.; Scully, M. O. "Delayed Choice" Quantum Eraser. *Phys. Rev. Lett.* **2000**, *84*, 1.
- [10] Kwiat, P.G.; Weinfurter, H.; Herzog, T.; Zeilinger, A. Interaction-free measurement. *Phys. Rev. Lett.* **1995**, *74*, 4763.
- [11] Barreto Lemos, G.; Borish, V.; Cole, G. D.; Ramelow, S.; Lapkiewicz, R.; Zeilinger, A. Quantum imaging with undetected photons. *Nature* **2014**, *512*, 409-412.
- [12] Altewischer, E.; van Exter, M. P.; Woerdman, J. P. Plasmon-assisted transmission of entangled photons. *Nature* **2002**, *418*, 304-306.
- [13] Chang, D. E.; Sorensen, A. S.; Hemmer, P. R.; Lukin, M. D. Quantum optics with surface plasmons. *Phys. Rev. Lett.* **2006**, *97*, 053002.
- [14] Chang, D. E.; Sorensen, A. S.; Demler, E. A.; Lukin, M. D. A single-photon transistor using nanoscale surface plasmons. *Nat. Phys.* **2007**, *3*, 807-812.
- [15] Dutta Gupta, S.; Agarwal, G. S. Two-photon quantum interference in plasmonics: theory and applications. *Opt. Lett.* **2014**, *39*, 2.
- [16] Heeres, R. W.; Kouwenhoven, L. P.; Zwiller, V. Quantum interference in plasmonic circuits. *Nat. Nanotech.* **2013**, *8*, 719-722.
- [17] Fakonas, J. S.; Lee, H.; Kelaita, Y. A.; Atwater, H. A. Two-plasmon quantum interference. *Nat. Photon.* **2014**, *8*, 317-320.
- [18] Dheur, M.-C.; Devaux, E.; Ebbesen, T. W.; Baron, A.; Rodier, J.-C.; Hugonin, J.-P.; Lalanne, P.; Greffet, J.-J.; Messin, G.; Marquier, F. Single-plasmon interferences. *Sci. Adv.* **2016**, *2*, e1501574.
- [19] Roger, T.; Vezzoli, S.; Bolduc, E.; Valente, J.; Heitz, J.J.F.; Jeffers, J.; Soci, C.; Leach, J.; Couteau, C.; Zheludev, N. I.; Faccio, D. Coherent perfect absorption in deeply subwavelength films in the single-photon regime. *Nat. Commun.* **2015**, *6*, 7031.
- [20] Huang, S.; Agarwal, G. S. Coherent perfect absorption of path-entangled single photons. *Opt. Exp.* **2014**, *22*, 020936.
- [21] Pysher, M. J.; Galvez, E. J.; Misra, K.; Wilson, K. R.; Melius, B. C.; Malik, M. Nonlocal labeling of paths in a single-photon interferometer. *Phys. Rev. A.* **2005**, *72*, 052327.
- [22] Zhang, J.; MacDonald, K. F.; Zheludev, N. I. Controlling light-with-light without nonlinearity. *Light. Sci. Appl.* **2012**, *1*, e18.
- [23] Kurtsiefer, C.; Oberparleiter, M.; Weinfurter, H. High efficiency entangled photon pair collection in type II parametric fluorescence. *Phys. Rev. A.* **2001**, *64*, 023802.
- [24] Clauser, J. F.; Horne, M. A.; Shimony, A.; Holt, R. A.

- 1 Proposed experiment to test local hidden-variable theo-
2 ries. *Phys. Rev. Lett.* **1969**, 23, 880-884 (1969).
- 3 [25] Kwiat, P. G.; Mattle, K.; Weinfurter, H.; Zeilinger, A.;
4 Sergienko, A. V.; Shih, Y. New high-intensity source
5 of polarization-entangled photon pairs. *Phys. Rev. Lett.*
6 **1995**, 75, 4337-4341.
- 7 [26] Herbert, N. FLASH-A superluminal communicator based
8 upon a new kind of quantum measurement. *N. Found.*
9 *Phys.* **1982**, 12, 1171.
- 10 [27] Peres, A. How the no-cloning theorem got its name.
11 eprint quant-ph/0205076 **2002**.
- 12 [28] Fang, X.; MacDonald, K. F.; Zheludev, N. I. Control-
13 ling light with light using coherent metadvice: all-
14 optical transistor, summator and inverter. *Light. Sci.*
15 *Appl.* **2015**, 4, e292.
- 16 [29] Papaioannou, M.; Plum, E.; Valente, J.; Rogers, E. T. F.;
17 Zheludev, N. I. All-optical multichannel logic based on
18 coherent perfect absorption in a plasmonic metamaterial.
19 *APL Photonics* **2016**, 1, 090801.
- 20 [30] Roger, T.; Restuccia, S.; Lyons, A.; Giovannini, D.;
21 Romero, J.; Jeffers, J.; Padgett, M.; Faccio, D. Coher-
22 ent absorption of N00N states. *Phys. Rev. Lett.* **2016**,
23 117, 023601.
- 24 [31] Vest, B.; Dheur, M-C.; Devaux, E.; Ebbesen, T. W.;
25 Baron, A.; Rousseau, E.; Hugonin, J-P.; Greffet, J-
26 J.; Messin, G.; Marquier, F. Coalescence and anti-
27 coalescence of surface plasmons on a lossy beamsplitter.
28 arXiv:1610.07479 **2016**.
- 29 [32] Pan, J-W.; Bouwmeester, D.; Weinfurter, H.; Zeilinger,
30 A. Experimental entanglement swapping: entangling
31 photons that never interacted. *Phys. Rev. Lett.* **1998**,
32 80, 3891.
- 33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60