



Heriot-Watt University
Research Gateway

Universality and tails of long-range interactions in one dimension

Citation for published version:

Valiente Cifuentes, M & Öhberg, P 2017, 'Universality and tails of long-range interactions in one dimension', *Physical Review A*, vol. 96, no. 1, 012701. <https://doi.org/10.1103/PhysRevA.96.012701>

Digital Object Identifier (DOI):

[10.1103/PhysRevA.96.012701](https://doi.org/10.1103/PhysRevA.96.012701)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Physical Review A

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Universality and tails of long-range interactions in one dimension

Manuel Valiente and Patrik Öhberg

SUPA, Institute of Photonics and Quantum Sciences, Heriot-Watt University, Edinburgh EH14 4AS, United Kingdom

(Received 16 May 2017; published 7 July 2017)

Long-range interactions and, in particular, two-body potentials with power-law long-distance tails are ubiquitous in nature. For two bosons or fermions in one spatial dimension, the latter case being formally equivalent to three-dimensional s -wave scattering, we show how generic asymptotic interaction tails can be accounted for in the long-distance limit of scattering wave functions. This is made possible by introducing a generalization of the collisional phase shifts to include space dependence. We show that this distance dependence is universal, in that it does not depend on short-distance details of the interaction. The energy dependence is also universal, and is fully determined by the asymptotic tails of the two-body potential. As an important application of our findings, we describe how to eliminate finite-size effects with long-range potentials in the calculation of scattering phase shifts from exact diagonalization. We show that even with moderately small system sizes it is possible to accurately extract phase shifts that would otherwise be plagued with finite-size errors. We also consider multichannel scattering, focusing on the estimation of open channel asymptotic interaction strengths via finite-size analysis.

DOI: [10.1103/PhysRevA.96.012701](https://doi.org/10.1103/PhysRevA.96.012701)

I. INTRODUCTION

Few-body systems play a central role in quantum mechanics, which arguably started with Schrödinger's solution to the hydrogen atom [1]. Traditionally, the scattering states of a two-body system or, more specifically, their asymptotic form, are used to predict cross sections in atomic, molecular, nuclear, and particle physics [2,3], which give useful information about the underlying interactions and structure of the colliding bodies. When the interactions have a short range, there is a technique that has been championed by nuclear physicists to extract effective nucleon-nucleon and multinucleon interactions using few-body observables only. This is the effective field theory (EFT) of nuclear forces [4–11], which began with Weinberg's seminal papers [12,13]. In its modern form, especially when the system is discretized on a lattice, it is combined with energetic methods, i.e., exact diagonalization or imaginary time evolution, rather than traditional scattering theory, to predict scattering observables (for a pedagogical review, see Ref. [14]). The methodology involved is a generalization of the ground-breaking work of Lüscher [15], who was the first to show the relationship between low-energy scattering observables and the two-body spectrum in a periodic box, at least in the weak-coupling limit. The advantages of this kind of approach are most obvious when at least one of the particles in the system is a composite object, since it is generally easier to extract the ground-state energy of a three- or four-body system in a finite volume than to solve the corresponding multichannel Faddeev or Yakubovsky equations [16], or in generic multichannel problems.

The use of EFT is not confined to nuclear physics. In ultracold atoms, the lowest-order EFT, corresponding to the Huang-Yang pseudopotential [17], is routinely used in the theory of Bose-Einstein condensates [18] and spin-1/2 Fermi gases [19]. It plays a particularly important role where it is most accurate, that is, near an s -wave two-body resonance in a two-component Fermi gas, for which a set of universal relations hold in all spatial dimensions [20–25]. Effects beyond

lowest-order EFT, including three-body effects, are patent in the three-boson problem, where these play a major role in Efimov physics [26], observed for the first time with ultracold atoms by Kraemer *et al.* [27]. The development of EFTs in reduced (one and two) spatial dimensions is somewhat behind that of the three-dimensional case. There are two well-known examples of lowest-order EFT in one dimension (1D), namely the Dirac- δ interaction for bosons and spin-1/2 fermions, which is UV regular, and its odd-wave dual (sometimes termed p -wave), which is UV divergent but renormalizable, as first discussed by Cheon and Shigehara [28] in the position representation, and later on in the momentum representation [29,30]. The EFT to next-to-leading order for two-body scattering in 1D was also discussed in Ref. [29].

The effective interactions discussed above are all concerned with low-energy scattering. In one spatial dimension, however, we have shown [31] that Luttinger liquids whose constituents are scalar particles may depend very little on low-energy interactions except for extremely low densities, since the relevant energy scale for two-particle collisions is twice the Fermi energy. When realistic interactions are involved, such as Born-Oppenheimer potentials, which have Van der Waals tails, and especially in multichannel problems, it can be considerably easier to numerically diagonalize the two-body Hamiltonian in a finite box rather than solving the Schrödinger or Lippmann-Schwinger equations in infinite space. The scattering phase shifts can then, in principle, be extracted from extensions to Lüscher's analysis, for which there exists a vast and comprehensive literature [32–37]. Unfortunately, these approaches typically¹ assume that the interaction tails are negligible, a perfectly valid assumption in the context of these works. This, however, is hardly the case in atomic and molecular physics: interactions between two neutral atoms, a neutral atom and an ion, and two dipolar molecules,

¹An exception is Ref. [38], which deals with emergent, periodised Coulomb interactions relevant in Nuclear Physics.

display r^{-6} , r^{-4} , and r^{-3} tails, respectively. Therefore, exact diagonalization in a finite box can introduce very significant errors if the tails are not properly accounted for in the finite-size analysis of the phase shifts.

Here we study scattering in one dimension, which, for the fermionic case, is formally equivalent to three-dimensional s -wave collisions, for interactions that exhibit a long-range tail. First, we show how the introduction of space-dependent phase shifts yields universal, energy-independent information relating the long-range tail and the spatially varying part of the phase shift. We then show how to obtain scattering phase shifts using finite-size energy considerations only, by taking into account this universal asymptotic behavior. We also generalize our results to multichannel scattering.

II. HAMILTONIAN OF THE SYSTEM

We consider the nonrelativistic two-body scattering problem whose dynamics is governed by the following position-represented Hamiltonian

$$\mathcal{H} = \frac{p_1^2}{2m} + \frac{p_2^2}{2m} + V(x_1 - x_2). \quad (1)$$

Above, m is the mass of the particles, while V is a generic interaction potential, assumed to have a vanishing asymptotic limit, i.e., $\lim_{x \rightarrow \pm\infty} V(x) = 0$. The stationary Schrödinger equation $\mathcal{H}\Psi = \mathcal{E}\Psi$ is separable in terms of center of mass $X = (x_1 + x_2)/2$ and relative $x = x_1 - x_2$ coordinates in the usual way, i.e.,

$$\Psi(X, x) = e^{iKX} \psi(x), \quad (2)$$

where $K = k_1 + k_2$ is the total momentum of the system, while $\psi(x)$ is the relative wave function. This satisfies the stationary Schrödinger equation $H\psi = E\psi$, where $E = \mathcal{E} - \hbar^2 K^2/4m$, and H is given by

$$H = \frac{p^2}{2\mu} + V(x) \quad (3)$$

with $\mu = m/2$ the reduced mass, and p the relative momentum operator. Positive relative energies $E = \hbar^2 k^2/2\mu$ correspond to scattering states, which we will focus on in the following.

III. GENERAL FRAMEWORK

In order to establish the universal and nonuniversal properties of two-body collisions of identical particles, we consider the asymptotic, or long-distance limit of the Schrödinger equation $H\psi = E\psi$. Since $V(x) \rightarrow 0$ at long distances, the asymptotic form of the stationary scattering states, ψ_k^{asympt} , must have the form

$$\psi_B^{\text{asympt}}(x) = \sin[k|x| + \theta_k^B(x)], \quad (4)$$

$$\psi_F^{\text{asympt}}(x) = \text{sgn}(x) \sin[k|x| + \theta_k^F(x)]. \quad (5)$$

Above, the subscript and superscript B (F) refers to bosons or spin-singlet fermions (spin-triplet or spinless fermions). Notice that we have given the phase shifts θ^B and θ^F explicit space dependence. This is crucial for the analysis that follows. We now replace the interaction potential by its asymptotic

form $V^\infty(x)$, and insert the asymptotic scattering states, Eq. (4) or (5), into the Schrödinger equation. In this way, the following asymptotic differential equation is obtained for the position-dependent phase shift (we drop the superscript B or F for ease of notation)

$$\frac{1}{2} \left(\frac{d\theta_k}{dx}(x) \right)^2 + k \text{sgn}(x) \frac{d\theta_k}{dx}(x) + \frac{\mu V^\infty(x)}{\hbar^2} = 0, \quad (6)$$

subject to the conditions that $|\theta_k''(x)/V^\infty(x)| \rightarrow 0$ and $|\theta_k''(x)/\theta_k'(x)| \rightarrow 0$. If, moreover, $\theta_k''(x)$ is much smaller (in absolute value) than $[\theta_k'(x)]^2$, as is usually the case (see below for power-law potentials and a more detailed derivation in Appendix A), then the equation that must be solved for the sake of consistency is even simpler,

$$k \text{sgn}(x) \frac{d\theta_k}{dx}(x) + \frac{\mu V^\infty(x)}{\hbar^2} = 0, \quad (7)$$

which implies

$$\theta_k(x) = \theta_k - \frac{\mu}{\hbar^2 k} \int^x dx \text{sgn}(x) V^\infty(x). \quad (8)$$

As a convention, we choose the integration constant in Eq. (8) such that $\theta_k = \lim_{x \rightarrow \infty} \theta_k(x)$ corresponds to the scattering phase shift.

IV. ASYMPTOTIC POWER-LAW INTERACTIONS

The most physically relevant interactions in nature typically exhibit long-distance power-law tails. Such is the case of Coulomb, dipolar, or van der Waals tails that dictate electron-electron or atom-atom interactions. Inverse square interactions are also relevant in the three-body problem, especially in Efimov physics [39–41], and have also received some attention recently [42,43].

Power-law interaction tails are described by the asymptotic potential

$$V^\infty(x) = \frac{g_\nu}{|x|^\nu}. \quad (9)$$

The strength of the tails can be arbitrarily large and both repulsive ($g_\nu > 0$) or attractive ($g_\nu < 0$). The space-dependent phase shift must be calculated in this case from Eq. (8) and is given by

$$\theta_k(x) = \theta_k + \frac{\mu}{\hbar^2 k} \frac{g_\nu}{\nu - 1} \frac{1}{|x|^{\nu-1}}, \quad \nu \neq 1, \quad (10)$$

$$\theta_k(x) = \theta_k - \frac{\mu}{\hbar^2 k} g_1 \log |2kx|, \quad \nu = 1. \quad (11)$$

The above results are remarkable. While the constant part of the phase shifts θ_k must be calculated microscopically, the space-varying asymptotes are fully universal: not only are their functional forms fixed by the asymptotic tails of the interaction, but also their prefactors are determined by these. In particular, the famous logarithmic phase shifts of the Coulomb potential [44] appear naturally within the current framework, and the only nonuniversal feature of the phase shift is the constant θ_k . Notice, once more, that the microscopic short-distance details of the interactions do not enter the asymptotes besides in the value of θ_k , and smoothed interactions at short distances play no role here. This type of universality is similar in spirit

to quantum defect theory (QDT) [45], where the nonuniversal short-range details of the interaction are encoded in a few parameters. Unlike QDT, however, where these parameters determine the short-distance behavior of the scattering wave functions [45–48], which therefore require exact solutions (usually for power-law interactions) at short-to-intermediate distances, our results deal with asymptotic behavior only. Although, as will be clear below, our results are not as accurate as QDT when used to calculate scattering phase shifts, they are more general since their universality does not rely on exact solutions beyond the asymptotic regime.

V. CALCULATING THE CONSTANT PART OF THE PHASE SHIFT

The above analysis is not only useful from a theoretical point of view, but can also be used to extract numerically, in a very straightforward manner, the microscopic scattering phase shifts θ_k for interactions falling off faster than $1/|x|$, and the so-called Coulomb phase shifts for Coulomb interactions. To see this, consider the following fictitious problem: place a single particle of mass μ in a finite box of size $2L$ with open boundary conditions, i.e., $\psi(L) = \psi(-L) = 0$, with the full (i.e., not just the asymptotic part) interaction potential $V(x)$ centered at $x = 0$. Diagonalize the single-particle problem numerically and extract the eigenvalues. For sufficiently large L , such that the asymptotic potential is accurate for $x \sim L$, the eigenfunctions have the asymptotic behavior in Eqs. (4) or (5) for bosons and fermions, respectively. The energy of the eigenstates is given by $\hbar^2 k^2/2\mu$, and k is quantized by applying open boundary conditions to the asymptotic eigenfunctions (4) or (5). The quantization reads

$$k = \frac{\pi n}{L} - \frac{\theta_k(L)}{L}, \quad n \in \mathbb{Z}_+. \quad (12)$$

Since the eigenvalues are known after numerical diagonalization, so is $k = \sqrt{2\mu E/\hbar^2}$. By rewriting the space-dependent phase shift as $\theta_k(x) = \theta_k + \Delta_k(x)$, where $\Delta_k(x)$ is the universal space-dependent part of the phase shift, Eq. (12) is already solved and gives

$$\theta_k = \pi n - kL - \Delta_k(L). \quad (13)$$

There is a particularly simple model, namely the Calogero-Sutherland model [49], for which it is possible to obtain all eigenvalues given one of them. In this model, which has inverse square interactions, the phase shift $\theta_k^{(0)} = -\pi(\lambda - 1)/2$ [49], with $g_2 = (\hbar^2/m)\lambda(\lambda - 1)$ is a constant [up to a $\text{sgn}(k)$ factor], due to the fact that the model in free space is scale invariant. Scale invariance is only broken by the existence of a boundary in the system. In Fig. 1 we show the scattering phase shifts calculated numerically by diagonalizing the system (see Appendix B) and either including or neglecting the universal space dependence of the phase shift. There, it is clear that the space dependence is absolutely necessary at low energies. The smaller the kL value is, the larger the deviation is between the exact value and the result using the position-dependent phase shift. However, it is worth stressing that at $kL = 3.99005$ (the lowest possible value of kL) the error is still a mere 0.28%.

The pure Calogero-Sutherland interaction, being scale invariant and exactly solvable, is, however, not the best




FIG. 1. Calculated phase shifts for the Calogero-Sutherland model with $2\mu g_2/\hbar^2 = 1$. Calculations using Eq. (13) with (red circles) and without (blue squares) space-dependent phase shifts are compared to the exact phase shift (black diamonds).

example to illustrate the power of the method. We can also use a softened version of the $1/x^2$ potential, given by

$$V(x) = \frac{k_0^2 g_2}{1 + (k_0 x)^2}, \quad (14)$$

for which the asymptotic potential is also $V^\infty(x) = g_2/x^2$. In Fig. 2, the results for the fermionic phase shift from exact diagonalization with and without Δ_k are compared with well-converged values obtained from a numerical solution to the Lippmann-Schwinger equation (see Appendix C). Again, the highest error is small, approximately a 0.5%. As a last application, in Fig. 3 we show the constant part of the Coulomb phase shift, calculated from exact diagonalization, and compared to the exact result [44] $\theta_k = \text{Arg} \Gamma(1 + i\mu g_1/\hbar^2 k)$.

VI. TAIL RENORMALIZATION IN MULTICHANNEL COLLISIONS

So far, we have discussed the standard single-channel scattering with long-range interactions. Single-channel problems,




FIG. 2. Calculated fermionic phase shifts with the interaction in Eq. (14), with $2\mu g_2/\hbar^2 = 3$. Calculations using Eq. (13) with (red circles) and without (blue squares) space-dependent phase shifts, compared to the phase shift from integration of the Lippmann-Schwinger equation (black diamonds). Data are obtained for two different sizes, L_1 and L_2 , and $k_0 L_1 = 1$. Positive values of θ_k have been shifted by $-\pi$ for ease of visualization.




FIG. 3. Calculated phase shifts for the Coulomb interaction with $2\mu g_1 L/\hbar^2 = 1$ and $k_0 L = 1$. Calculations using Eq. (13) with space-dependent phase shifts (red circles) are compared to the exact phase shifts (black diamonds) obtained from Coulomb functions (see text).

however, can be quite efficiently and accurately solved by means of a brute force numerical implementation of the Lippmann-Schwinger equation. The solution to this yields the corresponding phase shifts and, complemented with the universal asymptotic spatially dependent phase shifts explained in the previous section, all relevant information about the scattering asymptotes can be extracted. Multichannel problems, however, are significantly more difficult to solve via brute-force integration of the Lippmann-Schwinger equation. In general, the incident waves of each channel must be carefully prepared before attempting a numerical solution. Moreover, if the multichannel nature of the system is due to multiparticle bound states, which effectively describe single particles, preparing the incident state and solving the integral equations numerically can prove to be quite challenging. An easy-to-visualize example would be the elastic collision of a two-body bound state with a heavy (static) particle. Notice that, even if all bare interactions in a given system have the same asymptotic tails, the strength of the tails of the open channel interactions, and in some cases also their functional form, are modified by the inclusion of all channels in the system. It is therefore of great interest to develop methodology that not only delivers scattering phase shifts, but also gives information about the tails of the effective open-channel interaction.

The general theory of multichannel asymptotes is, fortunately, not very different from the single-channel picture. We denote the different physical channels by α_i , and assume that the open channel (α_1) where elastic collisions are to be investigated has the lowest energy in its noninteracting ground state. The asymptotic form of the effective interaction in the elastic channel (α_1, α_1) is denoted by $V_{\alpha_1, \alpha_1}^\infty(x)$, such that the asymptotic scattering state in the open channel, $\psi_{\alpha_1}(x)$, satisfies

$$-\frac{\hbar^2}{2\mu} \frac{\partial^2 \psi_{\alpha_1}(x)}{\partial x^2} + V_{\alpha_1, \alpha_1}^\infty(x) \psi_{\alpha_1}(x) = (E - \mathcal{E}_{\alpha_1}) \psi_{\alpha_1}(x), \quad (15)$$

where \mathcal{E}_{α_1} is the noninteracting ground-state energy, on the infinite line, of the two-body elastic channel (α_1, α_1).

The most physically relevant situation corresponds to a power-law tail $V_{\alpha_1, \alpha_1}^\infty = g_v^{\alpha_1}/|x|^v$. This is the case, for instance, in multichannel models of Feshbach resonances [50]. If only

a finite number of channels are present, then the power ν and strength $g_v^{\alpha_1}$ of the tails can be easily calculated, as in the example shown below. When this is the case, the phase shifts can be calculated just as in single-channel scattering. If, on the other hand, there is an infinite number of channels (e.g., in models of confinement-induced resonances [51–56]), the exponent ν may change, and the strength must be calculated nonperturbatively. It is therefore important, especially for infinitely many channels, to devise a way to extract the strength of the tail in the open channel by energetic arguments only. To do this, take two boxes of lengths $2L_1$ and $2L_2$ ($L_1 \neq L_2$) both having one eigenstate corresponding to a certain momentum k , linked with the integers n_1 and n_2 ($n_1 \neq n_2$), respectively [see Eq. (12)]. Since both states have the same momentum, their phase shifts θ_k are identical. By equating the corresponding conditions for θ_k with lengths $2L_1$ and $2L_2$ in Eq. (13), and using that $\Delta_k(x) = \mu g_v^{\alpha_1}/[\hbar^2 k(\nu - 1)|x|^{\nu-1}]$, the following is found for the strength of the tail

$$g_v^{\alpha_1} = (\nu - 1) \frac{\hbar^2 k}{\mu} \frac{(L_1 L_2)^{\nu-1}}{L_1^{\nu-1} - L_2^{\nu-1}} [\pi(n_2 - n_1) + k(L_1 - L_2)]. \quad (16)$$

To illustrate and benchmark the above results in multichannel calculations, we consider a simple two-channel model, which is similar to what is sometimes used in modeling magnetic Feshbach resonances [50]. The Schrödinger equation has the form

$$-\frac{\hbar^2}{2\mu} \frac{\partial^2 \phi_1(x)}{\partial x^2} + V_1(x) \phi_1(x) - J \phi_2(x) = E \phi_1(x), \quad (17)$$

$$-\frac{\hbar^2}{2\mu} \frac{\partial^2 \phi_2(x)}{\partial x^2} + V_2(x) \phi_2(x) - J \phi_1(x) = E \phi_2(x). \quad (18)$$

Above, $J > 0$ is a constant that represents the coupling between the bare channel wave functions ϕ_1 and ϕ_2 . The bare channels, 1 and 2, must be reexpressed in terms of physical channels, A and B , by means of the transformation $\psi_A = \phi_1 + \phi_2$ and $\psi_B = \phi_1 - \phi_2$, up to a global normalization constant. Defining $V_A(x) = V_B(x) = [V_1(x) + V_2(x)]/2$ and $J_A(x) = J_B(x) = [V_1(x) - V_2(x)]/2$, the multichannel equations become

$$-\frac{\hbar^2}{2\mu} \frac{\partial^2 \psi_A(x)}{\partial x^2} + V_A(x) \psi_A(x) + J_A(x) \psi_B(x) = (E - \mathcal{E}_A) \psi_A(x), \quad (19)$$

$$-\frac{\hbar^2}{2\mu} \frac{\partial^2 \psi_B(x)}{\partial x^2} + V_A(x) \psi_B(x) + J_A(x) \psi_A(x) = (E - \mathcal{E}_B) \psi_B(x), \quad (20)$$

where $\mathcal{E}_A = -J$ ($\mathcal{E}_B = +J$) is the open (closed) channel's noninteracting ground-state energy. In the numerical example, we use fermions and choose $\nu = 3$ (dipolarlike tails), and $V_i(x) = k_0^3 g_3^{(i)}/[1 + (k_0 x)^3]$, $i = 1, 2$, with $g_3^{(1)}/g_3^{(2)} = 2$. From Eq. (19), we clearly have $g_3^A = 3g_3^{(1)}/4$. In order to test how accurate the estimation of the tails in the open channel can be in more involved calculations (given finite-size effects and numerical uncertainty in exact diagonalization), we set L_1 to a

constant, for which its ground-state momentum $k = 0.3406k_0$ is extracted. Then, we numerically find three other values, L_2, L_3 , and L_4 such that the first, second, and third excited states, respectively, correspond to momentum k . Using these values in Eq. (16), we obtain six different estimates of g_3^A , giving the result $g_3^A/g_3^{(1)} = 0.770 \pm 0.028$ (see Appendix D). The exact value of $3/4$ is reproduced within error bars. Higher accuracy can be obtained by using many different values of L_1 , due to the energy independence of the tail strengths in this case. The phase shift is then estimated using Eq. (13) to be $\theta_k = -0.2693 \pm 9 \times 10^{-4}$.

VII. CONCLUSIONS

We have studied the asymptotic form of scattering states with interactions that have a long-range tail in one dimension, which, in the odd-wave channel, also describe three-dimensional s -wave scattering. We have shown how a generalization of phase shifts in the asymptotic scattering states yields a universal space dependence of these, independent of energy and short-distance details of the interaction. These results have been used to generalize widespread methods in nuclear physics to extract scattering information via finite-size analysis to the atomic situation where long-range tails are very important. In particular, we have exemplified our findings with different power-law tails, including the Coulomb interaction, and found that it is possible to extract phase shifts accurately using exact, finite-size diagonalization, even for relatively small system sizes. We have studied generic multichannel problems, for which the single-channel results are valid as well, and obtained an expression that relates finite-size information with the strength of the asymptotic tails of the interaction in the open elastic channel. We studied an example of a two-channel model showing that our method is a viable alternative to coupled-channel numerical calculations of scattering states. Our results and methods may be especially relevant for multichannel problems with an infinite number of coupled channels, as is the case in dimensional reduction and confinement-induced resonances. The methods we have introduced can also be of great use for the nonexpert in few-body physics, as it is generally easier to implement than other approaches.

ACKNOWLEDGMENTS

The authors acknowledge support from EPSRC Grant No. EP/M024636/1. The work of M.V. was initiated at the Aspen Center for Physics, which is supported by National Science Foundation Grant No. PHY-1066293.

APPENDIX A: DERIVATION OF THE DIFFERENTIAL EQUATION

Here, we present a detailed derivation of the differential equations for the space-dependent phase shifts in the asymptotic limit, Eqs. (6) and (7). For concreteness, we derive these for $x > 0$, and the case $x < 0$ follows analogously. For $x > 0$, both bosonic and fermionic asymptotic wave functions have the form $\psi^{\text{asympt}}(x) = \sin[kx + \theta_k(x)]$. We insert the scattering wave function into the stationary Schrödinger

equation $H\psi^{\text{asympt}} = (\hbar^2k^2/2\mu)\psi^{\text{asympt}}$, with H in Eq. (3) and $V(x)$ replaced by its asymptotic value $V^\infty(x)$. The following equation is obtained

$$-\left[\left(\frac{d\theta_k}{dx}(x)\right)^2 + 2k\frac{d\theta_k}{dx}(x) + \frac{\mu V^\infty(x)}{\hbar^2}\right]\psi^{\text{asympt}}(x) + \frac{d^2\theta_k}{dx^2}(x)\cos[kx + \theta(x)] = 0. \quad (\text{A1})$$

From Eq. (A1), we obtain Eq. (6) if both $|\theta_k''(x)/V^\infty(x)| \rightarrow 0$ and $|\theta_k''(x)/\theta_k'(x)| \rightarrow 0$ as $x \rightarrow \infty$. We need to ensure consistency in the order of approximation as well. Quite generally, e.g., with power-law or exponentially decaying potentials, $[\theta'(x)]^2$ is much smaller than $|\theta''(x)|$. Therefore, in Eq. (A1) we must drop the first term in the bracket, which yields Eq. (7). It is easy to see that the above conditions, for the case of power-law potentials $V^\infty(x) \sim |x|^{-\nu}$, simplify to $k|x| \gg 1$, i.e., the asymptotic limit.

APPENDIX B: EXACT DIAGONALIZATION

We explain here the details of the exact diagonalization used to obtain the results in the main text. We have chosen to discretize the system on a grid, since this is the simplest possible method, and is capable of giving accurate results if analyzed properly.

We discretized the Laplacian using a third-order quadrature rule. We use L_s equally spaced grid points with a lattice spacing d such that the length of the box is $L_s d$. The stationary Schrödinger equation is therefore discretized as

$$-\sum_{\mu=1}^3 J_{|\mu|}[\psi(n+\mu) + \psi(n-\mu)] + 2(J_1 + J_2 + J_3 + V(n) - E)\psi(n) = 0, \quad (\text{B1})$$

where $J_1 = (3/2d^2)\hbar^2/2\mu$, $J_2 = -(3/20d^2)\hbar^2/2\mu$ and $J_3 = (1/90d^2)\hbar^2/2\mu$. We diagonalize the Hamiltonian numerically for L_s between 201 and 801, with d adjusted so that L is kept constant, and the energy of a given state as a function of the number of grid points is $E(L_s)$. We then fit the continuum limit E_∞ as

$$E(L_s) = E_\infty + \alpha L_s^{-1} + \beta L_s^{-2}. \quad (\text{B2})$$

The error in the least-square fits to E_∞ are of the order of $10^{-6}\%$ in all cases we studied. The same types of fits are done for the phase shifts and the extracted values of the incident momentum of the scattering states, with similar fitting errors.

APPENDIX C: LIPPMANN-SCHWINGER EQUATION

The Lippmann-Schwinger equation for bosons and fermions in one dimension reads (see, for instance, Ref. [57])

$$\psi(x) = \psi_0(x) + \frac{\mu}{\hbar^2 k} \int_{-\infty}^{\infty} dy \sin(k|x-y|)V(y)\psi(y), \quad (\text{C1})$$

where $\psi_0(x) = \cos(kx)$ for bosons and $\psi_0(x) = \sin(kx)$ for fermions. Notice that the Lippmann-Schwinger equation for one-dimensional fermions and three-dimensional s -wave scattering, after rearrangement of the integration limits in Eq. (C1), are equivalent [3]. The phase shifts θ_k^{B} and θ_k^{F} for bosons and

fermions, respectively, are obtained from the scattering wave functions as

$$\cot \theta_k^B = \frac{\mu}{\hbar^2 k} \int_{-\infty}^{\infty} dy V(y) \cos(ky) \psi_k(y), \quad (\text{C2})$$

$$\tan \theta_k^F = -\frac{\mu}{\hbar^2 k} \int_{-\infty}^{\infty} dy V(y) \sin(ky) \psi_k(y). \quad (\text{C3})$$

The numerical solution of Eq. (C1) is obtained by discretizing the integral in Eq. (C1) using Gauss quadrature with a large distance cutoff Λ , and solving the resulting system of linear equations. The phase shifts, from Eqs. (C2) and (C3), are obtained using Gauss quadrature.

APPENDIX D: DETAILS OF THE MULTICHANNEL CALCULATION

The two interaction potentials V_1 and V_2 of the bare channels in Eqs. (17) and (18) are given by

$$V_i(x) = \frac{k_0^3 g_3^{(i)}}{1 + (k_0 x)^3}. \quad (\text{D1})$$

In the example in the text, we chose $2\mu k_0 g_3^{(1)}/\hbar^2 = 1$ and $g_3^{(2)} = g_3^{(1)}/2$. The length L_1 is chosen such that $k_0 L_1 = 10$, which is large enough that the interaction potentials $V_i(x)$ are well in the asymptotic regime at $x = L_1$. The value of the interchannel coupling constant is $J = \hbar^2 k_0^2 / 2\mu$. The ground-state momentum [i.e., corresponding to $n = 1$ in Eq. (12)] as obtained by numerically diagonalizing the two-channel Hamiltonian in the bare $\{1, 2\}$ basis is given in this case by $k/k_0 = 0.34060$. We then found three other lengths L_i , $i = 2, 3, 4$ such that the $(i - 1)$ th excited state corresponds to $k/k_0 = 0.34060$. These lengths were found to be $L_2 = 1.92362L_1$, $L_3 = 2.84622L_1$, and $L_4 = 3.76868L_1$. Using Eq. (16) for all pairs with different (L_i, L_j) , six different estimates for $g_3^A/g_3^{(1)}$ are found: $\{0.786361, 0.794766, 0.783350, 0.783521, 0.742459, 0.726599\}$. Their average $\langle g_3^A \rangle / g_3^{(1)}$ and standard deviation σ are therefore given by

$$\frac{\langle g_3^A \rangle}{g_3^{(1)}} = 0.770, \quad (\text{D2})$$

$$\sigma = 0.028, \quad (\text{D3})$$

which are the values reported in Sec. VI.

-
- [1] E. Schrödinger, *Ann. Phys. (NY)* **384**, 361 (1926).
 [2] J. R. Taylor, *Scattering Theory* (Dover Publications, Mineola, 2006).
 [3] C. J. Joachain, *Quantum Collision Theory* (North-Holland, Amsterdam, 1975).
 [4] R. Machleidt and D. R. Entem, *Phys. Rep.* **503**, 1 (2011).
 [5] E. Epelbaum, [arXiv:1001.3229v1](https://arxiv.org/abs/1001.3229v1).
 [6] U. van Kolck, *Prog. Part. Nucl. Phys.* **43**, 337 (1999).
 [7] H.-W. Hammer, A. Nogga, and A. Schwenk, *Rev. Mod. Phys.* **85**, 197 (2013).
 [8] E. Epelbaum, H.-W. Hammer, and U.-G. Meissner, *Rev. Mod. Phys.* **81**, 1773 (2009).
 [9] D. R. Phillips, S. R. Beane, and T. D. Cohen, *Nucl. Phys. A* **631**, 447 (1998).
 [10] D. B. Kaplan, M. J. Savage, and M. B. Wise, *Phys. Lett. B* **424**, 390 (1998).
 [11] D. B. Kaplan, M. J. Savage, and M. B. Wise, *Nucl. Phys. B* **534**, 329 (1998).
 [12] S. Weinberg, *Phys. Lett. B* **251**, 288 (1990).
 [13] S. Weinberg, *Nucl. Phys. B* **363**, 3 (1991).
 [14] A. Nicholson, in *An Advanced Course in Computational Nuclear Physics*, edited by M. Hjorth-Jensen, M. P. Lombardo, and U. van Kolck, Lecture Notes in Physics (Springer International Publishing, Cham, 2017), Vol. 936, pp. 155–235.
 [15] M. Lüscher, *Commun. Math. Phys.* **105**, 153 (1986).
 [16] M. Pine, D. Lee, and G. Rupak, *Eur. Phys. J. A* **49**, 151 (2013).
 [17] K. Huang and C. N. Yang, *Phys. Rev.* **105**, 767 (1957).
 [18] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999).
 [19] S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* **80**, 1215 (2008).
 [20] S. Tan, *Ann. Phys. (NY)* **323**, 2952 (2008).
 [21] S. Tan, *Ann. Phys. (NY)* **323**, 2971 (2008).
 [22] S. Tan, *Ann. Phys. (NY)* **323**, 2987 (2008).
 [23] M. Valiente, N. T. Zinner, and K. Mølmer, *Phys. Rev. A* **84**, 063626 (2011).
 [24] M. Valiente, N. T. Zinner, and K. Mølmer, *Phys. Rev. A* **86**, 043616 (2012).
 [25] F. Werner and Y. Castin, *Phys. Rev. A* **86**, 013626 (2012).
 [26] P. F. Bedaque, H.-W. Hammer, and U. van Kolck, *Phys. Rev. Lett.* **82**, 463 (1999).
 [27] T. Kraemer *et al.*, *Nature (London)* **440**, 315 (2006).
 [28] T. Cheon and T. Shigehara, *Phys. Rev. Lett.* **82**, 2536 (1999).
 [29] M. Valiente and N. T. Zinner, *Few-Body Syst.* **56**, 845 (2015).
 [30] X. Cui, *Phys. Rev. A* **94**, 043636 (2016).
 [31] M. Valiente and P. Öhberg, *Phys. Rev. A* **94**, 051606 (2016).
 [32] S. R. Beane, P. F. Bedaque, A. Parreño, and M. J. Savage, *Phys. Lett. B* **585**, 106 (2004).
 [33] M. Fukugita, Y. Kuramashi, M. Okawa, H. Mino, and A. Ukawa, *Phys. Rev. D* **52**, 3003 (1995).
 [34] S. Tan, *Phys. Rev. A* **78**, 013636 (2008).
 [35] M. Valiente and N. T. Zinner, *Sci. China Phys. Mech. Astron.* **59**, 114211 (2016).
 [36] M. T. Hansen and S. R. Sharpe, *Phys. Rev. D* **93**, 014506 (2016).
 [37] S. R. Beane, *Phys. Rev. A* **82**, 063610 (2010).
 [38] S. R. Beane and M. J. Savage, *Phys. Rev. D* **90**, 074511 (2014).
 [39] P. Naidon and S. Endo, *Rep. Prog. Phys.* **80**, 056001 (2017).
 [40] C. H. Greene, P. Giannakeas, and J. Perez-Rios, [arXiv:1704.02029](https://arxiv.org/abs/1704.02029).
 [41] E. Nielsen, D. V. Fedorov, A. S. Jensen, and E. Garrido, *Phys. Rep.* **347**, 373 (2001).
 [42] S. Moroz, J. P. D’Incao, and D. S. Petrov, *Phys. Rev. Lett.* **115**, 180406 (2015).
 [43] N. L. Guevara, Y. Wang, and B. D. Esry, *Phys. Rev. Lett.* **108**, 213202 (2012).
 [44] M. Abramowitz and I. A. Stegun (eds.), *Handbook of Mathematical Functions* (Dover Publications, Mineola, 1965).

- [45] C. Greene, U. Fano, and G. Strinati, *Phys. Rev. A* **19**, 1485 (1979).
- [46] B. Gao, *Phys. Rev. A* **64**, 010701 (2001).
- [47] J. Joger, A. Negretti, and R. Gerritsma, *Phys. Rev. A* **89**, 063621 (2014).
- [48] Z. Idziaszek, T. Calarco, and P. Zoller, *Phys. Rev. A* **76**, 033409 (2007).
- [49] B. Sutherland, *Beautiful Models* (World Scientific, Singapore, 2004).
- [50] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, *Rev. Mod. Phys.* **82**, 1225 (2010).
- [51] M. Olshanii, *Phys. Rev. Lett.* **81**, 938 (1998).
- [52] S. Sala, P.-I. Schneider, and A. Saenz, *Phys. Rev. Lett.* **109**, 073201 (2012).
- [53] S. Sala and A. Saenz, *Phys. Rev. A* **94**, 022713 (2016).
- [54] S. Sala, G. Zürn, T. Lompe, A. N. Wenz, S. Murmann, F. Serwane, S. Jochim, and A. Saenz, *Phys. Rev. Lett.* **110**, 203202 (2013).
- [55] M. Valiente and K. Mølmer, *Phys. Rev. A* **84**, 053628 (2011).
- [56] E. Haller, M. J. Mark, R. Hart, J. G. Danzl, L. Reichsöllner, V. Melezhik, P. Schmelcher, and H.-C. Nägerl, *Phys. Rev. Lett.* **104**, 153203 (2010).
- [57] M. Valiente, L. G. Phillips, N. T. Zinner, and P. Öhberg, [arXiv:1505.03519](https://arxiv.org/abs/1505.03519).