



Heriot-Watt University
Research Gateway

A self-assembled nanotube supported by halogen bonding interactions

Citation for published version:

Kennedy, SR, Main, MU, Pulham, CR, Ling, I & Dalgarno, SJ 2019, 'A self-assembled nanotube supported by halogen bonding interactions', *CrystEngComm*, vol. 21, no. 5, pp. 786-790.
<https://doi.org/10.1039/C8CE01824C>

Digital Object Identifier (DOI):

[10.1039/C8CE01824C](https://doi.org/10.1039/C8CE01824C)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Peer reviewed version

Published In:

CrystEngComm

Publisher Rights Statement:

© The Royal Society of Chemistry 2019

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

A self-assembled nanotube supported by halogen bonding interactions

 Stuart R. Kennedy,^a Mawgan U. Main,^b Colin R. Pulham,^{*a} Irene Ling^{*c} and Scott J. Dalgarno^{*b}

 Received 00th January 20xx,
 Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Di-propoxycalix[4]arene is known to self assemble into a nanotube in the solid state. The introduction of two bromine atoms at distal positions across the upper-rim of the calixarene framework modulates nanotube packing, a feature supported by complementary halogen bonding interactions.

Controlled self-assembly remains a fundamental challenge for the supramolecular chemist, especially when one considers that many types of intermolecular interaction can play a role in directing the final outcome. Calix[4]arenes (C[4]s) are cyclic polyphenols, many of which adopt cone or pinched-cone conformations depending on the degree of functionalisation (e.g. alkylation) at the lower-rim.¹ C[4]s have played a pivotal role in the development of supramolecular chemistry, and much has been learned from detailed single crystal X-ray diffraction (SCXRD) studies; an excellent example of this is the emergence and understanding of porosity in the seemingly non-porous sublimate of *p*-¹Bu-calix[4]arene.² In the vast majority of cases, including the aforementioned example, SCXRD studies have revealed bi-layer arrangements in which the molecules alternate in an up-down anti-parallel manner. This is due to the formation of many non-covalent interactions (e.g. CH... π and π -stacking) between the constituent molecules within these highly favourable arrangements (Fig. 1A).³

Although this is the case, there are a small number of reported examples in which cone-shaped C[4]s buck the trend, packing in an alternative parallel fashion to (typically) afford spherical or tubular assemblies; parallel packing invokes

curvature in any prevailing structure because of the cone shape of the constituent building blocks (Fig. 1B).⁴ Di-propoxycalix[4]arene, **1**, is a molecule that has been shown to pack as a triply helical nanotube in the solid state (Fig. 1C).⁵ The central core of the nanotube is compact, and the lower-rim propoxy chains (that point inwards) are arranged such that there is only a very small channel present. Neighbouring nanotubes pack through 'cog-like' interdigitation, with symmetry equivalent (s.e.) molecules self-including as shown in Figure 1C; this interdigitation is stabilised by complementary π -stacking and CH... π interactions within the cavities of s.e. of **1**. In previous work we showed that it was possible to modulate the packing of this nanotube motif through the introduction of carboxylic acids at the upper-rim of **1**.⁶ Subsequent crystallisation from pyridine had the effect of increasing inter-tubule spacing by from ~ 20 Å to ~ 27 Å due to the formation of complementary host-guest heterosynthons between neighbouring nanotubes (Fig. 1D).




Figure 1. A) Schematic of anti-parallel bilayer C[4] packing. B) Schematic of parallel C[4] packing with induced curvature. C) Expanded structure of **1** showing nanotube spacing and interdigitation.⁵ D) Heterosynthon formation between the *p*-carboxylate analogue of **1** upon crystallisation from pyridine.⁶ Dashed lines represent hydrogen bonding interactions. H atoms omitted for clarity in C and D except for those involved in H-bonding interactions. Colour code: C – grey, O – red, N – blue, H – white. Figures not to scale.

Halogen bonding has emerged as an excellent addition to the toolbox of interactions that the supramolecular chemist can exploit when seeking to control or influence self-assembly and

^a School of Chemistry, Joseph Black Building, David Brewster Road, Edinburgh, EH9 3FJ, UK. Email: C.R.Pulham@ed.ac.uk

^b Institute of Chemical Sciences, William Perkin Building, Heriot-Watt University, Riccarton, Edinburgh, Scotland EH14 4AS, UK. Email: S.J.Dalgarno@hw.ac.uk

^c School of Science, Monash University Malaysia, Jalan Lagoon Selatan, Bandar Sunway, 46150 Selangor, Malaysia. E-mail: ireneling@monash.edu

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: Details of the Hirshfeld analyses, additional figures to support discussion and the Crystallographic information file (CIF) for **2**. See DOI: 10.1039/x0xx00000x

/ or host-guest chemistry.⁷ With this in mind, we have begun to investigate how this type of interaction may affect the assembly behaviour of C[4] building blocks, in particular those that exhibit preferences to form tubular assemblies. In this contribution we report our initial findings concerning the effect of halogen introduction at the upper-rim of the framework in **1**, notably the ability to modulate nanotube packing through the formation of complementary halogen bonding interactions. We also present a concise summary of these building blocks, some of which exhibit a tendency to undergo parallel packing; this is with a view to outlining the potential to utilise a range of intermolecular interactions (including halogen bonding) to drive or control the formation of such challenging assemblies.

Upper-rim halogenation of the C[4] framework can depend on a number of different factors, an example being the presence or absence of directing groups at the lower-rim that subsequently influence functionalisation. Given that the nanotube assembly of **1** was our target for modulation, we synthesised readily accessible halogenated derivatives **2** – **4** as shown in Figure 2.^{8–12} Crystallisation of **2** and **3** was undertaken in a range of common laboratory solvents. Single crystals were obtained in just one case, that being compound **2** from acetone, although they were not found to be a solvate (vide infra). In stark contrast, compound **4** was found to be poorly soluble, and as such it was not possible to obtain single crystals to monitor any effect of upper-rim iodination.




Figure 2. Upper-rim halogenated derivatives synthesised and used in the present study.

Colourless single crystals of **2** that were suitable for diffraction studies were obtained upon dissolution in acetone and standing over a number of weeks.[‡] The crystals were found to be of trigonal symmetry and structure solution was carried out in the space group *R*-3. The asymmetric unit comprises one molecule of **2**, and symmetry expansion at the upper-rim reveals that s.e. molecules self-include in a cog-like manner (Fig. 3A) akin to that of **1** (Fig. 1C), despite the fact that two H atoms at the upper-rim have been replaced with bromines. Comparison of the self-included dimers in **1** and **2** reveals that the distance between centroids generated between the lower-rim oxygens has increased from 8.85 Å to 10.03 Å respectively as a direct result of halogenation coupled with self-inclusion.⁵ Further expansion of the structure shows a concomitant increase in the intertubule spacing, moving from ~20 Å to ~23 Å (compare Figs. 1C and 3B). This represents approximately half of the modulation achieved via the introduction of carboxyl groups to the framework of **1** as outlined above.⁶

Given the versatility observed for this nanotube system, we thought it pertinent to survey the Cambridge Structural Database (CSD) with a view to establishing how rare these combined features are with respect to assembly. A search of the CSD¹³ for all structures containing lower-rim di-alkoxy C[4]s reveals that dimeric or cog-like self-inclusion is a relatively common assembly motif (41 hits).^{5,14,15} This self-inclusion phenomenon is also found to be tolerant towards the presence or a variety of upper-rim groups, as long as they are able to form complementary host-guest interactions with the cavity of a s.e. molecule.




Figure 3. A) Self-included dimer found in the single crystal X-ray structure of **2**. B) Cog-like interdigitation of nanotubes of **2** showing the intertubule spacing of ~23 Å. C) Picture from the Hirshfeld analysis showing the Br...arene interaction found in the self-included dimer as a red spot on the cavity interior surface. H atoms omitted for clarity in A and B. Colour code: C – grey, O – red, H – white, Br – maroon. Figures not to scale.

Hirshfeld analysis¹⁶ of **2** (see Supporting Information) reveals a crystallographically unique Br... π interaction as shown by the red spot on the cavity interior surface in Figure 3C; this occurs with a Br...C distance of 3.486 Å. Examination of the regions

between the nanotubes reveals chains of Ar-H...Br interactions but little evidence of Br...Br interactions. Given that there were no directly relevant literature examples for structural comparison with **2**, we expanded our search criteria accordingly as outlined below.

A search for C[4]s that are doubly substituted at the upper-rim with distally positioned halogens returned just 11 hits, 5 of which deviate from the cone conformation and thus are not useful for comparison. Four of the six remaining hits (CSD codes GUDMEY, QADTUL, WAZZOO and XIGLEE) are lower-rim tetra-substituted C[4]s,¹⁷ meaning that they necessarily adopt pinched cone conformations in the solid state, and as a consequence of this do not have cavities occupied by either guest molecules or s.e. C[4]s. One of the two remaining structures (KEHTEX) is a heterobimetallic cluster in which a lithium ion and ligated THF occupy the C[4] cavity,¹⁸ leaving just one relevant hit (XIGLAA) in which there is a di-halogenated C[4] with an open cavity suitable for forming host-guest interactions.^{17d} Inspection of this structure reveals that the upper-rim substitution pattern is analogous to that of compound **3**, as bromination has been performed *para*- to the lower-rim hydroxyl groups rather than the alkoxy groups; the C[4] in XIGLAA differs from **3** in that it possesses lower-rim benzyloxy rather than propoxy groups. Symmetry expansion of the ASU in XIGLAA reveals self-inclusion (Fig. S1) in an analogous manner to that found in **1**, and not involving Br... π interactions. Considering the structure of **2**, the self-inclusion in XIGLAA may be a direct result of alternative halogenation positions, but this will require broader investigation.

Expansion of the search criteria to include C[4]s halogenated at all four upper-rim positions returned a total of 17 hits.¹³ Similar elimination of non-cone conformers reduced this to 10 hits, 8 of which have the C[4] in a pinched-cone conformation due to lower-rim tetra-substitution (CSD codes FAJFAX, FOQDAR, HACYIU, HACYOA, KARNIB01, NIGPOI, QADTOF and QUWLOK).^{17b, 19} The 2 remaining hits, (CIPQEY²⁰ and QEVBEZ²¹) both contain 5,11,17,23-tetrabromo-25,27-dimethoxy-26,28-dihydroxycalix[4]arene. The cavity of the C[4] in CIPQEY is occupied by a different guest molecule so can be eliminated for the purposes of comparison with **2**. Symmetry expansion of the ASU in QEVBEZ does generate a self-included dimer and, interestingly, Hirshfeld analysis¹⁶ reveals that there are no short Br... π interactions (Fig. S2) as is the case for **2**. This is surprising, and appears to be due to slight shifting of the C[4]s such that they are aligned in a more symmetric manner (i.e. less offset within the confines of the cavity).

Further expansion to include calix[4]arene and a halobenzene fragment returned 11 hits,¹³ none of which are relevant or useful for comparison with the halogen interactions present in the structure of **2**. Finally, expansion of criteria to include other halogenated guests, as well as other structurally related hosts, returned a number of hits reported in a study by Diederich and co-workers, all of which relate to the elegant encapsulation of monohalo- and (\pm)-*trans*-1,2-dihalocyclohexanes in enantiopure alleno-acetylnic cages.²² Although markedly different with respect to host structure, these complexes are useful for comparison with the structure

of **2**, as halogen... π interactions occur to varying extents depending on the particular guest being encapsulated. In these examples, a Br... π distance of 3.6 Å is recorded for a well confined guest in the resorcinarene-based cavitand host, and the average interaction distance can be correlated to polarizability of the halogen.

The shorter Br... π distance observed in **2** may be due to a series of factors that likely includes the propensity of **1** to pack in a parallel manner / in nanotubes. That said, a wide-ranging study of halogenated C[4] derivatives, if one can overcome solubility / synthetic issues, would provide great insight into halogen polarizability effects with respect to self-assembly with these multi-component systems.

Conclusions

To conclude, we have reported the modulation of a self-assembled C[4] nanotube through the formation of halogen bonding interactions. A detailed survey of the CSD reveals that there is relatively little structural information available for useful comparison, suggesting that much is yet to be unearthed with respect to halogen bonding and its use in the controlled self-assembly of C[4]-based building blocks. This is particularly interesting as one may potentially utilise the C[4] cavity in concert with synthetic alteration at various regions of the molecular framework in order to promote the construction of targeted assemblies. For example, the ability to control or drive parallel packing of C[4]s in order to reliably form nanotube or spherical assemblies is an extremely challenging goal, but one that may be achieved once an enhanced understanding of how specific interactions within the C[4] cavity guide assembly protocols. Future work will focus on expanding our library of halogenated C[4] derivatives, addressing solubility issues, and using halogen bonding interactions to promote the formation of systems that deviate from anti-parallel bi-layer packing.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank the Royal Society Newton Mobility Fund for supporting this work under grant reference NI170151. We would also like to thank Dr Gary Nichol for assistance with collection of single crystal diffraction data.

Notes and references

‡ Compounds **2** – **4** were synthesised according to literature procedures.¹⁰⁻¹² Crystallisation of **2**: Compound **2** was dissolved in acetone and diffraction quality single crystals grew upon standing over a number of weeks. **Crystal data for compound 2 (CCDC 1875214)**: C₃₄H₃₄Br₂O₄, *M* = 666.43 g/mol, trigonal, space group R-3, *a* = 37.3853(7) Å, *c* = 11.2109(2) Å, *V* = 13569.8(6) Å³, *Z* = 18, *T* = 120.01(10) K, μ (Mo K α) = 2.725 mm⁻¹, *D*_{calc} = 1.468 g/cm³, 50040 reflections measured (5.766° \leq 2 θ \leq 59.31°), 7953 unique (*R*_{int} =

0.0668, $R_{\text{sigma}} = 0.0602$) which were used in all calculations. The final R_1 was 0.0523 ($I > 2\sigma(I)$) and wR_2 was 0.1164 (all data).

- For example of synthetic alteration and subsequent conformational control over the calix[4]arene framework see: C. D. Gutsche, Chapter 1 in *Calixarenes 2001*, Kluwer Academic Publishers, Dordrecht, 2001; I. Thondorf, A. Shivanyuk and V. Böhmer, Chapter 1 in *Calixarenes 2001*, Kluwer Academic Publishers, Dordrecht, 2001.
- S. J. Dalgarno, P. K. Thallapally, L. J. Barbour and J. L. Atwood, *Chem. Soc. Rev.*, 2007, **36**, 236, and refernces therein.
- For a relevant review of calix[4]arene bi-layer formation see: J. L. Atwood, L. J. Barbour, M. J. Hardie and C. L. Raston, *Coord. Chem. Rev.*, 2001, **222**, 3;
- For example see: G. W. Orr, L. J. Barbour, J. L. Atwood, *Science*, 1999, **285**, 1049; I. Ling, H. Kumari, M. Mirzamani, A. N. Sobolev, C. J. Garvey, J. L. Atwood and C. L. Raston, *Chem. Commun.*, 2018, **54**, 10824.
- L. G. Kuz'mina, G. G. Sadikov, J. A. K. Howard, E. A. Shokova and V. V. Kovalev, *Kristallografiya*, 2003, **48**, 272; S. Pakhomova, J. Ondracek, M. Vindys and I. Stibor, *Z. Kristallogr.-New Cryst. Struct.*, 1997, **212**, 459.
- S. Kennedy and S. J. Dalgarno, *Chem. Commun.*, 2009, 5275; S. Kennedy, P. Cholewa, R. D. McIntosh and S. J. Dalgarno, *CrystEngComm*, 2013, **15**, 1520.
- For examples of recent reviews please see: A. Mukherjee, S. Tothadi and G. R. Desiraju, *Acc. Chem. Res.*, 2014, **47**, 2514; G. Cavallo, P. Metrangolo, R. Milani, T. Pilati, A. Priimagi, G. Resnati and G. Terraneo, *Chem. Rev.*, 2016, **116**, 2478; K. Rissanen, *Chem. Soc. Rev.*, 2017, **46**, 2638.
- Unfortunately it is not possible to isolate the entire series of analogues due to synthetic challenges, in particular the chloro derivatives. This would have been ideal for a full system analysis, but is unavoidable.
- To our knowledge it is not possible to isolate fully upper-rim halogenated analogues of **1**.
- For the synthesis of **2** see: O. Hudecek, P. Curinova, J. Budka and P. Lhoták, *Tetrahedron*, 2011, **67**, 5213.
- For the synthesis of **3** see: V. Stastny, P. Lhoták, V. Michlová, I. Stibor and J. Sykora, *Tetrahedron*, 2002, **67**, 7207.
- For the synthesis of **4** see: B. Klenke and W. Friedrichsen, *J. Chem. Soc., Perkin Trans. 1*, 1998, 3377.
- Correct as of 16/10/18, CSD version 5.39 (2018).
- a) M. Duta, Z. Asfari, A. Hagege, P. Thuery, M. Leroy, *Supramol. Chem.*, 2004, **16**, 205; b) A. N. Lazar, N. Dupont, A. Navaza, A. W. Coleman, *Chem. Commun.*, 2006, 1076; c) K. No, H. J. Lee, K. M. Park, S. S. Lee, K. H. Noh, S. K. Kim, J. Y. Lee, J. S. Kim, *J. Heterocyclic Chem.*, 2004, **41**, 211; d) S. Banthia, A. Samanta, *Org. Biomol. Chem.*, 2005, **3**, 1428; e) B. Bensenane, Z. Asfari, C. Platas-Iglesias, D. Esteban-Gómez, F. Djafri, M. Elhabiri, L. J. Charbonnière, *Dalton Trans.*, 2016, **45**, 15211; f) Z-G. Luo, Y. Zhao, C. Ma, L. Cao, S-H. Ai, J-S. Hu, X-M. Xu, *Jiegou Huaxue*, 2014, **33**, 1117; g) P. J. A. Kenis, E. G. Kerver, B. H. M. Snellink-Ruel, G. J. van Hummel, S. Harkema, M. C. Flipse, R. H. Woudenberg, J. F. J. Engbersen, D. N. Reinhoudt, *Eur. J. Org. Chem.*, 1998, 1089; h) P. Seigle-Ferrand, S. B. Sdira, C. Felix, R. Lamartine, C. Bavoux, B. Fenet, F. Bayard, F. Vocanson, *Mat. Sci. Eng. C*, 2006, **26**, 181; i) P. G. Jones, M. Freytag, *CSD Commun.*, 2010; j) S. Patra, R. Gunupuru, R. Lo, E. Suresh, B. Ganguly, P. Paul, *New J. Chem.*, 2012, **36**, 988; k) K. Takenaka, Y. Obora, Y. Tsuji, *Inorg. Chim. Acta*, 2004, **357**, 3895; l) F. Vita, M. Vorti, G. Orlandini, V. Zanichelli, C. Massera, F. Ugozzoli, A. Arduini, A. Secchi, *CrystEngComm*, 2016, **18**, 5017; m) M. Duta, Z. Asfari, P. Thuery, *CSD Commun.*, 2006; n) H. Halouani, I. Dumazet-Bonnamour, M. Perrin, R. Lamartine, *J. Org. Chem.*, 2004, **69**, 6521; o) C. Redshaw, O. Rowe, D. L. Hughes, A.-M. Fuller, I. A. Ibarra, S. M. Humphrey, *Dalton Trans.*, 2013, **42**, 1983; p) X. Zeng, X. Han, L. Chen, Q. Li, F. Xu, X. He, Z-Z. Zhang, *Tetrahedron Lett.*, 2002, **43**, 131; q) X. Zeng, L. Weng, L. Chen, F. Xu, Q. Li, X. Leng, X. He, Z-Z. Zhang, *Tetrahedron*, 2002, **58**, 2647; r) Z. Bo, Y-Z. Li, X-F. Lu, G-Y. Lu, *J. Chem. Crystallography*, 2005, **35**, 281; Z. Bo, C-Z. Zhang, G-Y. Lu, F. Liu, *Chinese J. Chem.*, 2006, **24**, 124; s) H. M. Chawla, S. P. Singh, S. Upreti, *Tetrahedron*, 2006, **62**, 9758; t) K. Stein, R. Schnorr, *CSD Commun*, 2015; u) P. Thuery, M. Lance, M. Nierlich, N. Reynier, V. Lamare, J. -F. Dozol, M. Saadioui, Z. Asfari, J. Vicens, *An. Quim.*, 1997, **93**, 324; v) M. Fehlinger, W. Abraham, *J. Incl. Phenom. Macrocyclic Chem.*, 2007, **58**, 263; w) P. Rashatasakhon, A. Jaiyu, R. Rojanathanes, N. Muangsin, N. Chaichit, M. Sukwattanasinitt, *J. Mol. Struct.*, 2010, **963**, 22; x) S. K. Kim, S. H. Kim, H. J. Kim, S. H. Lee, S. W. Lee, J. Ko, R. A. Bartsch, J. S. Kim, *Inorg. Chem.*, 2005, **44**, 786; y) D. Maity, M. Bhatt, M. L. Desai, E. Suresh, M. K. Si, V. P. Boricha, B. Ganguly, P. Paul, *Supramol. Chem.*, 2017, **29**, 600; z) S. Ullmann, R. Schnorr, C. Laube, B. Abel, B. Kersting, *Dalton Trans.*, 2018, **47**, 5801.
- a) S. Ullmann, R. Schnorr, M. Handke, C. Laube, B. Abel, J. Matysik, M. Findeisen, R. Rüger, T. Heine, B. Kersting, *Chem. Eur. J.*, 2017, **23**, 3824; b) A. Guelzim, S. Khriif, F. Baert, M. Saadioui, Z. Asfari, J. Vicens, *Acta Crystallogr. C*, 199, **53**, 1958; c) Z-Y. Xie, N-T. Hou, Y-Z. Zhu, H-B. Song, J-Y. Zheng, *Chem. Lett.*, 2008, **37**, 478; d) A. McConnell, C. Serpell, P. D. Beer, *New J. Chem.*, 2012, **36**, 102; e) H. M. Chawla, S. P. Singh, S. N. Sahu, S. Upreti, *Tetrahedron*, 2006, **62**, 7854; f) J. P. Ward, J. M. White, C. G. Young, *Tetrahedron*, 2013, **69**, 8824; g) V. Ramakrishna, S. Patra, E. Suresh, A. K. Bhatt, P. A. Bhatt, A. Hussain, P. Paul, *Inorg. Chem. Commun.*, 2012, **22**, 85; h) A. M. Reichwein, W. Verboom, S. Harkema, A. L. Spek, D. N. Reinhoudt, *J. Chem. Soc. Perkin Trans. 2*, 1994, 1167; i) N. Dupont, A. N. Lazar, F. Perret, O. Danylyuk, K. Suwinska, A. Navaza, A. W. Coleman, *CrystEngComm*, 2008, **10**, 975; j) H. Halouani, I. Dumazet-Bonnamour, C. Duchamp, C. Bavoux, N. Ehlinger, M. Perrin, R. Lamartine, *Eur. J. Org. Chem.*, 2002, 4202; k) P. Kuhn, D. Semeril, C. Jeunesse, D. Matt, P. J. Lutz, R. Louis, M. Neuburger, *Dalton Trans.*, 2006, 3647; l) S. J. Coles, C. W. Hall, M. B. Hursthouse, *Acta Cryst. C: Cryst. Struct. Commun.*, 2002, **58**, o29; m) F. Perret, A. N. Lazar, O. Shkurenko, K. Suwinska, N. Dupont, A. Navaza, A. W. Coleman, *CrystEngComm*, 2006, **8**, 890.
- J. J. McKinnon, M. A. Spackman and A. S. Mitchell, *Acta Cryst. B*, 2004, **60**, 627; M. A. Spackman and P. G. Byrom, *Chem. Phys. Lett.*, 1997, **267**, 215.
- a) L. Grubert, H. Henning, W. Abraham, *Tetrahedron*, 2009, **65**, 5936; b) D. R. Evans, M. Huang, J. C. Fettinger, T. L. Williams, *Inorg. Chem.*, 2002, **41**, 5986; c) J. Han, F.-L. Wang, Y.-X. Liu, F.-Y. Zhang, J.-B. Meng, Z.-J. He, *ChemPlusChem*, 2012, **77**, 196; d) K. Takenaka, Y. Obora, L.-H. Jiang, Y. Tsuji, *Organometallics*, 2002, **21**, 1158.
- L. Liu, L. N. Zakharov, J. A. Golen, A. L. Rheingold, W. H. Watson, T. A. Hanna, *Inorg. Chem.*, 2006, **45**, 4247.
- L. J. Barbour, G. W. Orr, J. L. Atwood, *Chem. Commun.*, 1998, 1901; M. Osipov, Q. Chu, S. J. Geib, D. P. Curran, S. G. Weber, *Beilstein J. Org. Chem.*, 2008, **4**, 36; S. E. Matthews, V. Felix, M. G. B. Drew, P. D. Beer, *Org. Biomol. Chem.*, 2003, **1**, 1232; P. G. Jones, M. Freytag, *CSD Communication*, 2010; L. J. Barbour, G. W. Orr, J. L. Atwood, *Chem. Commun.*, 1997, 1439.
- F. Maharaj, D. C. Craig, M. L. Scudder, R. Bishop, N. Kumar, *J. Incl. Phenom. Macro.*, 2007, **59**, 17.
- Y.-H. Luo, X.-J. Hu, J.-K. Liu, H. Zhang, Y. Li, H.-J. Yang, R.-J. Want, *Z. Krist.-New Cryst. St.*, 2006, 221, 327.
- C. Gropp, T. Husch, N. Trapp, M. Reiher and F. Diederich, *J. Am. Chem. Soc.*, 2017, **139**, 12190.