



Heriot-Watt University  
Research Gateway

## Stochastic compressible Euler equations and inviscid limits

### Citation for published version:

Breit, D & Mensah, PR 2019, 'Stochastic compressible Euler equations and inviscid limits', *Nonlinear Analysis: Theory, Methods and Applications*, vol. 184, pp. 218-238. <https://doi.org/10.1016/j.na.2019.02.013>

### Digital Object Identifier (DOI):

[10.1016/j.na.2019.02.013](https://doi.org/10.1016/j.na.2019.02.013)

### Link:

[Link to publication record in Heriot-Watt Research Portal](#)

### Document Version:

Peer reviewed version

### Published In:

Nonlinear Analysis: Theory, Methods and Applications

### General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [open.access@hw.ac.uk](mailto:open.access@hw.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Stochastic compressible Euler equations and inviscid limits

Dominic Breit      Prince Romeo Mensah

January 31, 2019

Department of Mathematics, Heriot-Watt University  
Riccarton Edinburgh EH14 4AS, UK

## Abstract

We prove the existence of a unique local strong solution to the stochastic compressible Euler system with nonlinear multiplicative noise. This solution exists up to a positive stopping time and is strong in both the PDE and probabilistic sense. Based on this existence result, we study the inviscid limit of the stochastic compressible Navier–Stokes system. As the viscosity tends to zero, any sequence of finite energy weak martingale solutions converges to the compressible Euler system.

**Keywords:** Euler system, Navier–Stokes system, compressible fluids, stochastic forcing, local strong solutions, inviscid limit

## 1 Introduction

We consider a stochastic variant of the *compressible barotropic Euler system* describing the time evolution of the mass density  $\varrho$  and the bulk velocity  $\mathbf{u}$  of a fluid driven by a nonlinear multiplicative noise. The system of equations reads

$$d\varrho + \operatorname{div}_x(\varrho\mathbf{u}) dt = 0, \quad (1.1)$$

$$d(\varrho\mathbf{u}) + [\operatorname{div}_x(\varrho\mathbf{u} \otimes \mathbf{u}) + a\nabla_x\varrho^\gamma] dt = \mathbb{G}(\varrho, \varrho\mathbf{u})dW. \quad (1.2)$$

Here  $\gamma > 1$  denotes the adiabatic exponent,  $a > 0$  is the squared reciprocal of the Mach number (the ratio between average velocity and speed of sound). The driving process  $W$  is a cylindrical Wiener process defined on some probability space  $(\Omega, \mathfrak{F}, \mathbb{P})$  and the coefficient  $\mathbb{G}$  is generally nonlinear and satisfies suitable growth assumptions, see Section 2 for the precise set-up. In order to eliminate the well-known difficulties related to the behaviour of fluid flows near the boundary of the underlying domain but still consider a physically meaningful situation, we study (1.1)–(1.2) on the whole space  $\mathbb{R}^n$ . We complement (1.1)–(1.2) with the far field condition

$$\varrho(x) \rightarrow \bar{\varrho}, \quad \mathbf{u}(x) \rightarrow 0, \quad |x| \rightarrow \infty, \quad (1.3)$$

for some  $\bar{\varrho} > 0$ . The initial conditions are random variables

$$\varrho(0, \cdot) = \varrho_0, \quad \mathbf{u}(0, \cdot) = \mathbf{u}_0, \tag{1.4}$$

with sufficient spatial regularity specified later. The main interest is the three-dimensional case, but  $n = 1, 2$  are also included in our theory (and obviously higher but non-physical dimensions). We remark that in contrast to the incompressible system, the one-dimensional situation makes sense for (1.1)–(1.2). In fact, martingale solutions to (1.1)–(1.2) for  $n = 1$  are studied in [4].

Our main result concerning the system (1.1)–(1.4) is the existence of a unique *maximal strong pathwise solution*. This solution is strong in the analytical sense (i.e., equations (1.1)–(1.2) are satisfied pointwise) and strong in the probabilistic sense (i.e., it is defined on a given probability space). It exists up to the hypothetical blow-up of the  $W^{1,\infty}$ -norm of the velocity  $\mathbf{u}$ . The precise formulation is given in Definition 2.3. The existence of a *maximal strong pathwise solution*, being defined on a maximal (random) time interval, follows from an extension of a *local strong pathwise solution*, see Definition 2.2, which lives up to a suitable stopping time. The main statement can be found in Theorem 2.4. Corresponding results in the deterministic case are classical and we refer to [1] and [3].

As in the incompressible case, global existence and uniqueness is a famous open problem. The presence of noise does not seem to change the situation. As solutions to nonlinear hyperbolic systems are known to develop singularities in finite time, the question about global well-posedness in the class of weak solutions has been analysed extensively. This is based on the method of convex integration which has been developed in the context of fluid mechanics by De Lellis and Székelyhidi [18]. The non-uniqueness of global-in-time weak solutions to (1.1)–(1.2) has recently been shown in [9] proceeding similar result in the deterministic case, cf. [21].

In contrast to the compressible system (1.1)–(1.2), its incompressible counterpart has been studied extensively. There are numerous results about the two-dimensional situation, see [5, 6, 13, 14, 27]. First results in three dimensions (treated on the whole space) can be found in [35, 28]. Similar to our main theorem, the existence of a unique local strong solutions is shown, however, only additive noise is allowed. The general three-dimensional case (with slip boundary conditions in a bounded domain and with nonlinear multiplicative noise) has finally been studied recently in [25].

A main idea in our existence proof is to rewrite (1.1)–(1.2) as a symmetric hyperbolic system by formally dividing (1.2) by  $\varrho$  similarly to [8]. In order to make the general framework from [29] for these systems available, we cut the noise in the critical range (that is, if  $\varrho$  is large or close to zero). The main tool in the limit procedure is an abstract Cauchy lemma from [24], see Lemma 2.1. We remark that the method from [8], used in the analysis of the Navier–Stokes, can to a certain extent be applied to (1.1)–(1.2) at least if periodic boundary conditions are considered. It does not, however, yield continuity of density and velocity in time – an advantage of the approach in the present paper.

In our second main result we are concerned with the relationship between the Navier–Stokes and Euler equations. Viscous compressible fluids subject to stochastic forcing can be

described by the Navier–Stokes system

$$d\rho + \operatorname{div}_x(\rho \mathbf{u}) dt = 0, \quad (1.5)$$

$$d(\rho \mathbf{u}) + [\operatorname{div}_x(\rho \mathbf{u} \otimes \mathbf{u}) + a \nabla_x \rho^\gamma] dt = \operatorname{div}_x \mathbb{S}(\nabla_x \mathbf{u}) dt + \mathbb{G}(\rho, \rho \mathbf{u}) dW. \quad (1.6)$$

Here  $\mathbb{S}(\nabla_x \mathbf{u})$  is the viscous stress tensor for which we assume Newton’s rheological law

$$\mathbb{S}(\nabla_x \mathbf{u}) = \nu \left( \nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{n} \operatorname{div}_x \mathbf{u} \mathbb{I} \right) + \lambda \operatorname{div}_x \mathbf{u} \mathbb{I}, \quad \nu > 0, \lambda \geq 0. \quad (1.7)$$

The study of the system (1.5)–(1.7) was first initiated in [11] where the global-in-time existence of *finite energy weak martingale solutions* is shown. These solutions are weak in the analytical sense (derivatives only exists in the sense of distributions) and weak in the probabilistic sense (the probability space is an integral part of the solution) as well. Moreover, the time-evolution of the energy can be controlled in terms of its initial state. The results from [11] – limited to periodic boundary conditions – have been extended to the whole space in [32]. So, a comparison between (1.1)–(1.2) and (1.5)–(1.7) is possible. Our main result, stated in Theorem 2.10, shows that any sequence of *finite energy weak martingale solutions* to (1.5)–(1.7) converges locally in time to the unique strong solution of (1.1)–(1.2) as  $\lambda, \nu \rightarrow 0$ . A similar strategy has been employed in [7] in order to study the inviscid-incompressible limit (where in addition,  $a = \frac{1}{\varepsilon^2}$  with  $\varepsilon \rightarrow 0$  is considered, where the limit system is the incompressible Euler system). A major difference to [7] is the generality of the noise coefficients we can consider now. Due to the incompressibility constraint on the target system only linear noise can be considered in [7]. In contrast to this, in the compressible case we can allow the full generality for the noise for which the existence theory applies.

The main tool in our proof is the relative energy inequality from [7]. It allows to compare a *finite energy weak martingale solutions* to (1.5)–(1.7) with a set of smooth comparison functions – in this case, the local solution to (1.1)–(1.2). The concept of relative energy inequality has a long history starting with the pioneering work of Dafermos [16]. In the context of compressible Navier–Stokes equation, it has been introduced in [23] and has also been used to study the inviscid limit of the compressible Navier–Stokes system in the deterministic case, cf. [36].

## 2 Preliminaries and main result

We start by introducing some notations and basic facts used in the text. To begin, we fix an arbitrary large time horizon  $T > 0$ .

### 2.1 Analytic framework

We will define the Sobolev space  $W^{s,2}(\mathbb{R}^n)$  for  $s \in \mathbb{R}$  as the set of tempered distributions for which the norm

$$\|v\|_{W^{s,2}(\mathbb{R}^n)} = \left( \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{v}(\xi)|^2 d\xi \right)^{\frac{1}{2}} \quad (2.1)$$

defined in frequency space is finite. Here  $\hat{v}$  denotes the Fourier transform of  $v$ . To shorten notation, we will write  $\|\cdot\|_{s,2}$  for  $\|\cdot\|_{W^{s,2}(\mathbb{R}^n)}$ . The following estimates are standard in the Moser-type calculus and can be found e.g. in Majda [31, Proposition 2.1].

1. For  $u, v \in W^{s,2} \cap L^\infty(\mathbb{R}^n)$  and  $|\alpha| \leq s$  we have

$$\|\partial_x^\alpha(uv)\|_2 \leq c_s (\|u\|_\infty \|\nabla_x^s v\|_2 + \|v\|_\infty \|\nabla_x^s u\|_2). \quad (2.2)$$

2. For  $u \in W^{s,2}(\mathbb{R}^n)$ ,  $\nabla_x u \in L^\infty(\mathbb{R}^n)$ ,  $v \in W^{s-1,2} \cap L^\infty(\mathbb{R}^n)$  and  $|\alpha| \leq s$  we have

$$\|\partial_x^\alpha(uv) - u\partial_x^\alpha v\|_2 \leq c_s (\|\nabla_x u\|_\infty \|\nabla_x^{s-1} v\|_2 + \|v\|_\infty \|\nabla_x^s u\|_2). \quad (2.3)$$

3. Let  $u \in W^{s,2} \cap C(\mathbb{R}^n)$  and let  $F$  be an  $s$ -times continuously differentiable function on an open neighborhood of the compact set  $G = \text{range}[u]$ . Then we have for all  $1 \leq |\alpha| \leq s$ ,

$$\|\partial_x^\alpha F(u)\|_{L^2(\mathbb{R}^n)} \leq c_s \|\partial_u F\|_{C^{s-1}(G)} \|u\|_{L^\infty(\mathbb{R}^n)}^{|\alpha|-1} \|\partial_x^\alpha u\|_{L^2(\mathbb{R}^n)}. \quad (2.4)$$

## 2.2 Stochastic framework

The driving process  $W$  is a cylindrical Wiener process on a separable Hilbert space  $\mathfrak{U}$  defined on some stochastic basis  $(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P})$  with a complete, right-continuous filtration. More specifically,  $W$  is given by a formal expansion

$$W(t) = \sum_{k \geq 1} e_k \beta_k(t),$$

where  $\{\beta_k\}_{k \geq 1}$  is a family of mutually independent real-valued Brownian motions with respect to  $(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P})$  and  $\{e_k\}_{k \geq 1}$  is an orthonormal basis of  $\mathfrak{U}$ . To give the precise definition of the diffusion coefficient  $\mathbb{G}$ , consider  $\rho \in L^2(\mathbb{R}^n)$ ,  $\rho \geq 0$ ,  $\mathbf{q} \in L^2(\mathbb{R}^n)$  and define it as follows

$$\mathbb{G}(\rho, \mathbf{q})e_k = \mathbf{G}_k(\cdot, \rho(\cdot), \mathbf{q}(\cdot)).$$

We suppose that the coefficients  $\mathbf{G}_k : \mathbb{R}^n \times [0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  are  $C^s$ -functions that satisfy uniformly in  $x \in \mathbb{R}^n$

$$\mathbf{G}_k(\cdot, 0, 0) = 0, \quad (2.5)$$

$$|\nabla^l \mathbf{G}_k(\cdot, \cdot, \cdot)| \leq \alpha_k, \quad \sum_{k \geq 1} \alpha_k < \infty \quad \text{for all } l \in \{1, \dots, s\}, \quad (2.6)$$

with  $s \in \mathbb{N}$  specified below. Finally, we assume that the  $\mathbf{G}_k$ s are compactly supported, i.e. there is  $\mathcal{K} \Subset \mathbb{R}^n$  such that

$$\text{spt}(\mathbf{G}_k) \Subset \mathcal{K} \quad \text{for all } k \in \mathbb{N}. \quad (2.7)$$

This is also assumed in the Navier–Stokes case in view of the far field condition, cf. [32]. A typical example we have in mind is

$$\mathbf{G}_k(x, \rho, \mathbf{q}) = \mathbf{a}_k(x)\rho + \mathbb{A}_k(x)\mathbf{q}, \quad (2.8)$$

where  $\mathbf{a}_k : \mathbb{R}^n \rightarrow \mathbb{R}^n$  and  $\mathbb{A}_k : \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$  are smooth functions which are compactly supported. However, our analysis applies to general nonlinear coefficients  $\mathbf{G}_k$ .

Observe that if  $\varrho, \mathbf{q}$  are  $(\mathfrak{F}_t)$ -progressively measurable  $L^2(\mathbb{R}^n)$ -valued processes such that

$$\varrho \in L^2\left(\Omega \times [0, T]; L^2(\mathbb{R}^n)\right), \quad \mathbf{q} \in L^2\left(\Omega \times [0, T]; L^2(\mathbb{R}^n; \mathbb{R}^n)\right),$$

and  $\mathbb{G}$  satisfies (2.5) and (2.6), then the stochastic integral

$$\int_0^t \mathbb{G}(\varrho, \varrho \mathbf{u}) \, dW = \sum_{k \geq 1} \int_0^t \mathbf{G}_k(\cdot, \varrho, \varrho \mathbf{u}) \, d\beta_k$$

is a well-defined  $(\mathfrak{F}_t)$ -martingale ranging in  $L^2(\mathbb{R}^n; \mathbb{R}^n)$ . Finally, we define an auxiliary space  $\mathfrak{U}_0 \supset \mathfrak{U}$  via

$$\mathfrak{U}_0 = \left\{ v = \sum_{k \geq 1} \alpha_k e_k; \sum_{k \geq 1} \frac{\alpha_k^2}{k^2} < \infty \right\},$$

endowed with the norm

$$\|v\|_{\mathfrak{U}_0}^2 = \sum_{k \geq 1} \frac{\alpha_k^2}{k^2}, \quad v = \sum_{k \geq 1} \alpha_k e_k.$$

Note that the embedding  $\mathfrak{U} \hookrightarrow \mathfrak{U}_0$  is Hilbert-Schmidt. Moreover, trajectories of  $W$  are  $\mathbb{P}$ -a.s. in  $C([0, T]; \mathfrak{U}_0)$ , cf. [15].

Let us complete this section by presenting a technical tool to pass to the limit in sequences of local strong solutions (which typically exists only up to a stopping time). It originates from [24, Lemma 5.1] in an abstract setting. For the present form, we refer to [25, Lemma 7.1].

**Lemma 2.1.** *Let  $(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P})$  be a stochastic basis with a complete right-continuous filtration. Let  $(u_R)$  be a sequence of  $W^{s,2}(\mathbb{R}^n)$ -valued continuous stochastic processes adapted to  $(\mathfrak{F}_t)$ . For  $N > 1$  and  $T > 0$ , we define the sequence of stopping times*

$$\mathfrak{t}_{R,N} = \inf \left\{ t \geq 0 : \|u_R(t)\|_{s,2} \geq N + \|u_R(0)\|_{s,2} \right\} \wedge T.$$

Assume that we have

$$\lim_{R \rightarrow \infty} \sup_{L \geq R} \mathbb{E} \sup_{0 \leq t \leq \mathfrak{t}_{R,N} \wedge \mathfrak{t}_{L,N}} \|u_R(t) - u_L(t)\|_{s,2} = 0, \quad (2.9)$$

$$\lim_{\delta \rightarrow 0} \sup_R \mathbb{P} \left\{ \sup_{0 \leq t \leq \mathfrak{t}_{R,N} \wedge \delta} \|u_R(t)\|_{s,2} > \|u_R(0)\|_{s,2} + N - 1 \right\} = 0. \quad (2.10)$$

Then there is a stopping time  $\mathfrak{t}$  such that  $\mathfrak{t} > 0$   $\mathbb{P}$ -a.s. and a  $W^{s,2}(\mathbb{R}^n)$ -valued  $(\mathfrak{F}_t)$ -progressively measurable process  $u$  satisfying

$$u(\cdot \wedge \mathfrak{t}) \in C([0, T]; W^{s,2}(\mathbb{R}^n)) \quad \mathbb{P}\text{-a.s.}$$

such that

$$\sup_{0 \leq t \leq \mathfrak{t}} \|u_R(t) - u(t)\|_{s,2} = 0 \quad (2.11)$$

$\mathbb{P}$ -a.s. as  $R \rightarrow \infty$  (at least for a subsequence).

### 2.3 Compressible Euler equations

Let us first introduce the notion of local strong pathwise solution. Such a solution is strong in both the PDE and probabilistic sense but possibly exists only locally in time. To be more precise, system (1.1)–(1.2) will be satisfied pointwise (not only in the sense of distributions) on the given stochastic basis associated to the cylindrical Wiener process  $W$ .

**Definition 2.2** (Local strong pathwise solution). Let  $(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P})$  be a stochastic basis with a complete right-continuous filtration. Let  $W$  be an  $(\mathfrak{F}_t)$ -cylindrical Wiener process and let  $\bar{\varrho} > 0$ . Let  $(\varrho_0, \mathbf{u}_0)$  be a  $\bar{\varrho} + W^{s,2}(\mathbb{R}^n) \times W^{s,2}(\mathbb{R}^n)$ -valued  $\mathfrak{F}_0$ -measurable random variable, and let  $\mathbb{G}$  satisfy (2.5) and (2.6) for some  $s \in \mathbb{N}$ . A triplet  $(\varrho, \mathbf{u}, \mathfrak{t})$  is called a local strong pathwise solution to the system (1.1)–(1.4) provided

(a)  $\mathfrak{t}$  is an a.s. strictly positive  $(\mathfrak{F}_t)$ -stopping time;

(b) the density  $\varrho$  is a  $\bar{\varrho} + W^{s,2}(\mathbb{R}^n)$ -valued  $(\mathfrak{F}_t)$ -progressively measurable process satisfying

$$\varrho(\cdot \wedge \mathfrak{t}) > 0, \quad \varrho(\cdot \wedge \mathfrak{t}) \in C([0, T]; \bar{\varrho} + W^{s,2}(\mathbb{R}^n)) \quad \mathbb{P}\text{-a.s.};$$

(c) the velocity  $\mathbf{u}$  is a  $W^{s,2}(\mathbb{R}^n)$ -valued  $(\mathfrak{F}_t)$ -progressively measurable process satisfying

$$\mathbf{u}(\cdot \wedge \mathfrak{t}) \in C([0, T]; W^{s,2}(\mathbb{R}^n)) \quad \mathbb{P}\text{-a.s.};$$

(d) there holds  $\mathbb{P}$ -a.s.

$$\begin{aligned} \varrho(t \wedge \mathfrak{t}) &= \varrho_0 - \int_0^{t \wedge \mathfrak{t}} \operatorname{div}_x(\varrho \mathbf{u}) \, ds, \\ (\varrho \mathbf{u})(t \wedge \mathfrak{t}) &= \varrho_0 \mathbf{u}_0 - \int_0^{t \wedge \mathfrak{t}} \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) \, ds - \int_0^{t \wedge \mathfrak{t}} a \nabla_x \varrho^\gamma \, ds + \int_0^{t \wedge \mathfrak{t}} \mathbb{G}(\varrho, \varrho \mathbf{u}) \, dW, \end{aligned}$$

for all  $t \in [0, T]$ .

In the above definition, we have tacitly assumed that  $s$  is large enough in order to provide sufficient regularity for the strong solutions. Classical solutions require spatial derivatives of  $\mathbf{u}$  and  $\varrho$  to be continuous  $\mathbb{P}$ -a.s. This motivates the following definition.

**Definition 2.3** (Maximal strong pathwise solution). Fix a stochastic basis with a cylindrical Wiener process and an initial condition as in Definition 2.2. A quadruplet

$$(\varrho, \mathbf{u}, (\mathfrak{t}_R)_{R \in \mathbb{N}}, \mathfrak{t})$$

is a maximal strong pathwise solution to system (1.1)–(1.4) provided

- (a)  $\mathfrak{t}$  is an a.s. strictly positive  $(\mathfrak{F}_t)$ -stopping time;
- (b)  $(\mathfrak{t}_R)_{R \in \mathbb{N}}$  is an increasing sequence of  $(\mathfrak{F}_t)$ -stopping times such that  $\mathfrak{t}_R < \mathfrak{t}$  on the set  $[\mathfrak{t} < T]$ ,  $\lim_{R \rightarrow \infty} \mathfrak{t}_R = \mathfrak{t}$  a.s. and

$$\sup_{t \in [0, \mathfrak{t}_R]} \|\mathbf{u}(t)\|_{1, \infty} \geq R \quad \text{on} \quad [\mathfrak{t} < T]; \quad (2.12)$$

- (c) each triplet  $(\varrho, \mathbf{u}, \mathfrak{t}_R)$ ,  $R \in \mathbb{N}$ , is a local strong pathwise solution in the sense of Definition 2.2.

The notion of a maximal pathwise solution has already appeared in the literature in the context of various SPDE or SDE models, see for instance [12, 20, 26, 34]. Finally, we have all in hand to formulate our main result.

**Theorem 2.4.** *Let  $s \in \mathbb{N}$  satisfy  $s > \frac{n}{2} + 2$  and let  $\bar{\varrho} > 0$ . Let the coefficients  $\mathbf{G}_k$  satisfy hypotheses (2.5), (2.6) and let  $(\varrho_0, \mathbf{u}_0)$  be an  $\mathfrak{F}_0$ -measurable,  $\bar{\varrho} + W^{s,2}(\mathbb{R}^n) \times W^{s,2}(\mathbb{R}^n)$ -valued random variable such that  $\varrho_0 > 0$   $\mathbb{P}$ -a.s. Then there exists a unique maximal strong pathwise solution  $(\varrho, \mathbf{u}, (\mathfrak{t}_R)_{R \in \mathbb{N}}, \mathfrak{t})$  to problem (1.1)–(1.4) in the sense of Definition 2.3 with the initial condition  $(\varrho_0, \mathbf{u}_0)$ .*

**Remark 2.5.** Starting with the pioneering work in [17], several counterexamples have been developed showing that the (deterministic) compressible Euler system is desperately ill-posed. Even if the initial data is smooth, the global existence and uniqueness of solutions can fail. Similar result for the stochastic compressible Euler system have been achieved recently in [9]. The existence of global strong solutions (i.e. the stopping time  $\mathfrak{t}$  in Definition 2.2 reaches  $T$ ) is not expected.

## 2.4 Compressible Navier–Stokes equations

In this section, we present the concept of finite energy weak martingale solutions to (1.5)–(1.7). It has been introduced in [11] and improved in [7] and [10]. Both papers complement (1.5)–(1.7) with periodic boundary conditions. A corresponding version on the whole space can be found in [32]. These solutions are weak in the analytical sense (derivatives only exists in the sense of distributions) and weak in the probabilistic sense (the probability space is an integral part of the solution) as well. Moreover, the time-evolution of the energy can be controlled in terms of its initial state. They exists globally in time.

**Definition 2.6** (Finite energy weak martingale solution). Let  $\bar{\rho} > 0$ . Let  $\Lambda$  be a Borel probability measure on  $L_{loc}^\gamma(\mathbb{R}^n) \times L_{loc}^{2\gamma/\gamma+1}(\mathbb{R}^n)$ . Then  $[(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P}); \rho, \mathbf{v}, W]$  is a finite energy weak martingale solution of (1.5)–(1.7) if

- (a)  $(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P})$  is a stochastic basis with a complete right-continuous filtration,
- (b)  $W$  is a  $(\mathfrak{F}_t)$ -cylindrical Wiener process,



- (c) the density  $\rho$  satisfies  $\rho \geq 0$ ,  $t \mapsto \langle \rho(t, \cdot), \phi \rangle \in C[0, T]$  for any  $\phi \in C_c^\infty(\mathbb{R}^n)$   $\mathbb{P}$ -a.s., the function  $t \mapsto \langle \rho(t, \cdot), \phi \rangle$  is progressively measurable and

$$\mathbb{E} \left[ \sup_{t \in (0, T)} \|\rho(t, \cdot)\|_{L^\gamma(K)}^p \right] < \infty$$

for all  $1 \leq p < \infty$  and all  $K \Subset \mathbb{R}^n$ ,

- (d) the velocity field  $\mathbf{v}$  is  $(\mathfrak{F}_t)$ -adapted and

$$\mathbb{E} \left[ \int_0^T \|\mathbf{v}\|_{W^{1,2}(K)}^2 dt \right]^p < \infty$$

for all  $1 \leq p < \infty$  and all  $K \Subset \mathbb{R}^n$ ,

- (e) the momentum  $\rho \mathbf{v}$  satisfies  $t \mapsto \langle \rho \mathbf{v}, \varphi \rangle \in C[0, T]$  for any  $\varphi \in C_c^\infty(\mathbb{R}^n)$   $\mathbb{P}$ -a.s., the function  $t \mapsto \langle \rho \mathbf{v}, \varphi \rangle$  is progressively measurable and

$$\mathbb{E} \left[ \sup_{t \in (0, T)} \|\rho \mathbf{v}\|_{L^{\frac{2\gamma}{\gamma+1}}(K)}^p \right] < \infty$$

for all  $1 \leq p < \infty$  and all  $K \Subset \mathbb{R}^n$ ,

- (f)  $\Lambda = \mathbb{P} \circ [\rho(0), \rho \mathbf{v}(0)]^{-1}$ ,

- (g) for all  $\phi \in C_c^\infty(\mathbb{R}^n)$  and  $\varphi \in C_c^\infty(\mathbb{R}^n)$  we have

$$\begin{aligned} \langle \rho(t), \phi \rangle &= \langle \rho(0), \phi \rangle - \int_0^t \langle \rho \mathbf{v}, \nabla_x \phi \rangle ds \\ \langle \rho \mathbf{v}(t), \varphi \rangle &= \langle (\rho \mathbf{v})(0), \varphi \rangle - \int_0^t \langle \rho \mathbf{v} \otimes \mathbf{v}, \nabla_x \varphi \rangle ds + \int_0^t \langle \mathbb{S}(\nabla_x \mathbf{v}), \nabla_x \varphi \rangle ds \\ &\quad - \int_0^t \langle a \rho^\gamma, \operatorname{div}_x \varphi \rangle ds + \int_0^t \langle \mathbb{G}(\rho, \rho \mathbf{v}), \varphi \rangle dW \end{aligned} \quad (2.13)$$

$\mathbb{P}$ -a.s. for all  $t \in [0, T]$ ,

- (h) the energy inequality

$$\begin{aligned} &\int_{\mathbb{R}^n} \left[ \frac{1}{2} \rho |\mathbf{v}|^2 + H(\rho, \bar{\rho}) \right] dx + \int_0^t \int_{\mathbb{R}^n} \mathbb{S}(\nabla_x \mathbf{v}) : \nabla_x \mathbf{v} dx ds \\ &\leq \int_{\mathbb{R}^n} \left[ \frac{1}{2} \frac{|\rho \mathbf{v}(0)|^2}{\rho(0)} + H(\rho(0), \bar{\rho}) \right] dx + \sum_{k=1}^{\infty} \int_0^t \left( \int_{\mathbb{R}^n} \mathbf{G}_k(\rho, \rho \mathbf{v}) \cdot \mathbf{v} dx \right) dW_k \\ &\quad + \frac{1}{2} \sum_{k=1}^{\infty} \int_0^t \int_{\mathbb{R}^n} \rho^{-1} |\mathbf{G}_k(\rho, \rho \mathbf{v})|^2 ds \end{aligned} \quad (2.14)$$

holds for a.e.  $t \in [0, T]$   $\mathbb{P}$ -a.s. Here  $M_E$  is a real-valued square integrable martingale and  $H(\rho, \bar{\rho})$  is the pressure potential given by

$$H(\rho, \bar{\rho}) = \frac{a}{\gamma - 1} \left( \rho^\gamma - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}) - \bar{\rho}^\gamma \right). \quad (2.15)$$

The following existence theorem is shown in [32, Theorem 1] (see also [33, Ch. 3] for a more detailed proof).

**Theorem 2.7.** *Let  $\bar{\rho} > 0$ ,  $\gamma > \frac{n}{2}$  and assume that  $\Lambda$  is a Borel probability measure on  $L_{loc}^\gamma(\mathbb{R}^n) \times L_{loc}^{2\gamma/\gamma+1}(\mathbb{R}^n)$  such that*

$$\Lambda \left\{ (\rho, \mathbf{q}) \in L_{loc}^\gamma(\mathbb{R}^n) \times L_{loc}^{2\gamma/\gamma+1}(\mathbb{R}^n) : 0 < M_1 \leq \rho \leq M_2, \mathbf{q}|_{\{\rho=0\}} = 0, \right\} = 1$$

with constants  $0 < M_1 < M_2$ . Furthermore, assume that the following moment estimate

$$\int_{L_x^\gamma \times L_x^{2\gamma/\gamma+1}} \left\| \frac{1}{2} \frac{|\mathbf{q}|^2}{\rho} + H(\rho, \bar{\rho}) \right\|_{L_x^1}^p d\Lambda(\rho, \mathbf{q}) < \infty \quad (2.16)$$

holds for all  $1 \leq p < \infty$ . Finally, assume that (2.5), (2.6) with  $s = 1$  and (2.7) hold. Then there exists a finite energy weak martingale solution of (1.5)–(1.7) in the sense of Definition 2.6 with initial law  $\Lambda$ .

**Remark 2.8.** Due to the assumptions on the initial law as well as (2.5), (2.6) with  $s = 1$  and (2.7) the energy inequality (2.14) implies the following moment estimates

$$\begin{aligned} \mathbb{E} \left[ \int_0^T \int_{\mathbb{R}^n} |\nabla \mathbf{v}|^2 dx dt \right]^p &< \infty, & \mathbb{E} \left[ \int_0^T \int_{\mathbb{R}^n} \rho |\mathbf{v}|^2 dx dt \right]^p &< \infty, \\ \mathbb{E} \left[ \int_0^T \int_{\mathbb{R}^n} \left( |\rho - \bar{\rho}|^2 \mathbb{I}_{|\rho - \bar{\rho}| \leq 1} + |\rho - \bar{\rho}|^\gamma \mathbb{I}_{|\rho - \bar{\rho}| > 1} \right) dx dt \right]^p &< \infty, \end{aligned}$$

for all  $1 \leq p < \infty$ . Note that the last integrand is a lower bound for  $H(\rho, \bar{\rho})$  due to the elementary inequality

$$H(\rho, r) \geq c(r) \begin{cases} |\rho - r|^2 & : \text{if } r/2 \leq \rho \leq 2r, \\ 1 + \rho^\gamma & : \text{else,} \end{cases} \quad (2.17)$$

which holds for any  $r > 0$ .

**Remark 2.9.** Although Theorem 2.7 was shown in [32] for  $n = 3$ , it also applies in general dimensions replacing the bound  $\gamma > \frac{3}{2}$  by  $\gamma > \frac{n}{2}$ .

## 2.5 Inviscid limit

In this section we give the main result concerning the relation of the systems (1.1)–(1.2) and (1.5)–(1.7). From a formal point of view, the Navier–Stokes system (1.5)–(1.7) converges to the Euler system (1.1)–(1.2) if  $\nu, \lambda \rightarrow 0$ . In order to make this idea rigorous, we have to analyse a singular limit. Singular limit arguments for analysing the interactions between fluid dynamic models arises from suitable change of variables in time and space or by using dimensional analysis. Such transformations are now standard and interested readers can refer to [2, 30] and the references within for further information.

To study the inviscid limit result for (1.5)–(1.7), we are interested in the transformation that leads to the following mappings:

$$\varrho \mapsto \rho_\varepsilon, \quad \mathbf{u} \mapsto \mathbf{v}_\varepsilon, \quad \nu \mapsto \varepsilon\nu, \quad \lambda \mapsto \varepsilon\lambda.$$

This yields the system

$$\begin{aligned} d\rho_\varepsilon + \operatorname{div}(\rho_\varepsilon \mathbf{v}_\varepsilon) dt &= 0, \\ d(\rho_\varepsilon \mathbf{v}_\varepsilon) + \left[ \operatorname{div}(\rho_\varepsilon \mathbf{v}_\varepsilon \otimes \mathbf{v}_\varepsilon) + a \nabla \rho_\varepsilon^\gamma \right] dt &= \varepsilon \operatorname{div} \mathbb{S}(\nabla \mathbf{v}_\varepsilon) dt + \mathbb{G}(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon) dW, \end{aligned} \quad (2.18)$$

where the parameter  $\varepsilon \in (0, 1]$  corresponds to the inverse of the *Reynolds number*. Our aim is to pass to the limit  $\varepsilon \rightarrow 0$ . We consider the following ill-prepared data that connects the inputs of Navier–Stokes and Euler system. We assume that the initial data  $(\rho_{0,\varepsilon}, \mathbf{v}_{0,\varepsilon})$  of the system (2.18) satisfy the following conditions

$$\mathbb{E} \int_{\mathbb{R}^n} H(\rho_{0,\varepsilon}, \bar{\varrho}) dx < \infty, \quad \rho_{0,\varepsilon} |\mathbf{v}_{0,\varepsilon}|^2 \in L^1(\mathbb{R}^n), \quad 0 < \varrho^- \leq \rho_{0,\varepsilon} \leq \varrho^+ \quad \mathbb{P}\text{-a.s.}, \quad (2.19)$$

where  $\varrho^-$  and  $\varrho^+$  are independent of  $\varepsilon$ . The initial data  $(\varrho_0, \mathbf{u}_0)$  of the limit system (1.1)–(1.2) satisfy

$$(\varrho_0, \mathbf{u}_0) \in \bar{\varrho} + W^{s,2}(\mathbb{R}^n) \times W^{s,2}(\mathbb{R}^n), \quad \varrho_0 \geq \varrho^- > 0 \quad \mathbb{P}\text{-a.s.} \quad (2.20)$$

Finally, we suppose that

$$\mathbb{E} \int_{\mathbb{R}^n} H(\rho_{0,\varepsilon}, \varrho_0) dx \xrightarrow{\varepsilon \searrow 0} 0, \quad \mathbb{E} \int_{\mathbb{R}^n} |\mathbf{v}_{0,\varepsilon} - \mathbf{u}_0|^2 dx \xrightarrow{\varepsilon \searrow 0} 0. \quad (2.21)$$

Our main result reads as follows.

**Theorem 2.10.** *Let  $\bar{\varrho} > 0$  be given and suppose that (2.5)–(2.7) hold with  $s > \frac{n}{2} + 2$ . Let  $(\Omega, \mathfrak{F}, \mathbb{P})$  be a complete probability space and  $W$  a cylindrical Wiener process on  $(\Omega, \mathfrak{F}, \mathbb{P})$ . Assume that*

$$\left[ (\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P}), \rho_\varepsilon, \mathbf{v}_\varepsilon, W \right]_{\varepsilon > 0} \quad (2.22)$$

*is a family of finite energy weak martingale solution to the system (2.18) in the sense of Definition 2.6 with  $\bar{\rho} = \bar{\varrho} > 0$ . On the same stochastic basis  $(\Omega, \mathfrak{F}, (\mathfrak{F}_t), \mathbb{P})$ , consider the unique maximal strong pathwise solution to the Euler system (1.1)–(1.2) given by  $(\varrho, \mathbf{u}, (\mathbf{t}_R)_{R \in \mathbb{N}}, \mathbf{t})$  driven by the same cylindrical Wiener process  $W$ . Assume that the initial data  $(\rho_{0,\varepsilon}, \mathbf{v}_{0,\varepsilon})$  and  $(\varrho_0, \mathbf{u}_0)$  are  $\mathfrak{F}_0$ -measurable and satisfies (2.19)–(2.21). Then we have*

$$\sup_{t \in (0, T)} \mathbb{E} \int_{\mathbb{R}^n} \left[ \frac{1}{2} \rho_\varepsilon |\mathbf{v}_\varepsilon - \mathbf{u}|^2 + H(\rho_\varepsilon, \varrho) \right] (t \wedge \mathbf{t}_R, \cdot) dx \rightarrow 0 \quad (2.23)$$

*as  $\varepsilon \rightarrow 0$  for all  $R \in \mathbb{N}$ .*

Before we proof Theorem 2.10, we remark that (2.23) implies that we have

$$\rho_\varepsilon(\cdot \wedge \mathbf{t}_R) \rightarrow \varrho(\cdot \wedge \mathbf{t}_R) \text{ in } L^{\bar{\gamma}}(\Omega \times (0, T); L^{\bar{\gamma}}_{loc}(\mathbb{R}^n)), \quad (2.24)$$

$$(\rho_\varepsilon \mathbf{v}_\varepsilon)(\cdot \wedge \mathbf{t}_R) \rightarrow (\varrho \mathbf{u})(\cdot \wedge \mathbf{t}_R) \text{ in } L^{\frac{2\bar{\gamma}}{\bar{\gamma}+1}}(\Omega \times (0, T); L^{\frac{2\bar{\gamma}}{\bar{\gamma}+1}}_{loc}(\mathbb{R}^n)), \quad (2.25)$$

where  $\bar{\gamma} = \min\{2, \gamma\}$ . The convergence (2.24) follows from inequality (2.17) whilst (2.25) follows from the identity

$$\rho_\varepsilon \mathbf{v}_\varepsilon - \varrho \mathbf{u} = (\rho_\varepsilon - \varrho) \mathbf{u} + \sqrt{\rho_\varepsilon} \sqrt{\rho_\varepsilon} (\mathbf{v}_\varepsilon - \mathbf{u})$$

and Hölder's inequality, cf. [36].

### 3 Proof of Theorem 2.4

#### 3.1 Approximation

On a formal level, it can be seen that (1.1)–(1.2) is equivalent to

$$dr + \mathbf{u} \cdot \nabla_x r \, dt + \frac{\gamma - 1}{2} r \operatorname{div}_x \mathbf{u} \, dt = 0, \quad (3.1)$$

$$d\mathbf{u} + [\mathbf{u} \cdot \nabla_x \mathbf{u} + r \nabla_x r] \, dt = \mathbb{F}(r, \mathbf{u}) dW, \quad (3.2)$$

where

$$\varrho(r) = \left( \frac{\gamma - 1}{2a\gamma} r^2 \right)^{\frac{1}{\gamma-1}}, \quad \mathbb{F}(r, \mathbf{u}) = \frac{1}{\varrho(r)} \mathbb{G}(\varrho(r), \varrho(r) \mathbf{u}),$$

cf. [8, Sec. 2.4]. This can be made rigorous as long as  $r$  (or equivalently,  $\varrho$ ) is strictly positive. As we will solve (1.1)–(1.2) with respect to far field conditions, we seek a solution  $r \in \bar{r} + W^{s,2}(\mathbb{R}^n)$ , where  $\bar{r} = \sqrt{\frac{2a\gamma}{\gamma-1} \varrho^{\frac{\gamma-1}{2}}}$ . In order to do so, we set  $\hat{r} := r - \bar{r}$  and aim to solve

$$d\hat{r} + \mathbf{u} \cdot \nabla_x \hat{r} \, dt + \frac{\gamma - 1}{2} (\hat{r} + \bar{r}) \operatorname{div}_x \mathbf{u} \, dt = 0, \quad (3.3)$$

$$d\mathbf{u} + [\mathbf{u} \cdot \nabla_x \mathbf{u} + (\hat{r} + \bar{r}) \nabla_x \hat{r}] \, dt = \hat{\mathbb{F}}(\hat{r}, \mathbf{u}) dW, \quad (3.4)$$

where  $\hat{\mathbb{F}}(\hat{r}, \mathbf{u}) = \mathbb{F}(\hat{r} + \bar{r}, \mathbf{u})$ . We remark that the left-hand side of (3.4) corresponds to a symmetric hyperbolic system, cf. Majda [31]. In the stochastic case such system have been studied in [29]. Unfortunately, the result from [29] does not apply to the general assumptions on  $\mathbb{G}$  we have in mind. In fact, the assumptions on the noise coefficient  $\mathbb{F}$  are violated for small (close to zero) or large values of  $r$ . Due to this we replace  $\mathbb{F}$  by

$$\mathbb{F}_R(r, \mathbf{u}) = \frac{1}{\varrho(r)} \varphi_R(\varrho(r)) \varphi_R(\varrho(r)^{-1}) \mathbb{G}(\varrho(r), \varrho(r) \mathbf{u}),$$

where  $\varphi_R : [0, \infty) \rightarrow [0, 1]$  are smooth cut-off functions satisfying

$$\varphi_R(y) = \begin{cases} 1, & 0 \leq y \leq R, \\ 0, & R + 1 \leq y, \end{cases} \quad (3.5)$$

and similarly  $\hat{\mathbb{F}}_R(\hat{r}, \mathbf{u}) = \mathbb{F}_R(\hat{r} + \bar{r}, \mathbf{u})$ . We now study the system

$$d\hat{r} + \mathbf{u} \cdot \nabla_x \hat{r} dt + \frac{\gamma - 1}{2} (\hat{r} + \bar{r}) \operatorname{div}_x \mathbf{u} dt = 0, \quad (3.6)$$

$$d\mathbf{u} + [\mathbf{u} \cdot \nabla_x \mathbf{u} + (\hat{r} + \bar{r}) \nabla_x \hat{r}] dt = \hat{\mathbb{F}}_R(\hat{r}, \mathbf{u}) dW. \quad (3.7)$$

We assume for the moment that

$$\|(\varrho_0, \mathbf{u}_0)\|_{W^{s,2}(\mathbb{R}^n)} < M, \quad \varrho_0 > \frac{1}{M} \text{ } \mathbb{P}\text{-a.s.} \quad (3.8)$$

for some deterministic constant  $M > 0$ . These assumptions will be relaxed later. By definition of  $\mathbb{F}_R$ , the noise disappears if  $\varrho(r)$  is larger than  $R+1$  or smaller than  $\frac{1}{R+1}$ . Consequently, (2.5) and (2.6) imply that  $\mathbb{F}_R$  (and  $\hat{\mathbb{F}}_R$ ) is globally Lipschitz continuous on  $W^{s,2}(\mathbb{R}^n) \times W^{s,2}(\mathbb{R}^n)$  for any fixed  $R$ . By [29, Thm. 1.2] (where  $u$  takes the role of  $(\hat{r}, \mathbf{u})$ ), there is a unique strong solution  $(\hat{r}_R, \mathbf{u}_R, \mathfrak{t}_R)$  to (3.6)–(3.7) in the following sense:<sup>1</sup>

- (i)  $(\hat{r}_R, \mathbf{u}_R)$  is a  $W^{s,2}(\mathbb{R}^n) \times W^{s,2}(\mathbb{R}^n)$ -valued right-continuous progressively measurable process;
- (ii)  $\mathfrak{t}_R$  is a stopping time with respect to  $(\mathfrak{F}_t)$  such that  $\mathbb{P}$ -a.s.

$$\mathfrak{t}_R = \lim_{N \rightarrow \infty} \mathfrak{t}_{R,N} \quad (3.9)$$

where

$$\mathfrak{t}_{R,N} = \inf \{0 \leq t < \infty : \|(\hat{r}_R, \mathbf{u}_R)(t)\|_{s,2} \geq N\} \quad (3.10)$$

with the convention that  $\mathfrak{t}_{R,N} = \infty$  if the set above is empty;

- (iii) there holds  $\mathbb{P}$ -a.s.

$$(\hat{r}_R, \mathbf{u}_R)(\cdot \wedge \mathfrak{t}_R) \in C([0, T]; W^{s,2}(\mathbb{R}^n) \times W^{s,2}(\mathbb{R}^n))$$

as well as

$$\begin{aligned} \hat{r}(t \wedge \mathfrak{t}_{R,N}) &= r_0 - \int_0^{t \wedge \mathfrak{t}_{R,N}} \mathbf{u} \cdot \nabla_x \hat{r} ds - \int_0^{t \wedge \mathfrak{t}_{R,N}} \frac{\gamma - 1}{2} (\hat{r} + \bar{r}) \operatorname{div}_x \mathbf{u} ds, \\ \mathbf{u}(t \wedge \mathfrak{t}_{R,N}) &= \mathbf{u}_0 - \int_0^{t \wedge \mathfrak{t}_{R,N}} \mathbf{u} \cdot \nabla_x \mathbf{u} ds - \int_0^{t \wedge \mathfrak{t}_{R,N}} (\hat{r} + \bar{r}) \nabla_x \hat{r} ds + \int_0^{t \wedge \mathfrak{t}_{R,N}} \hat{\mathbb{F}}_R(\hat{r}, \mathbf{u}) dW, \end{aligned}$$

for all  $t \in [0, T]$  and all  $N \geq 1$ , where  $r_0 = \sqrt{\frac{2a\gamma}{\gamma-1}} \varrho_0^{\frac{\gamma-1}{2}} - \bar{r}$ .

---

<sup>1</sup>The result from [29] requires the assumption  $s > \frac{n}{2} + 2$ .

Given  $(\hat{r}_R, \mathbf{u}_R)$  we see that  $(r_R, \mathbf{u}_R) = (\hat{r}_R + \bar{r}, \mathbf{u}_R)$  solves  $\mathbb{P}$ -a.s.

$$dr_R + \mathbf{u}_R \cdot \nabla_x r_R \, dt + \frac{\gamma-1}{2} r_R \operatorname{div}_x \mathbf{u}_R \, dt = 0, \quad (3.11)$$

$$d\mathbf{u}_R + [\mathbf{u}_R \cdot \nabla_x \mathbf{u}_R + r_R \nabla_x r_R] \, dt = \mathbb{F}_R(r_R, \mathbf{u}_R) dW. \quad (3.12)$$

The aim in the following is to pass to the limit  $R \rightarrow \infty$  in (3.11) and (3.12). This will be done by verifying the assumptions of Lemma 2.1. We start by showing certain a priori estimates as a consequence of which we obtain (2.10). Eventually we show uniqueness of (3.1)–(3.2) which implies (2.9).

### 3.2 A priori estimates

We immediately see that  $(\varrho_R, \mathbf{u}_R) := \left( \left( \frac{\gamma-1}{2\alpha\gamma} r_R^2 \right)^{\frac{1}{\gamma-1}}, \mathbf{u}_R \right)$  solves  $\mathbb{P}$ -a.s.

$$d\varrho_R + \operatorname{div}_x(\varrho_R \mathbf{u}_R) \, dt = 0 \quad (3.13)$$

with  $\varrho_R(0) = \varrho_0$  and far field  $\bar{\varrho}$ . The standard maximum principle (see, e.g., [19, eq. (7)] or [22, eq. (2.8)]) applied to (3.13) yields

$$\inf_{\mathbb{R}^n} \varrho_0 \exp \left( - \int_0^{t_{R,N}} \|\operatorname{div}_x \mathbf{u}_R\|_\infty ds \right) \leq \varrho_R(t, x) \leq \sup_{\mathbb{R}^n} \varrho_0 \exp \left( \int_0^{t_{R,N}} \|\operatorname{div}_x \mathbf{u}_R\|_\infty ds \right) \quad (3.14)$$

$\mathbb{P}$ -a.s. for all  $(t, x) \in (0, t_{R,N}) \times \mathbb{R}^n$ . Consequently, the definition of  $t_{R,N}$ , the embedding  $W^{s,2}(\mathbb{R}^n) \hookrightarrow C^{1,\alpha}(\mathbb{R}^n)$  (for  $s > \frac{n}{2} + 1$  and some  $\alpha > 0$ ), as well as (3.8) implies that

$$c_N^{-1} \inf_{\mathbb{R}^n} \varrho_0 \leq \varrho_R(t, x) \leq c_N \sup_{\mathbb{R}^n} \varrho_0 \quad (3.15)$$

$\mathbb{P}$ -a.s. for all  $(t, x) \in (0, t_{R,N}) \times \mathbb{R}^n$  and for a positive constant  $c_N = c_N(\bar{\varrho}, M)$  which is independent of  $R$ . For a given  $N \in \mathbb{N}$  there is  $R_N$  such that (3.15) implies that

$$R_N^{-1} \leq \varrho_{R_N} \leq R_N \quad (3.16)$$

$\mathbb{P}$ -a.s. in  $(0, t_{R_N, N}) \times \mathbb{R}^n$ .

Let  $\alpha$  be a multiindex such that  $|\alpha| \leq s$ . Differentiating (3.11) in the  $x$ -variable, we obtain

$$\begin{aligned} d\partial_x^\alpha r_R + \left[ \mathbf{u}_R \cdot \nabla_x \partial_x^\alpha r_R + \frac{\gamma-1}{2} r_R \operatorname{div}_x \partial_x^\alpha \mathbf{u}_R \right] dt \\ = \left[ \mathbf{u}_R \cdot \partial_x^\alpha \nabla_x r_R - \partial_x^\alpha (\mathbf{u}_R \cdot \nabla_x r_R) \right] dt \\ + \frac{\gamma-1}{2} \left[ r_R \partial_x^\alpha \operatorname{div}_x \mathbf{u}_R - \partial_x^\alpha (r_R \operatorname{div}_x \mathbf{u}_R) \right] dt \\ =: T_1^R dt + T_2^R dt. \end{aligned} \quad (3.17)$$

Similarly, we differentiate (3.12) and deduce that

$$\begin{aligned}
& d\partial_x^\alpha \mathbf{u}_R + [\mathbf{u}_R \cdot \nabla_x \partial_x^\alpha \mathbf{u}_R + r_R \nabla_x \partial_x^\alpha r_R] dt \\
&= [\mathbf{u}_R \cdot \partial_x^\alpha \nabla_x \mathbf{u}_R - \partial_x^\alpha (\mathbf{u}_R \cdot \nabla_x \mathbf{u}_R)] dt \\
&+ [r_R \partial_x^\alpha \nabla_x r_R - \partial_x^\alpha (r_R \nabla_x r_R)] dt \\
&+ \partial_x^\alpha \mathbb{F}_R(r_R, \mathbf{u}_R) dW \\
&=: T_3^R dt + T_4^R dt + \partial_x^\alpha \mathbb{F}_R(r_R, \mathbf{u}_R) dW.
\end{aligned} \tag{3.18}$$

It follows from (2.3) that the ‘‘error’’ terms may be handled as

$$\begin{aligned}
\|T_1^R\|_2 &\lesssim \left[ \|\nabla_x \mathbf{u}_R\|_\infty \|\nabla_x^s r_R\|_2 + \|\nabla_x r_R\|_\infty \|\nabla_x^s \mathbf{u}_R\|_2 \right], \\
\|T_2^R\|_2 &\lesssim \left[ \|\nabla_x r_R\|_\infty \|\nabla_x^s \mathbf{u}_R\|_2 + \|\operatorname{div}_x \mathbf{u}_R\|_\infty \|\nabla_x^s r_R\|_2 \right], \\
\|T_3^R\|_2 &\lesssim \|\nabla_x \mathbf{u}_R\|_\infty \|\nabla_x^s \mathbf{u}_R\|_2, \\
\|T_4^R\|_2 &\lesssim \|\nabla_x r_R\|_\infty \|\nabla_x^s r_R\|_2.
\end{aligned} \tag{3.19}$$

Multiplying (3.17) by  $\partial_x^\alpha (r_R - \bar{r})$ , we observe

$$\begin{aligned}
& \|\partial_x^\alpha (r_R(t) - \bar{r})\|_2^2 + (\gamma - 1) \int_0^t \int_{\mathbb{R}^n} r_R \operatorname{div}_x \partial_x^\alpha \mathbf{u}_R \partial_x^\alpha (r_R - \bar{r}) dx d\sigma \\
&\leq \|\partial_x^\alpha (r_0 - \bar{r})\|_2^2 + c \int_0^t (\|\nabla_x \mathbf{u}_R\|_\infty \|r_R - \bar{r}\|_{s,2} + \|\nabla_x (r_R - \bar{r})\|_\infty \|\mathbf{u}_R\|_{s,2}) \|\partial_x^\alpha (r_R - \bar{r})\|_2 d\sigma
\end{aligned} \tag{3.20}$$

provided  $|\alpha| \leq s$ . Here, we took into account

$$\int_{\mathbb{R}^n} \mathbf{u}_R \cdot \nabla_x \partial_x^\alpha r_R \partial_x^\alpha (r_R - \bar{r}) dx = -\frac{1}{2} \int_{\mathbb{R}^n} \operatorname{div}_x \mathbf{u}_R |\partial_x^\alpha (r_R - \bar{r})|^2 dx$$

as well as (3.19). To apply the same treatment to (3.18), we apply Itô’s formula to the function  $\int_{\mathbb{R}^n} |\partial_x^\alpha \mathbf{u}_R|^2 dx$  and gain

$$\begin{aligned}
& \|\partial_x^\alpha \mathbf{u}_R(t)\|_2^2 dx + 2 \int_0^t \int_{\mathbb{R}^n} [\mathbf{u}_R \cdot \nabla_x \partial_x^\alpha \mathbf{u}_R + r_R \nabla_x \partial_x^\alpha (r_R - \bar{r})] \cdot \partial_x^\alpha \mathbf{u}_R dx d\sigma \\
&= \|\partial_x^\alpha \mathbf{u}_0\|_2^2 + 2 \int_0^t \int_{\mathbb{R}^n} [T_3^R + T_4^R] \cdot \partial_x^\alpha \mathbf{u}_R dx d\sigma \\
&+ 2 \int_0^t \int_{\mathbb{R}^n} \partial_x^\alpha \mathbb{F}_R(r_R, \mathbf{u}_R) \cdot \partial_x^\alpha \mathbf{u}_R dW \\
&+ \sum_{k \geq 1} \int_0^t \int_{\mathbb{R}^n} |\partial_x^\alpha \mathbf{F}_{R,k}(r_R, \mathbf{u}_R)|^2 dx d\sigma.
\end{aligned} \tag{3.21}$$

Integrating by parts yields

$$\begin{aligned}
& \int_{\mathbb{R}^n} [\mathbf{u}_R \cdot \nabla_x \partial_x^\alpha \mathbf{u}_R + r_R \nabla_x \partial_x^\alpha (r_R - \bar{r})] \cdot \partial_x^\alpha \mathbf{u}_R dx \\
&= -\frac{1}{2} \int_{\mathbb{R}^n} |\partial_x^\alpha \mathbf{u}_R|^2 \operatorname{div}_x \mathbf{u}_R dx - \int_{\mathbb{R}^n} r_R \operatorname{div}_x \partial_x^\alpha \mathbf{u}_R \partial_x^\alpha (r_R - \bar{r}) dx \\
&- \int_{\mathbb{R}^n} \nabla_x (r_R - \bar{r}) \cdot \partial_x^\alpha \mathbf{u}_R \partial_x^\alpha (r_R - \bar{r}) dx
\end{aligned}$$

Now we combine (3.19)–(3.21) (and multiply (3.21) by  $\gamma - 1$ ) and observe that the term containing  $r_R \partial_x^\alpha (r_R - \bar{r}) \operatorname{div}_x \partial_x^\alpha \mathbf{u}_R$  on the left hand side cancels out. We therefore conclude that

$$\begin{aligned}
& \|(r_R(t) - \bar{r}, \mathbf{u}_R(t))\|_{s,2}^2 \\
& \leq \|(r_0 - \bar{r}, \mathbf{u}_0)\|_{s,2}^2 + c \int_0^t \|\nabla \mathbf{u}_R\|_\infty \left( \|r_R - \bar{r}\|_{s,2}^2 + \|\mathbf{u}_R\|_{s,2}^2 \right) d\sigma \\
& \quad + c \int_0^t \|\nabla(r_R - \bar{r})\|_\infty \left( \|r_R - \bar{r}\|_{s,2}^2 + \|\mathbf{u}_R\|_{s,2}^2 \right) d\sigma \\
& \quad + c \int_0^t \int_{\mathbb{R}^n} \partial_x^\alpha \mathbb{F}_R(r_R, \mathbf{u}_R) \cdot \partial_x^\alpha \mathbf{u}_R dx dW \\
& \quad + c \sum_{k \geq 1} \int_0^t \int_{\mathbb{R}^n} |\partial_x^\alpha \mathbf{F}_{R,k}(r_R, \mathbf{u}_R)|^2 dx d\sigma \\
& = (I)_0 + (I)_1 + \cdots + (I)_4.
\end{aligned} \tag{3.22}$$

We choose  $R \geq R_N$ , where  $R_N$  is chosen in a way that the cut-offs in the definition of  $\mathbb{F}_R$  are not seen for  $t \leq \mathfrak{t}_{R,N}$ , recall (3.16). Now we take the supremum over  $0 \leq t \leq \mathfrak{t}_{R,N} \wedge \delta$ , where  $\delta > 0$ , and apply expectations. Using the definition of  $\mathfrak{t}_{R,N}$  we easily obtain

$$\mathbb{E} \sup_{0 \leq t \leq \mathfrak{t}_{R,N} \wedge \delta} [(I)_1 + (I)_2] \leq c(N) \mathbb{E} \int_0^{\mathfrak{t}_{R,N} \wedge \delta} \|(r_R - \bar{r}, \mathbf{u}_R)\|_{s,2}^2 dt \leq c(N)\delta.$$

In accordance with (3.16), the cut-offs in the definition of  $\mathbb{F}_R$  are not seen for  $t \leq \mathfrak{t}_{R,N}$  if  $R \geq R_N$ . Moreover, the norms of  $\partial_x^\alpha \mathbb{F}_R$  can be controlled by  $N$ . First of all, we have by (2.5)–(2.7)

$$\begin{aligned}
\mathbb{E} \sup_{0 \leq t \leq \mathfrak{t}_{R,N} \wedge \delta} (I)_4 &= \mathbb{E} \sum_{k \geq 1} \int_0^{\mathfrak{t}_{R,N} \wedge \delta} \int_{\mathcal{K}} |\partial_x^\alpha \mathbf{F}_{R,k}(r_R, \mathbf{u}_R)|^2 dx d\sigma \\
&\leq c \mathbb{E} \int_0^{\mathfrak{t}_{R,N} \wedge \delta} \int_{\mathcal{K}} (|\partial_x^\alpha r_R|^2 + |\partial_x^\alpha \mathbf{u}_R|^2) dx d\sigma \\
&\leq c \mathbb{E} \int_0^{\mathfrak{t}_{R,N} \wedge \delta} \int_{\mathcal{K}} (1 + |\partial_x^\alpha (r_R - \bar{r})|^2 + |\partial_x^\alpha \mathbf{u}_R|^2) dx d\sigma \\
&\leq c \mathbb{E} \int_0^{\mathfrak{t}_{R,N} \wedge \delta} (1 + \|(r_R - \bar{r}, \mathbf{u}_R)\|_{s,2}^2) d\sigma \leq c(N)\delta.
\end{aligned}$$

Using the Burkholder-Davis-Gundy inequality, we can estimate the stochastic integral in the same fashion. After applying expectations, we gain using (2.5)–(2.7)

$$\begin{aligned}
& \mathbb{E} \left[ \sup_{0 \leq t \leq \mathfrak{t}_{R,N} \wedge \delta} \left| \int_0^t \int_{\mathbb{R}^n} \partial_x^\alpha \mathbb{F}(r_R, \mathbf{u}_R) \cdot \partial_x^\alpha \mathbf{u}_R dx dW \right| \right] \\
& \lesssim \mathbb{E} \left[ \sum_{k \geq 1} \int_0^{\mathfrak{t}_{R,N} \wedge \delta} \left( \int_{\mathbb{R}^n} \partial_x^\alpha \mathbf{F}_k(r_R, \mathbf{u}_R) \cdot \partial_x^\alpha \mathbf{u}_R dx \right)^2 dt \right]^{\frac{1}{2}}
\end{aligned}$$



$$\begin{aligned}
&\lesssim \mathbb{E} \left[ \int_0^{t_{R,N} \wedge \delta} \left( \sum_{k \geq 1} \|\partial_x^\alpha \mathbf{F}_k(r_R, \mathbf{u}_R)\|_{s,2}^2 \right) \|\mathbf{u}_R\|_{s,2}^2 dt \right]^{\frac{1}{2}} \\
&\lesssim \mathbb{E} \left[ \int_0^{t_{R,N} \wedge \delta} \int_{\mathcal{K}} (|\partial_x^\alpha r_R|^2 + |\partial_x^\alpha \mathbf{u}_R|^2) dx \|\mathbf{u}_R\|_{s,2}^2 dt \right]^{\frac{1}{2}} \\
&\lesssim \mathbb{E} \left[ \int_0^{t_{R,N} \wedge \delta} \int_{\mathcal{K}} (1 + |\partial_x^\alpha (r_R - \bar{r})|^2 + |\partial_x^\alpha \mathbf{u}_R|^2) dx \|\mathbf{u}_R\|_{s,2}^2 dt \right]^{\frac{1}{2}} \\
&\lesssim \mathbb{E} \left[ \int_0^{t_{R,N} \wedge \delta} (1 + \|r_R - \bar{r}\|_{s,2}^2 + \|\mathbf{u}_R\|_{s,2}^2) \|\mathbf{u}_R\|_{s,2}^2 dt \right]^{\frac{1}{2}} \\
&\leq c(N) \sqrt{\delta}.
\end{aligned}$$

Plugging all together, we have shown

$$\mathbb{E} \sup_{0 \leq t \leq t_{R,N} \wedge \delta} \|(r_R(t) - \bar{r}, \mathbf{u}_R(t))\|_{s,2}^2 \leq \mathbb{E} \|(r_0 - \bar{r}, \mathbf{u}_0)\|_{s,2}^2 + c(N)(\sqrt{\delta} + \delta).$$

We obtain

$$\lim_{\delta \rightarrow 0} \sup_R \mathbb{P} \left( \sup_{0 \leq t \leq t_{R,N} \wedge \delta} \|(r_R(t) - \bar{r}, \mathbf{u}_R(t))\|_{s,2}^2 > \|(r_0 - \bar{r}, \mathbf{u}_0)\|_{s,2}^2 + 1 \right) = 0$$

which is (2.10).

### 3.3 Pathwise uniqueness

We mimick the approach of the last subsection and have a look at the difference of two solutions  $(r_R, \mathbf{u}_R)$  and  $(r_L, \mathbf{u}_L)$  which satisfies

$$\begin{aligned}
d\partial_x^\alpha (r_R - r_L) &= -\partial_x^\alpha \left( \mathbf{u}_R \cdot \nabla_x r_R + \frac{\gamma-1}{2} r_R \operatorname{div}_x \mathbf{u}_R \right) dt \\
&\quad + \partial_x^\alpha \left( \mathbf{u}_L \cdot \nabla_x r_L + \frac{\gamma-1}{2} r_L \operatorname{div}_x \mathbf{u}_L \right) dt,
\end{aligned} \tag{3.23}$$

and

$$\begin{aligned}
d\partial_x^\alpha (\mathbf{u}_R - \mathbf{u}_L) &= -\partial_x^\alpha \left( \mathbf{u}_R \cdot \nabla_x \mathbf{u}_R + r_R \nabla_x r_R \right) dt + \partial_x^\alpha \left( \mathbf{u}_L \cdot \nabla_x \mathbf{u}_L + r_L \nabla_x r_L \right) dt \\
&\quad + \left[ \partial_x^\alpha \mathbb{F}(r_R, \mathbf{u}_R) - \partial_x^\alpha \mathbb{F}(r_L, \mathbf{u}_L) \right] dW.
\end{aligned}$$

Multiplying (3.23) by  $\partial_x^\alpha (r_R - r_L)$ , we get

$$\begin{aligned}
\frac{1}{2} d |\partial_x^\alpha (r_R - r_L)|^2 &= -\partial_x^\alpha \left( \mathbf{u}_R \cdot \nabla_x r_R + \frac{\gamma-1}{2} r_R \operatorname{div}_x \mathbf{u}_R \right) \partial_x^\alpha (r_R - r_L) dt \\
&\quad + \partial_x^\alpha \left( \mathbf{u}_L \cdot \nabla_x r_L + \frac{\gamma-1}{2} r_L \operatorname{div}_x \mathbf{u}_L \right) \partial_x^\alpha (r_R - r_L) dt.
\end{aligned} \tag{3.24}$$

Similarly, using Itô's product rule, we obtain

$$\begin{aligned}
& \frac{1}{2} d |\partial_x^\alpha (\mathbf{u}_R - \mathbf{u}_L)|^2 \\
&= -\partial_x^\alpha \left( \mathbf{u}_R \cdot \nabla_x \mathbf{u}_R + r_R \nabla_x r_R \right) \cdot \partial_x^\alpha (\mathbf{u}_R - \mathbf{u}_L) dt \\
&+ \partial_x^\alpha \left( \mathbf{u}_L \cdot \nabla_x \mathbf{u}_L + r_L \nabla_x r_L \right) \cdot \partial_x^\alpha (\mathbf{u}_R - \mathbf{u}_L) dt \\
&+ \left[ \partial_x^\alpha \mathbb{F}_R(r_R, \mathbf{u}_R) - \partial_x^\alpha \mathbb{F}_L(r_L, \mathbf{u}_L) \right] \cdot \partial_x^\alpha (\mathbf{u}_R - \mathbf{u}_L) dW \\
&+ \frac{1}{2} \sum_{k \geq 1} \left( \partial_x^\alpha \mathbf{F}_{R,k}(r_R, \mathbf{u}_R) - \partial_x^\alpha \mathbf{F}_{L,k}(r_L, \mathbf{u}_L) \right)^2 dt.
\end{aligned} \tag{3.25}$$

We sum (3.24) and (3.25), integrate over the physical space, and perform the same estimates as in the previous section. Note that the highest order terms in (3.24) read

$$\begin{aligned}
& \int_{\mathbb{R}^n} (\mathbf{u}_R \cdot \nabla_x \partial_x^\alpha r_R - \mathbf{u}_L \cdot \nabla_x \partial_x^\alpha r_L) \partial_x^\alpha (r_R - r_L) dx \\
&+ \frac{\gamma - 1}{2} \int_{\mathbb{R}^n} (r_R \operatorname{div}_x \partial_x^\alpha \mathbf{u}_R - r_L \operatorname{div}_x \partial_x^\alpha \mathbf{u}_L) \partial_x^\alpha (r_R - r_L) dx \\
&= \int_{\mathbb{R}^n} \left( (\mathbf{u}_R - \mathbf{u}_L) \cdot \nabla_x \partial_x^\alpha r_R \partial_x^\alpha (r_R - r_L) + \frac{1}{2} \operatorname{div}_x \mathbf{u}_L |\partial_x^\alpha (r_R - r_L)|^2 \right) dx \\
&+ \frac{\gamma - 1}{2} \int_{\mathbb{R}^n} (r_R - r_L) \operatorname{div}_x \partial_x^\alpha \mathbf{u}_L \partial_x^\alpha (r_R - r_L) dx \\
&+ \frac{\gamma - 1}{2} \int_{\mathbb{R}^n} r_R \operatorname{div}_x \partial_x^\alpha (\mathbf{u}_R - \mathbf{u}_L) \partial_x^\alpha (r_R - r_L) dx.
\end{aligned}$$

Here, the last integral cancels out after integration by parts, with its counterpart in (3.25). Summing over all  $\alpha$  with  $|\alpha| \leq s - 1$  we deduce

$$\begin{aligned}
& d \left( \|r_R - r_L\|_{W^{s-1,2}}^2 + \|\mathbf{u}_R - \mathbf{u}_L\|_{W^{s-1,2}}^2 \right) \\
&\leq c(R, L) \left[ (1 + \|(r_R, \mathbf{u}_R)\|_{W^{s,2}}^2 + \|(r_L, \mathbf{u}_L)\|_{W^{s,2}}^2) \left( \|r_R - r_L\|_{W^{s-1,2}}^2 + \|\mathbf{u}_R - \mathbf{u}_L\|_{W^{s-1,2}}^2 \right) \right] dt \\
&+ \left[ \partial_x^\alpha \mathbb{F}_R(r_R, \mathbf{u}_R) - \partial_x^\alpha \mathbb{F}_L(r_L, \mathbf{u}_L) \right] \cdot \partial_x^\alpha (\mathbf{u}_R - \mathbf{u}_L) dW \\
&+ \frac{1}{2} \sum_{k \geq 1} \left( \partial_x^\alpha \mathbf{F}_{R,k}(r_R, \mathbf{u}_R) - \partial_x^\alpha \mathbf{F}_{L,k}(r_L, \mathbf{u}_L) \right)^2 dt,
\end{aligned} \tag{3.26}$$

where  $s > \frac{n}{2} + 2$ . As the initial data coincide, we obtain by Gronwall's lemma for  $R, L$  large enough

$$\mathbb{E} \sup_{0 \leq t \leq t_{R,N} \wedge t_{L,N}} \left( \|r_R - r_L\|_{W^{s-1,2}}^2 + \|\mathbf{u}_R - \mathbf{u}_L\|_{W^{s-1,2}}^2 \right) = 0.$$

This certainly yields

$$\mathbb{E} \sup_{0 \leq t \leq t_{R,N} \wedge t_{L,N}} \left( \|r_R - r_L\|_{W^{s,2}}^2 + \|\mathbf{u}_R - \mathbf{u}_L\|_{W^{s,2}}^2 \right) = 0.$$

which implies (2.9).

### 3.4 Conclusion

As shown in the last two subsections, we are in the position to apply Lemma 2.1. We infer the existence of a  $\mathbb{P}$ -a.s. positive stopping time  $\mathfrak{t}$  and a predictable process  $(\mathfrak{t}, r, \mathbf{u})$  such that

$$\mathbb{E} \sup_{0 \leq t \leq \mathfrak{t}} \left( \|r_R - r\|_{W^{s,2}}^2 + \|\mathbf{u}_R - \mathbf{u}\|_{W^{s,2}}^2 \right) = 0 \quad (3.27)$$

as  $R \rightarrow \infty$ . By (3.27), it is easy to pass to the limit in (3.11) and (3.12). By Itô's formula, we conclude that  $(\mathfrak{t}, \varrho, \mathbf{u}) := \left( \mathfrak{t}, \left( \frac{\gamma-1}{2a\gamma} r^2 \right)^{\frac{1}{\gamma-1}}, \mathbf{u} \right)$  is a solution to (1.1)–(1.2) in the sense of Definition 2.2.

Assume that  $[\varrho^1, \mathbf{u}^1, \mathfrak{t}^1]$  and  $[\varrho^2, \mathbf{u}^2, \mathfrak{t}^2]$  are two local strong solutions with the same initial datum. Then we obtain that  $[\varrho^1, \mathbf{u}^1]$  and  $[\varrho^2, \mathbf{u}^2]$  coincide a.s. as a direct consequence of (3.26) (note that the systems (1.1)–(1.2) and (3.1)–(3.2) are equivalent up to the stopping time). This also implies that the blow-up time for two maximal strong solutions (in the sense of Definition 2.3) coincide. So, maximal strong solutions are unique. So far, we have assumed (3.8) which is quite restrictive. This assumption can be removed as in [8, Sec. 4.3]. Finally, it is standard to extend the local strong solution to a maximal strong solution, cf. [12, 20, 26, 34]. For our purposes, the method from [8, Sec. 4.4] can be used. The proof of Theorem 2.4 is complete.

## 4 Proof of Theorem 2.10

Let

$$[(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \geq 0}, \mathbb{P}), \rho_\varepsilon, \mathbf{v}_\varepsilon, W]_{\varepsilon > 0}$$

be a sequence of finite energy weak martingale solutions to (2.18), existence of which is guaranteed by Theorem 2.7. Our aim is to pass to the limit  $\varepsilon \rightarrow 0$ .

### 4.1 Relative energy inequality

The *relative energy inequality* is a tool which enables us to compare  $\rho_\varepsilon, \mathbf{v}_\varepsilon$  with some smooth comparison functions. Let  $(f, \mathbf{U})$  be a pair of stochastic processes which are adapted to the filtration  $(\mathfrak{F}_t)_{t \geq 0}$  and which satisfies

$$\begin{aligned} df &= D_t^d f dt + \mathbb{D}_t^s f dW, \\ d\mathbf{U} &= D_t^d \mathbf{U} dt + \mathbb{D}_t^s \mathbf{U} dW. \end{aligned} \quad (4.1)$$

In the above,  $D_t^d f, D_t^d \mathbf{U}$  are functions of  $(\omega, t, x)$  and  $\mathbb{D}_t^s f, \mathbb{D}_t^s \mathbf{U}$  belong to  $L_2(\mathfrak{U}; L^2(\mathbb{R}^n))$  for a.e  $(\omega, t) \in \Omega \times [0, T]$ . For the *relative energy functional*

$$\mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | f, \mathbf{U}) = \int_{\mathbb{R}^n} \left[ \frac{1}{2} \rho_\varepsilon |\mathbf{v}_\varepsilon - \mathbf{U}|^2 + H(\rho_\varepsilon, f) \right] dx,$$

we have that for any  $t \in (0, T)$ ,

$$\begin{aligned} \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | f, \mathbf{U})(t) &+ \int_0^t \int_{\mathbb{R}^n} \varepsilon [\mathbb{S}(\nabla_x \mathbf{v}_\varepsilon) - \mathbb{S}(\nabla_x \mathbf{U})] : (\nabla_x \mathbf{v}_\varepsilon - \nabla_x \mathbf{U}) \, dx \, ds \\ &\leq \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | f, \mathbf{U})(0) + M_{RE}(t) + \int_0^t \mathcal{R}(\rho_\varepsilon, \mathbf{v}_\varepsilon | f, \mathbf{U})(s) \, ds \end{aligned} \quad (4.2)$$

$\mathbb{P}$ -a.s., where

$$\begin{aligned} \mathcal{R}(\rho_\varepsilon, \mathbf{v}_\varepsilon | f, \mathbf{U}) &= \int_{\mathbb{R}^n} \varepsilon \mathbb{S}(\nabla_x \mathbf{U}) : (\nabla_x \mathbf{U} - \nabla_x \mathbf{v}_\varepsilon) \, dx \\ &+ \int_{\mathbb{R}^n} \rho_\varepsilon (D_t^d \mathbf{U} + \mathbf{v}_\varepsilon \cdot \nabla_x \mathbf{U}) \cdot (\mathbf{U} - \mathbf{v}_\varepsilon) \, dx \\ &+ \int_{\mathbb{R}^n} [(f - \rho_\varepsilon) P''(f) D_t^d f + \nabla_x P'(f) \cdot (f \mathbf{U} - \rho_\varepsilon \mathbf{v}_\varepsilon)] \, dx \\ &+ \int_{\mathbb{R}^n} [p(f) - p(\rho_\varepsilon)] \operatorname{div}_x(\mathbf{U}) \, dx \\ &+ \frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathbb{R}^n} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \mathbb{D}_t^s \mathbf{U}(e_k) \right|^2 \, dx. \end{aligned} \quad (4.3)$$

Here,  $M_{RE}$  is a real valued square integrable martingale and  $P(\varrho) = \frac{a}{\gamma-1} \varrho^\gamma$  is the pressure potential. Let us finally specify the appropriate smoothness assumptions we require for  $D_t^d f$ ,  $D_t^d \mathbf{U}$ ,  $\mathbb{D}_t^s f$  and  $\mathbb{D}_t^s \mathbf{U}$ . We suppose that

$$\begin{aligned} (f - \bar{\rho}) &\in C_c^\infty([0, T] \times \mathbb{R}^n), \quad \mathbf{U} \in C_c^\infty([0, T] \times \mathbb{R}^n), \quad \mathbb{P}\text{-a.s.}, \\ \mathbb{E} \left[ \sup_{t \in [0, T]} \|f - \bar{\rho}\|_{W^{1,q}(\mathbb{R}^n)}^2 \right]^q &+ \mathbb{E} \left[ \sup_{t \in [0, T]} \|\mathbf{U}\|_{W^{1,q}(\mathbb{R}^n)}^2 \right]^q \leq c(q) \quad \forall 2 \leq q < \infty, \end{aligned} \quad (4.4)$$

$$0 < \underline{f} \leq f(t, x) \leq \bar{f} \quad \mathbb{P}\text{-a.s.} \quad (4.5)$$

Moreover,  $f, \mathbf{U}$  satisfy

$$\begin{aligned} D^d f, D^d \mathbf{U} &\in L^q(\Omega; L^q(0, T; W^{1,q}(\mathbb{R}^n))), \quad \mathbb{D}^s f, \mathbb{D}^s \mathbf{U} \in L^2(\Omega; L^2(0, T; L_2(\mathfrak{U}; L^2(\mathbb{R}^n))))), \\ \left( \sum_{k \geq 1} |\mathbb{D}^s f(e_k)|^q \right)^{\frac{1}{q}}, \left( \sum_{k \geq 1} |\mathbb{D}^s \mathbf{U}(e_k)|^q \right)^{\frac{1}{q}} &\in L^q(\Omega; L^q(0, T; L^q(\mathbb{R}^n))). \end{aligned} \quad (4.6)$$

The *relative energy inequality* (4.2) is a consequence of the *energy inequality* (2.13). The proof relies on a sophisticated application of Itô's formula in infinite dimensions. The latter one can be found in [7, Lemma 3.1] in case of periodic boundary conditions. For a corresponding statement in the current setting where the underlying domain is  $\mathbb{R}^n$  we refer to [33, Sec. 3.6]. In order to prove Theorem 2.10, we choose  $(f, \mathbf{U}) = (\varrho(\cdot \wedge \mathbf{t}_R), \mathbf{u}(\cdot \wedge \mathbf{t}_R))$  where  $(\varrho, \mathbf{u}, (\mathbf{t}_R)_{R \in \mathbb{N}}, \mathbf{t})$  is the unique maximal strong pathwise solution to (1.1)–(1.2) which exists by Theorem 2.4

(note that the  $C_c^\infty$ -assumption in (4.4) can be relaxed by a standard approximation argument). Recall that the stopping time  $\mathfrak{t}_R$  announces the blow-up and satisfies

$$\sup_{t \in [0, \mathfrak{t}_R]} \|\mathbf{u}(t)\|_{1, \infty} \geq R \quad \text{on} \quad [t < T];$$

Moreover,  $(f, \mathbf{U}) = (\varrho, \mathbf{u})$  satisfies an equation of the form (4.1), where

$$D_t^d f = -\operatorname{div}_x(\varrho \mathbf{u}), \quad \mathbb{D}_t^s f = 0, \quad D_t^d \mathbf{U} = -\mathbf{u} \cdot \nabla_x \mathbf{u} - \frac{1}{\varrho} \nabla_x p(\varrho), \quad \mathbb{D}_t^s \mathbf{U} = \frac{1}{\varrho} \mathbb{G}(\varrho, \varrho \mathbf{u}).$$

By Theorem 2.4 and (2.5)–(2.7), it is easy to see that (4.5) and (4.6) are satisfied for  $t \leq \mathfrak{t}_R$ . Note in particular the lower bound for  $\varrho$  which follows from the maximum principle (3.14) and (2.19). So, (4.2) holds and the remainder takes the form

$$\begin{aligned} \mathcal{R}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u}) &= \int_{\mathbb{R}^n} \varepsilon \mathbb{S}(\nabla_x \mathbf{u}) : (\nabla_x \mathbf{u} - \nabla_x \mathbf{v}_\varepsilon) \, dx \\ &+ \int_{\mathbb{R}^n} \rho_\varepsilon \left( -\mathbf{u} \cdot \nabla_x \mathbf{u} - \frac{1}{\varrho} \nabla_x p(\varrho) + \mathbf{v}_\varepsilon \cdot \nabla_x \mathbf{u} \right) \cdot (\mathbf{u} - \mathbf{v}_\varepsilon) \, dx \\ &+ \int_{\mathbb{R}^n} \left[ -(\varrho - \rho_\varepsilon) P''(\varrho) \operatorname{div}_x(\varrho \mathbf{u}) + \nabla_x P'(\varrho) \cdot (\varrho \mathbf{u} - \rho_\varepsilon \mathbf{v}_\varepsilon) \right] \, dx \\ &+ \int_{\mathbb{R}^n} [p(\varrho) - p'(\varrho)(\varrho - \rho_\varepsilon) - p(\rho_\varepsilon)] \operatorname{div}_x(\mathbf{u}) \, dx + \int_{\mathbb{R}^n} p'(\varrho)(\varrho - \rho_\varepsilon) \operatorname{div}_x(\mathbf{u}) \, dx \\ &+ \frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathbb{R}^n} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right|^2 \, dx. \end{aligned} \quad (4.7)$$

Note that we can write

$$\varrho - \rho_\varepsilon = (\varrho - \bar{\varrho}) - (\rho_\varepsilon - \bar{\varrho}) \mathbb{I}_{|\rho_\varepsilon - \bar{\varrho}| \leq 1} - (\rho_\varepsilon - \bar{\varrho}) \mathbb{I}_{|\rho_\varepsilon - \bar{\varrho}| > 1},$$

where we have  $\mathbb{P}$ -a.s.

$$\begin{aligned} (\rho_\varepsilon - \bar{\varrho}) \mathbb{I}_{|\rho_\varepsilon - \bar{\varrho}| \leq 1} &\in L^\infty(0, T; L^2(\mathbb{R}^n)), \\ (\rho_\varepsilon - \bar{\varrho}) \mathbb{I}_{|\rho_\varepsilon - \bar{\varrho}| > 1} &\in L^\infty(0, T; L^\gamma(\mathbb{R}^n)), \end{aligned}$$

see Remark (2.8). Consequently, all terms in (4.7) are well-defined due to the regularity of  $(\varrho, \mathbf{u})$ .

## 4.2 Estimating the remainder

In order to estimate the remainder in (4.7) we follow ideas from [7, Sec. 4]. We tacitly assume that  $t \leq \mathfrak{t}_R$  such that  $\mathbf{u}, \nabla \mathbf{u}, \varrho$  and  $\varrho^{-1}$  can be bounded in terms of  $R$ . By using the identities

$$\varrho \nabla_x P'(\varrho) = \nabla_x p(\varrho), \quad \varrho \partial_t P'(\varrho) = \partial_t p(\varrho), \quad -\partial_t \varrho = \operatorname{div}_x(\varrho \mathbf{u}),$$

it holds that

$$\int_{\mathbb{R}^n} \left[ \frac{\rho_\varepsilon}{\varrho} \nabla_x p(\varrho) \cdot (\mathbf{v}_\varepsilon - \mathbf{u}) - (\varrho - \rho_\varepsilon) P''(\varrho) \operatorname{div}_x(\varrho \mathbf{u}) + \nabla_x P'(\varrho) \cdot (\varrho \mathbf{u} - \rho_\varepsilon \mathbf{v}_\varepsilon) \right] \, dx$$

$$\begin{aligned}
&= \int_{\mathbb{R}^n} \left[ \frac{\rho_\varepsilon}{\varrho} \nabla_x p(\varrho) \cdot (\mathbf{v}_\varepsilon - \mathbf{u}) + (\varrho - \rho_\varepsilon) \partial_t [P'(\varrho)] + \nabla_x p(\varrho) \cdot \mathbf{u} - \frac{\rho_\varepsilon}{\varrho} \nabla_x p(\varrho) \cdot \mathbf{v}_\varepsilon \right] dx \\
&= \int_{\mathbb{R}^n} \left[ \partial_t p(\varrho) - \frac{\rho_\varepsilon}{\varrho} \partial_t p(\varrho) + \nabla_x p(\varrho) \cdot \mathbf{u} - \frac{\rho_\varepsilon}{\varrho} \nabla_x p(\varrho) \cdot \mathbf{u} \right] dx \\
&= \int_{\mathbb{R}^n} \left( \frac{\varrho - \rho_\varepsilon}{\varrho} \right) (\partial_t p(\varrho) + \nabla_x p(\varrho) \cdot \mathbf{u}) dx.
\end{aligned}$$

However, since  $(\varrho, \mathbf{u})$  is a strong solution to the continuity equation, it satisfies the *strong* renormalized continuity equation

$$\partial_t p(\varrho) + \nabla_x p(\varrho) \cdot \mathbf{u} = -\gamma p(\varrho) \operatorname{div}_x(\mathbf{u}).$$

By combining this with the identity  $\varrho p'(\varrho) = \gamma p(\varrho)$  yields

$$\int_{\mathbb{R}^n} \left( \frac{\varrho - \rho_\varepsilon}{\varrho} \right) (\partial_t p(\varrho) + \nabla_x p(\varrho) \cdot \mathbf{u}) dx \leq \int_{\mathbb{R}^n} (\rho_\varepsilon - \varrho) p'(\varrho) \operatorname{div}_x(\mathbf{u}) dx$$

since  $1/\gamma < 1$ . So, by collecting the above estimates, we can now deduce from (4.7) that for each  $R \in \mathbb{N}$ ,

$$\begin{aligned}
&\mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u})(t \wedge \mathbf{t}_R) + \varepsilon \int_0^{t \wedge \mathbf{t}_R} \int_{\mathbb{R}^n} \left( \mathbb{S}(\nabla_x \mathbf{v}_\varepsilon) - \mathbb{S}(\nabla_x \mathbf{u}) \right) : (\nabla_x \mathbf{v}_\varepsilon - \nabla_x \mathbf{u}) dx ds \\
&\leq \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u})(0) + M_{RE}(t \wedge \mathbf{t}_R) + \int_0^{t \wedge \mathbf{t}_R} \tilde{\mathcal{R}}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u})(s) ds,
\end{aligned} \tag{4.8}$$

where now

$$\begin{aligned}
\tilde{\mathcal{R}}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u}) &= \varepsilon \int_{\mathbb{R}^n} \mathbb{S}(\nabla_x \mathbf{u}) : (\nabla_x \mathbf{u} - \nabla_x \mathbf{v}_\varepsilon) dx \\
&\quad + \int_{\mathbb{R}^n} \rho_\varepsilon (\mathbf{v}_\varepsilon - \mathbf{u}) \cdot \nabla_x \mathbf{u} \cdot (\mathbf{u} - \mathbf{v}_\varepsilon) dx \\
&\quad - \int_{\mathbb{R}^n} [p(\rho_\varepsilon) - (\rho_\varepsilon - \varrho) p'(\varrho) - p(\varrho)] \operatorname{div}_x(\mathbf{u}) dx \\
&\quad + \frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathbb{R}^n} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right|^2 dx.
\end{aligned} \tag{4.9}$$

Now we observe that

$$\begin{aligned}
\left| \int_{\mathbb{R}^n} \rho_\varepsilon (\mathbf{v}_\varepsilon - \mathbf{u}) \cdot \nabla_x \mathbf{u} \cdot (\mathbf{u} - \mathbf{v}_\varepsilon) dx \right| &\leq \|\nabla_x \mathbf{u}\|_{L_x^\infty} \int_{\mathbb{R}^n} \rho_\varepsilon |\mathbf{v}_\varepsilon - \mathbf{u}|^2 dx \\
&\leq c(R) \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u})
\end{aligned} \tag{4.10}$$

$\mathbb{P}$ -a.s. Since the identity

$$p(\rho_\varepsilon) - (\rho_\varepsilon - \varrho) p'(\varrho) - p(\varrho) = (\gamma - 1) H(\rho_\varepsilon, \varrho)$$

holds, it follows that

$$\begin{aligned}
& \left| \int_{\mathbb{R}^n} [p(\rho_\varepsilon) - (\rho_\varepsilon - \varrho)p'(\varrho) - p(\varrho)] \operatorname{div}_x(\mathbf{u}) \, dx \right| \\
& \leq c \|\operatorname{div}_x \mathbf{u}\|_{L_x^\infty} \int_{\mathbb{R}^n} H(\rho_\varepsilon, \varrho) \, dx \\
& \leq c(R) \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u}).
\end{aligned} \tag{4.11}$$

$\mathbb{P}$ -a.s. Similarly, by Young's inequality for bilinear forms

$$\begin{aligned}
& \left| \int_{\mathbb{R}^n} \varepsilon \mathbb{S}(\nabla_x \mathbf{u}) : (\nabla_x \mathbf{v}_\varepsilon - \nabla_x \mathbf{u}) \, dx \right| \\
& \leq \frac{1}{2} \varepsilon \int_{\mathbb{R}^n} \left( \mathbb{S}(\nabla_x \mathbf{v}_\varepsilon) - \mathbb{S}(\nabla_x \mathbf{u}) \right) : (\nabla_x \mathbf{v}_\varepsilon - \nabla_x \mathbf{u}) \, dx + c \varepsilon \int_{\mathbb{R}^n} |\mathbb{S}(\nabla_x \mathbf{u})|^2 \, dx \\
& \leq \frac{1}{2} \varepsilon \int_{\mathbb{R}^n} \left( \mathbb{S}(\nabla_x \mathbf{v}_\varepsilon) - \mathbb{S}(\nabla_x \mathbf{u}) \right) : (\nabla_x \mathbf{v}_\varepsilon - \nabla_x \mathbf{u}) \, dx + c(R) \varepsilon.
\end{aligned} \tag{4.12}$$

$\mathbb{P}$ -a.s. Lastly, we rewrite

$$\begin{aligned}
& \frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathbb{R}^n} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right|^2 \, dx \\
& = \frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathcal{K}} \chi_{\{\rho_\varepsilon \leq \varrho/2\}} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right|^2 \, dx \\
& + \frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathcal{K}} \chi_{\{\varrho/2 < \rho_\varepsilon < 2\varrho\}} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right|^2 \, dx \\
& + \frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathcal{K}} \chi_{\{\rho_\varepsilon \geq 2\varrho\}} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right|^2 \, dx \\
& =: I_1 + I_2 + I_3
\end{aligned} \tag{4.13}$$

for  $\mathcal{K} \Subset \mathbb{R}^3$  (recall (2.7)). We can now use the inequality  $\rho \leq 1 + \rho^\gamma$  and (2.17) to conclude that

$$\begin{aligned}
I_1 & \leq c \sum_{k \in \mathbb{N}} \int_{\mathcal{K}} \chi_{\{\rho_\varepsilon \leq \varrho/2\}} \left( \frac{1}{\rho_\varepsilon} |\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)|^2 + \frac{\rho_\varepsilon}{\varrho^2} |\mathbf{G}_k(\varrho, \varrho \mathbf{u})|^2 \right) \, dx \\
& \leq c \int_{\mathcal{K}} \chi_{\{\rho_\varepsilon \leq \varrho/2\}} (\rho_\varepsilon + \rho_\varepsilon |\mathbf{v}_\varepsilon|^2 + \rho_\varepsilon |\mathbf{u}|^2) \, dx \\
& \leq c(R) \int_{\mathcal{K}} \chi_{\{\rho_\varepsilon \leq \varrho/2\}} (1 + \rho_\varepsilon^\gamma + \rho_\varepsilon |\mathbf{v}_\varepsilon - \mathbf{u}|^2) \, dx \\
& \leq c(R) \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon \mid \varrho, \mathbf{u}).
\end{aligned} \tag{4.14}$$

Similarly, we obtain by (2.5), (2.6), (2.17), the bounds on  $\varrho$  and the mean-value theorem

$$I_2 \leq \frac{1}{2} \sum_{k \geq 1} \int_{\mathbb{R}^n} \chi_{\frac{\varrho}{2} \leq \rho_\varepsilon \leq 2\varrho} \rho_\varepsilon \left( \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\varrho} \right)^2 \, dx$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{k \geq 1} \int_{\mathbb{R}^n} \chi_{\frac{\varrho}{2} \leq \rho_\varepsilon \leq 2\varrho} \rho_\varepsilon \left( \frac{\mathbf{G}_k(\varrho, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\varrho} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right)^2 dx \\
& \leq c(R) \int_{\mathbb{R}^n} \chi_{\frac{\varrho}{2} \leq \rho_\varepsilon \leq 2\varrho} \left( |\rho_\varepsilon - \varrho|^2 (1 + |\rho_\varepsilon \mathbf{v}_\varepsilon - \varrho \mathbf{u}|^2) \right) dx \\
& \leq c(R) \int_{\mathbb{R}^n} \chi_{\frac{\varrho}{2} \leq \rho_\varepsilon \leq 2\varrho} \left( |\rho_\varepsilon - \varrho|^2 (1 + |\mathbf{u}|^2) + |\rho_\varepsilon (\mathbf{v}_\varepsilon - \mathbf{u})|^2 \right) dx \\
& \leq c(R) \int_{\mathbb{R}^n} \chi_{\frac{\varrho}{2} \leq \rho_\varepsilon \leq 2\varrho} |\rho_\varepsilon - \varrho|^2 dx + \int_{\mathbb{R}^n} \rho_\varepsilon |\mathbf{v}_\varepsilon - \mathbf{u}|^2 dx \\
& \leq c(R) \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u}).
\end{aligned}$$

Estimating  $I_3$  in (4.13) is similar to (4.14). So, we can conclude from (4.13) that

$$\frac{1}{2} \sum_{k \in \mathbb{N}} \int_{\mathbb{R}^n} \rho_\varepsilon \left| \frac{\mathbf{G}_k(\rho_\varepsilon, \rho_\varepsilon \mathbf{v}_\varepsilon)}{\rho_\varepsilon} - \frac{\mathbf{G}_k(\varrho, \varrho \mathbf{u})}{\varrho} \right|^2 dx \leq c(R) \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u}). \quad (4.15)$$

### 4.3 Conclusion

Collecting the estimates (4.10)–(4.15), we have shown that

$$\int_0^{t \wedge t_R} \mathcal{R}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u}) ds \leq c(R) \left( \int_0^{t \wedge t_R} \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u})(s) ds + \varepsilon \right). \quad (4.16)$$

Combining (4.16) and (4.8) and applying Gronwall's lemma yields

$$\begin{aligned}
\sup_{t \in (0, T)} \mathbb{E} \left[ \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u})(t \wedge t_R) \right] + \varepsilon \mathbb{E} \left[ \int_0^{T \wedge t_R} \int_{\mathbb{R}^n} [\mathbb{S}(\nabla_x \mathbf{v}_\varepsilon) - \mathbb{S}(\nabla_x \mathbf{u})] : (\nabla_x \mathbf{v}_\varepsilon - \nabla_x \mathbf{u}) dx ds \right] \\
\leq c(R) \mathbb{E} \left[ \mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u})(0) + \varepsilon \right].
\end{aligned} \quad (4.17)$$

Note that we have

$$\mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u})(0) = \int_{\mathbb{R}^n} \frac{1}{2} \rho_{0, \varepsilon} |\mathbf{v}_{0, \varepsilon} - \mathbf{u}_0|^2 dx + \int_{\mathbb{R}^n} H(\varepsilon \rho_{0, \varepsilon}, \varrho_0) dx$$

which converges to zero in expectation by (2.19)–(2.21). Consequently, we obtain

$$\mathcal{E}(\rho_\varepsilon, \mathbf{v}_\varepsilon | \varrho, \mathbf{u})(0) \rightarrow 0 \quad (4.18)$$

as  $\varepsilon \rightarrow 0$ . The convergence (2.23) then follows from passing to the limit  $\varepsilon \rightarrow 0$  in (4.17).  $\square$

## References

- [1] R. Agemi : The initial-boundary value problem for inviscid barotropic fluid motion. *Hokkaido Math. J.*, 10(1):156–182, 1981.



- [2] T. Alazard : Low Mach number limit of the full Navier-Stokes equations. *Arch. Ration. Mech. Anal.*, 180(1):1–73, 2006.
- [3] H. Beirão da Veiga : On the barotropic motion of compressible perfect fluids. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 8(2):317–351, 1981.
- [4] F. Berthelin, J. Vovelle : Stochastic isentropic Euler equations. *arXiv preprint arXiv:1310.8093*, 2013.
- [5] H. Bessaih : Martingale solutions for stochastic Euler equations. *Stochastic Anal. Appl.*, 17(5):713–725, 1999.
- [6] H. Bessaih, F. Flandoli : 2-D Euler equation perturbed by noise. *NoDEA Nonlinear Differential Equations Appl.*, 6(1):35–54, 1999.
- [7] D. Breit, E. Feireisl, M. Hofmanová : Compressible fluids driven by stochastic forcing: the relative energy inequality and applications. *Comm. Math. Phys.*, 350(2):443–473, 2017.
- [8] D. Breit, E. Feireisl, M. Hofmanová : Local strong solutions to the stochastic compressible Navier-Stokes system. *Commun. PDE* 43, 313–345. (2018).
- [9] D. Breit, E. Feireisl, M. Hofmanová : On solvability and ill-posedness of the compressible Euler system subject to stochastic forces. *arXiv preprint arXiv:1705.08097*, 2017.
- [10] D. Breit, E. Feireisl, M. Hofmanová : *Stochastically forced compressible fluid flows. De Gruyter Series in Applied and Numerical Mathematics.* De Gruyter, Berlin/Munich/Boston, (2018).
- [11] D. Breit, M. Hofmanová : Stochastic Navier-Stokes equations for compressible fluids. *Indiana Univ. Math. J.*, 65(4):1183–1250, 2016.
- [12] Z. Brzeźniak, B. Masłowski, J. Seidler : Stochastic nonlinear beam equations. *Probab. Theory Related Fields*, 132(1):119–149, 2005.
- [13] Z. Brzeźniak, S. Peszat : Stochastic two dimensional Euler equations. *Ann. Probab.*, 29(4):1796–1832, 2001.
- [14] M. Capiński, N. J. Cutland : Stochastic Euler equations on the torus. *Ann. Appl. Probab.*, 9(3):688–705, 1999.
- [15] G. Da Prato, J. Zabczyk : *Stochastic equations in infinite dimensions*, volume 44 of *Encyclopedia of Mathematics and its Applications.* Cambridge University Press, Cambridge, 1992.
- [16] C. M. Dafermos : The second law of thermodynamics and stability. *Arch. Rational Mech. Anal.*, 70(2):167–179, 1979.
- [17] C. De Lellis and L. Székelyhidi, Jr. On admissibility criteria for weak solutions of the Euler equations. *Arch. Ration. Mech. Anal.*, **195**(1):225–260, 2010.

- [18] C. De Lellis, L. Székelyhidi, Jr : The  $h$ -principle and the equations of fluid dynamics. *Bull. Amer. Math. Soc. (N.S.)*, 49(3):347–375, 2012.
- [19] R. J. DiPerna, P.-L. Lions: Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.* 98, 511–547, 1989.
- [20] K. D. Elworthy : *Stochastic differential equations on manifolds*, volume 70 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge-New York, 1982.
- [21] E. Feireisl : Weak solutions to problems involving inviscid fluids. In *Mathematical fluid dynamics, present and future*, volume 183 of *Springer Proc. Math. Stat.*, pages 377–399. Springer, Tokyo, 2016.
- [22] E. Feireisl, A. Novotný, H. Petzeltová: On the existence of globally defined weak solutions to the Navier–Stokes equations of compressible isentropic fluids. *J. Math. Fluid. Mech.* 3, 358–392, 2001.
- [23] E. Feireisl, B. J. Jin, A. Novotný : Relative entropies, suitable weak solutions, and weak-strong uniqueness for the compressible Navier-Stokes system. *J. Math. Fluid Mech.*, 14(4):717–730, 2012.
- [24] N. Glatt-Holtz, M. Ziane : Strong pathwise solutions of the stochastic Navier-Stokes system. *Adv. Differential Equations*, 14(5-6):567–600, 2009.
- [25] N. E. Glatt-Holtz, V. C. Vicol : Local and global existence of smooth solutions for the stochastic Euler equations with multiplicative noise. *Ann. Probab.*, 42(1):80–145, 2014.
- [26] J. Jacod : *Calcul stochastique et problèmes de martingales*, volume 714 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979.
- [27] J. U. Kim : On the stochastic Euler equations in a two-dimensional domain. *SIAM J. Math. Anal.*, 33(5):1211–1227, 2002.
- [28] J. U. Kim : Existence of a local smooth solution in probability to the stochastic Euler equations in  $\mathbf{R}^3$ . *J. Funct. Anal.*, 256(11):3660–3687, 2009.
- [29] J. U. Kim : On the stochastic quasi-linear symmetric hyperbolic system. *J. Differential Equations*, 250(3):1650–1684, 2011.
- [30] P.-L. Lions : *Mathematical topics in fluid mechanics. Vol. 1*, volume 3 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press, Oxford University Press, New York, 1996.
- [31] A. Majda : *Compressible fluid flow and systems of conservation laws in several space variables*, volume 53 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1984.
- [32] P. R. Mensah : Existence of martingale solutions and the incompressible limit for stochastic compressible flows on the whole space. *Ann. Mat. Pura Appl. (4)*, 196(6):2105–2133, 2017.

- [33] P. R. Mensah : The stochastic compressible Navier-Stokes system on the whole space and some singular limits. *PhD thesis*, Heriot-Watt University, 2018.
- [34] R. Mikulevicius, B. L. Rozovskii : Stochastic Navier-Stokes equations for turbulent flows. *SIAM J. Math. Anal.*, 35(5):1250–1310, 2004.
- [35] R. Mikulevicius, G. Valiukevicius : On stochastic Euler equation in  $\mathbf{R}^d$ . *Electron. J. Probab.*, 5:no. 6, 20, 2000.
- [36] F. Sueur : On the inviscid limit for the compressible Navier-Stokes system in an impermeable bounded domain. *J. Math. Fluid Mech.*, 16(1):163–178, 2014.