



Heriot-Watt University  
Research Gateway

# Modulation instability of discrete angular momentum in fiber rings

## Citation for published version:

Maitland, C, Faccio, D & Biancalana, F 2019, Modulation instability of discrete angular momentum in fiber rings. in *CLEO: Applications and Technology 2019.*, JW2A.39, OSA Publishing, CLEO: Applications and Technology 2019, San Jose, United States, 5/05/19. [https://doi.org/10.1364/CLEO\\_AT.2019.JW2A.39](https://doi.org/10.1364/CLEO_AT.2019.JW2A.39)

## Digital Object Identifier (DOI):

[10.1364/CLEO\\_AT.2019.JW2A.39](https://doi.org/10.1364/CLEO_AT.2019.JW2A.39)

## Link:

[Link to publication record in Heriot-Watt Research Portal](#)

## Document Version:

Peer reviewed version

## Published In:

CLEO

## Publisher Rights Statement:

© 2019 Optical Society of America. One print or electronic copy may be made for personal use only. Systematic reproduction and distribution, duplication of any material in this paper for a fee or for commercial purposes, or modifications of the content of this paper are prohibited.

## General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

## Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [open.access@hw.ac.uk](mailto:open.access@hw.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Modulation Instability of Discrete Angular Momentum in Fiber Rings

Calum Maitland<sup>1\*</sup> Daniele Faccio<sup>2</sup> and Fabio Biancalana<sup>1</sup>

<sup>1</sup> School of Engineering and Physical Sciences, Heriot-Watt University, Edinburgh EH14 4AS, UK

<sup>2</sup> School of Physics and Astronomy, Kelvin Building, University of Glasgow, Glasgow G12 8QQ, UK

\*cm350@hw.ac.uk

**Abstract:** We present an analysis of modulation instability in a ring of coupled optical fibers. Plane waves are shown to be unstable to perturbations carrying discrete angular momenta, both for normal and anomalous group velocity dispersion. © 2019 The Author(s)

**OCIS codes:** 190.3100, 190.4370.

Modulation instability (MI) of plane waves in anomalously-dispersive fibers with a self-focusing Kerr nonlinearity is one of the most well-known phenomena in nonlinear optics [1]. It has been observed in higher dimensional systems, in particular as an azimuthal instability of optical vortices in continuous self-focusing Kerr media [2]. Circular arrays of coupled optical fibres have been shown to support supermodes carrying angular momentum as discrete optical vortices [3]. The nonlinear optical properties of fiber rings have been examined previously, particularly in the context of  $\mathcal{PT}$ -symmetry breaking [4] and the stability of modes in fibers with only a few cores [5]. Here we show that plane wave supermodes of fiber arrays as shown in figure 1 are unstable in the presence of perturbations and that the gain spectra of these perturbations depends on their angular momenta.




Fig. 1. Six core fibre ring. Light couples between cores through the overlap of their evanescent fields.

We model light propagation in the N-core fiber array with N coupled (1+1) dimensional nonlinear Schrödinger equations; the equation for the field  $E_n$  in the  $n^{\text{th}}$  fiber is

$$i\partial_z E_n = \frac{\beta_2}{2} \partial_t^2 E_n - \gamma |E_n|^2 E_n - \Delta (E_{n+1} + E_{n-1}) \quad (1)$$

Following the standard MI derivation in Agrawal [1], we consider a plane-wave solution of (1)

$$\bar{E}_n = \sqrt{P_0} \exp(i2\pi mn/N + ik_0 z) \quad (2)$$

as a pump, perturbed by a pair of weak excitations coupled through the nonlinearity

$$a_n(z, t) = a_S \exp(i(Kz + 2\pi ln/N - \Omega t)) + a_I \exp(-i(Kz + 2\pi(l - 2m)n/N - \Omega t)) \quad (3)$$

These perturbations will grow exponentially if their wavenumber has an imaginary component, which requires the perturbation's detuning  $\Omega$  from the pump to lie between two critical frequencies,  $\Omega_{C1}$  and  $\Omega_{C2}$ . If this holds, then the perturbations grow exponentially in power at a rate

$$G(\Omega, \Delta) = \text{Re} \left( \sqrt{4\gamma^2 P_0^2 - (\beta_2 \Omega^2 + 2\gamma P_0 + 2\Delta_{l,m} + 2\Delta_{2m-l,m})^2} \right) \quad (4)$$




Fig. 2. Gain spectra for perturbations carrying angular momentum  $|l| = 1$  as a function of the detuning  $\Omega$  and the fiber coupling strength  $\Delta$ . **(a)** Anomalous GVD  $\beta_2 = -1$ ; **(b)** Normal GVD  $\beta_2 = 1$

where  $\Delta_{x,m} = \Delta (\cos(2\pi x/N) - \cos(2\pi m/N))$ . Example plots of gain spectra in a six-core fibre ring are shown in figure 2, given pump and probe angular momentum  $m = 0$  and  $|l| = 1$  respectively. In contrast to standard MI, here we see MI can occur for both normal and anomalous GVD. If  $\beta_2 < 0$ , perturbations with non-zero  $l$  see suppressed instability above a threshold coupling strength. With  $\beta_2 > 0$ , gain initially grows stronger with increasing coupling, until it forks into two sidebands. Higher order angular momenta experience similar instabilities; figure 3 compares the gain sidebands for  $|l| = 1, 2, 3$  **(a)** predicted by eq. (4) in a six-core ring with the spectra in a single core **(b)** obtained by numerically integrating eq. (1). We will also explore how twisting the array can tune the MI gain.




Fig. 3. **(a)** Predicted MI gain for perturbations carrying different angular momenta  $|l| = 1, 2, 3$  in an homogeneously pumped six-core fiber ring with  $\beta_2 = 1\text{ps}^2/\text{km}$ ,  $\Delta = 1/\text{cm}$ ,  $\gamma P_0 = 0.1/\text{cm}$  **(b)** Spectrum  $|E_1(\Omega, z)|^2$  in a single core of the ring from a numerical simulation of (1). Blue, magenta and red dashed lines indicate the MI critical frequencies for  $|l| = 1, 2, 3$ , verifying the accuracy of eq. (4).

## References

1. G. P. Agrawal, *Nonlinear Fiber Optics (3rd ed.)* (Academic, 2001).
2. L. T. Vuong *et al.*, “Collapse of Optical Vortices”, *Phys. Rev. Lett.* **96**, 133901 (2006).
3. C. N. Alexeyev, A. V. Volyar and M. A. Yavorsky, “Linear azimuthons in circular fiber arrays and optical angular momentum of discrete optical vortices”, *Phys. Rev. A* **80**, 063821 (2009).
4. C. Castro-Castro, Y. Shen, G. Srinivasan, A. B. Aceves and P. G. Kevrekidis, “Light dynamics in nonlinear trimers and twisted multicore fibers”, *J. Nonlinear Opt. Phys. Mater.* **25**, 1650042 (2016).
5. J. H. Li, H. Zhou, K. S. Chiang and S. R. Xiao, “Modulation instabilities in equilateral three-core optical fibers”, *J. Opt. Soc. Am. B* **33** (11), 2357 (2016).