



Heriot-Watt University  
Research Gateway

## Rhamnolipid, a naturally produced oil dispersant, may improve oil spill remediation

### Citation for published version:

Nikolova, C, Ijaz, U, Magill, C, Kleindienst, S, Joye, SB & Gutierrez, T, *Rhamnolipid, a naturally produced oil dispersant, may improve oil spill remediation*, 2021, Digital or Visual Products.  
<https://doi.org/10.21203/rs.3.rs-967247/v1>

### Digital Object Identifier (DOI):

[10.21203/rs.3.rs-967247/v1](https://doi.org/10.21203/rs.3.rs-967247/v1)

### Link:

[Link to publication record in Heriot-Watt Research Portal](#)

### Document Version:

Early version, also known as pre-print

### General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [open.access@hw.ac.uk](mailto:open.access@hw.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Rhamnolipid, a naturally produced oil dispersant, may improve oil spill remediation

Christina Nikolova

Umer Ijaz

Clayton Magill

Sara Kleindienst

Samantha B. Joye

Tony Gutierrez

---

## Video Byte

**Keywords:** dispersant, synthetic dispersant, biosurfactant, rhamnolipid, marine environment, marine, crude oil, hydrocarbons, biodegradation, oil spill, bacteria, microorganisms, microbes, Finasol, remediation, aromatic hydrocarbons, oil degradation, Faroe-Shetland Channel, Atlantic Ocean, Microbiome

**Posted Date:** October 13th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-967247/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

Oil spills have devastating effects on the environment, and thousands, of varying size, occur each year. Spilled oil can be removed from the environment in numerous ways, such as with the use of dispersants to break up oil slicks on the water surface. But while oil spills themselves pose well-known threats to marine life, the methods used for oil cleanup can also have unintended consequences. To examine these effects, researchers recently investigated how treatment of oil with dispersants produced synthetically (Finasol) and by bacteria (rhamnolipid) impact microbial communities and their ability to break down oil from the subarctic Atlantic Ocean. They found that cold-loving bacteria initially dominated the bacterial communities when both dispersants were used, but some key species of bacteria that specialize in breaking down aromatic hydrocarbons, which are the major and most toxic components of crude oil, became abundant over time in only the presence of rhamnolipid. These bacteria were not observed at all when Finasol was used, and this synthetic dispersant had an overall negative effect on microbial biodiversity. While these results were obtained under laboratory conditions, they indicate that rhamnolipid, the more biodegradable and less toxic of the investigated dispersants, may offer an advantage over Finasol, providing valuable guidance for the future use of these dispersants in oil spill remediation.