



Heriot-Watt University
Research Gateway

Effects of herbal growth promoters on common carp (*Cyprinus carpio*)

Citation for published version:

Effendi, I, Yoswaty, D, Syawal, H, Austin, B, Lyndon, AR, Kurniawan, R & Wahyuni, S 2022, 'Effects of herbal growth promoters on common carp (*Cyprinus carpio*)', *Aquaculture, Aquarium, Conservation and Legislation - International Journal of the Bioflux Society*, vol. 15, no. 3, pp. 1096-1103.

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:


Aquaculture, Aquarium, Conservation and Legislation - International Journal of the Bioflux Society

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Effects of herbal growth promoters on common carp (*Cyprinus carpio*)

¹Irwan Effendi, ¹Dessy Yoswaty, ¹Henni Syawal, ²Brian Austin, ³Alastair R. Lyndon, ¹Ronal Kurniawan, ¹Sri Wahyuni

¹ Faculty of Fisheries and Marine Sciences, University of Riau, Indonesia; ² Stirling University, Stirling, United Kingdom; ³ Heriot-Watt University, Edinburgh, United Kingdom. Corresponding author: D. Yoswaty, dyoswaty13@gmail.com

Abstract. The common carp (*Cyprinus carpio*) is the most widely farmed freshwater fish in the world. Various aspects of cultivation continue to be developed to create an improved aquaculture business. Increasing fish growth and improving feed conversion efficiency are part of the popular topics, and the addition of herbal growth promoters to the diet is one of the promising approaches. This study aimed to determine the effect of adding mango leaves (*Mangifera indica*), guava leaves (*Psidium guajava*), and noni leaves (*Morinda citrifolia*) as growth promoters for carp. The method used was an experimental method using a 1-factor completely randomized design (addition of a growth promoter to commercial feed) with 4 treatment levels and 3 replications. The treatment levels were: T0 - commercial pellets without growth promoter (experimental control); T1 - mango leaves at a dose of 1.5 g 100 g⁻¹ commercial feed; T2 - guava leaves at a dose of 1.5 g 100 g⁻¹ commercial feed; T3 - noni leaves dosage of 1.5 g 100 g⁻¹ commercial feed. Fish rearing was carried out in 12 units of floating net cages measuring 2x2x1 m. The density of fish per cage was 25 fish m⁻³. The standard and formulated feeds were administered as much as 5% of body weight every day. Fish were maintained for 2 months, measuring length and weight every 10 days. The results showed that the herbal growth promoter of guava leaf (T2) showed the best effect, with an absolute weight growth of 69.67 g, specific growth rate of 2.90%, feed efficiency of 83.54% and feed conversion ratio of 1.14. It was followed by treatments T3, T1 and T0. This study recommends the addition of guava leaves into common carp feeds at a dose of 1.5 g 100 g⁻¹ to increase growth rate and improve feed conversion ratio for carp culture.

Key Words: feed efficiency, herbal supplement, guava, mango, noni.

Introduction. Carp (*Cyprinus carpio*) is known as a freshwater fish, although recently it was discovered that it is also able to live in brackish water up to 15% salinity (Salati et al 2011; Singh et al 2018). This fish is widely cultivated in Asian countries and some European countries. China is the largest producer, with 70% of the world carp production (FAO 2019). In Indonesia, common carp started to be cultivated around the 1920s. Intensive monoculture production based on artificial feed can be carried out in cages, irrigation reservoirs, and ponds and running water tanks, or in recirculation systems. Its adaptability to the environment is relatively good, maintenance is easy, it can eat commercial feed, the maintenance cycle is relatively short, the taste of meat is preferred and the price is affordable (Khan et al 2016). Common carp is considered as a very important cultivated fish species in Indonesia. Raising common carp on a large scale is a lucrative business opportunity. Market demand for this fish continues to increase yearly. They can be produced in large quantities and in a short period of time (FAO 2019).

Medicinal plants have been reported to increase the growth of cultured fish. They increased digestive enzymes, thereby increasing the survival and growth of aquatic animals. *Spirulina platensis* in fish diet improved the growth of stellate sturgeon *Acipenser stellatus* (Salehi-Farsani et al 2014). Traditional Chinese medicine has a beneficial effect on the growth of carp (Jian & Wu 2004). Apines-Amar et al (2013) reported the benefits of dietary onion and ginger in promoting growth in brown marbled grouper, *Epinephelus fuscoguttatus*. Three herbs (*Alteranthera sessilis*, *Eclipta alba* and *Cissus quadrangularis*) have astringent effects and increase the activity of digestive enzymes (protease, amylase

and lipase) in freshwater shrimp. This leads to increased feed utilization and ultimately to better growth rates, indicated by evidence of increased concentrations of vitamins, proteins, essential amino acids, unsaturated fatty acids and minerals. The content of vitamins C and E in the hepatopancreas as well as the content of sodium and potassium, and muscle in freshwater shrimp increased when the shrimp were fed herbal supplements (Radhakrishnan et al 2014).

Green tea, caraway seed and American ginseng increase the growth rate and feed utilization of tilapia, *Oreochromis niloticus* (Abdel-Tawwab et al 2010; Ahmad & Abdel-Tawwab 2011; Abdel-Tawwab 2012). Papain enzyme-enriched diet improved feed efficiency and growth of fingerlings of Sangkuriang catfish, *Clarias gariepinus* (Rachmawati et al 2019). Fennel seed flour improves the growth performance and feed utilization in tilapia (Ahmad & Abdel-Tawwab 2011). Caffeine increases the growth of *Sparus aurata* at concentrations greater than 0.1% in feed (Chatzifotis et al 2008). Mixing the herbs *Massa Medicata Fermentata*, *Crataegi fructus*, *Artemisia capillary* and *Cnidium officinale* increased the growth and utilization of fatty acids in Japanese flounder, *Paralichthys olivaceus* (Ji et al 2007). *Quillaja saponins* has the ability to increase the growth rate and decrease the metabolic rate of tilapia (Francis et al 2005).

Mango (*Mangifera indica*), guava (*Psidium guajava*) and noni (*Morinda citrifolia*) are tropical plants that are widely grown everywhere, both cared for and free-growing in nature. In addition to the fruit, the leaves, bark, roots and other parts have also been used as medicinal supplements, including for fisheries needs. This study aimed to determine the effect of mango, noni, and guava leaves as growth promoters, to increase the growth performance of common carp.

Material and Method

Time and place of the study. This research was carried out from August to October 2021. Fish rearing was carried out at the floating net cage cultivation center in the Koto Panjang Hydroelectric Power Plant Reservoir, Kampar, Indonesia.

Materials. The fish tested were carp (*Cyprinus carpio*) with an average length of 10 cm and a weight of 15 g, purchased from local farmer. Before starting the experiment, the fish were acclimatized for 7 days. Commercial fish pellets Hi-Pro-Vite 781-2 produced by PT Central Panganpertiwi Tbk., Karawang, West Java, Indonesia, were used in this study. The growth-stimulating herbal ingredients were mango, guava and noni leaves, which grew around the Riau University Campus, Pekanbaru, Indonesia. Proximate analysis was carried out based on Ahmed et al (2013) and growth promoter phytochemical analysis was carried out according to Syawal et al (2021).

Preparation of fish feed. Growth promoter materials (guava, noni, and mango leaves) were collected and dried in the room at room temperature (± 7 days). After drying, the leaves were mashed using a blender and sifted to get a fine leaf powder. Furthermore, 1.5 g of each leaf flour was added and mixed in 100 g of commercial pellets, and stirred homogeneously. The feed mix was milled and pelleted according to the size of the fish mouth opening and dried in the sun for 3-4 days at a temperature of 30-35°C.

Research methodology. This experiment used a 1-factor completely randomized design (addition of a growth promoter to commercial feed) with 4 treatment levels and 3 replications. The treatments were: T0 - commercial pellets without growth promoter (experimental control); T1 - mango leaves at a dose of 1.5 g 100 g⁻¹ commercial feed; T2 - guava leaves at a dose of 1.5 g 100 g⁻¹ commercial feed; T3 - noni leaves dosage of 1.5 g 100 g⁻¹ commercial feed. Fish rearing was carried out in 12 units of floating net cages measuring 2x2x1 m. The density of fish per cage was 25 fish m⁻³. The formulated feeds were fed to fish as much as 5% of body weight daily. Fish were kept for 2 months, with weight measurements every 10 days.

Absolut growth rate. The absolute growth rate was calculated using the following formula:

$$AGR = W_t - W_o$$

Where: AGR - absolute growth rate (g); W_t - average weight of fish at the end of the study (g); W_o - average weight of the fish at the beginning of the study (g).

Specific growth rate. The specific growth rate is calculated using the formula:

$$SGR = \frac{\ln W_t - \ln W_o}{t} \times 100$$

Where: SGR - daily growth rate (% day⁻¹); W_t - weight of fish at the end of the study (g); W_o - weight of fish at the beginning of the study (g); T - length of study (days).

Feed conversion rate. Feed conversion was calculated using the following formula:

$$FCR = \frac{\Sigma F}{(B_t + B_m) - B_o}$$

Where: FCR - feed conversion rate; ΣF - amount of feed given during rearing (g); B_t - fish biomass at the end of rearing (g); B_o - fish biomass at the start of rearing (g); B_m - dead fish biomass during rearing (g).

Feed efficiency. Feed efficiency was calculated using the following formula:

$$FE = \frac{((B_t + B_d) - B_o)}{F} \times 100$$

Where: FE - feed efficiency; ΣF - amount of feed given during rearing (g); B_t - fish biomass at the end of rearing (g); B_d - biomass of fish that died during the study (g); B_o - fish biomass at the start of rearing (g).

Data analysis. Data obtained from physiological measurements were collected and tabulated. They were analyzed statistically using the SPSS version 23 software. The data were analyzed by using one way ANOVA at a 95% confidence level. The Student Newman Keuls test was used when differences were observed.

Water quality. The water quality parameters measured were temperature, pH, dissolved oxygen and ammonia by using a Horiba U-52G-2 Multi-Parameter Water Quality Meter and DLAB Spectrophotometer SP-V1100. Measurements were collected in the afternoon every 10 days during the study.

Results and Discussion

Proximate analysis. The results of the proximate analysis of commercial feed and growth promoter ingredients were normal (Table 1).

Table 1
Proximate analysis of the feed containing the herbal growth promoters

<i>Treatment</i>	<i>Protein (%)</i>	<i>Fat (%)</i>	<i>Crude fiber (%)</i>	<i>Water (%)</i>
Commercial pellets (T0)	31.00	6.00	5.00	10
Mango leaves (T1)	30.97	9.53	6.07	9.25
Guava leaves (T2)	31.27	8.79	7.73	9.12
Noni leaves (T3)	31.44	10.66	5.70	9.84

Water quality. Water quality during the study was in a range adequate to the growth of carp in the Koto Panjang Hydroelectric Power Station reservoir. Water temperature ranged from 28 to 31°C, pH was between 6.4-6.9, and ammonia between 0.0015-0.0018 mg L⁻¹. The optimal temperature for carp growth in the region is 28-32°C and pH between 6.4-7 (Budijono et al 2021). From this data, it can be concluded that the quality of the water used to keep the fish was in normal conditions.

Growth performance. Feeds containing growth promoters had a significant effect ($p < 0.05\%$) on absolute growth rate, specific growth rate, feed efficiency and feed conversion rate. The absolute growth rate ranged from 37.8 to 69.67 g. T2 provided the best AGR (69.67 g), while the lowest was observed in T0 (37.8 g). The best specific growth rate was also recorded in T2 (2.9%), followed by T3 (2.70%), T1 (2.45%) and finally T0 (2.12%). Feed efficiency ranged from 62.98 to 83.54%, with the highest value noted in T2 (83.54%), followed by T3 (82.01%), T1 (71.13%) and finally T0 (62.98%). A similar pattern was also recorded in the variable feed conversion ratio. The lowest feed conversion ratio was recorded in T2 (1.14), followed by T3 (1.15), T1 (1.31) and finally T0 (1.43). More detailed data are presented in Table 2.

Table 2
Absolute growth rate, specific growth rate, feed efficiency and feed conversion rate of carp (*C. carpio*)

Experimental treatments	Absolut growth rate (g)	Specific growth rate (% day ⁻¹)	Feed efficiency (%)	Feed conversion
T0	37.8±0.9 ^a	2.12±0.03 ^a	62.98±1.84 ^a	1.43±0.02 ^c
T1	49.37±1.56 ^b	2.45±0.04 ^b	71.13±1.69 ^b	1.31±0.03 ^b
T2	69.67±1.76 ^d	2.9±0.03 ^d	83.54±1.62 ^c	1.14±0.02 ^a
T3	59.9±3.34 ^c	2.7±0.08 ^c	82.01±2.17 ^c	1.15±0.01 ^a

Note: superscripts on the same row show significant differences ($p < 0.05$).

The results of statistical analysis showed that the addition of herbal growth promoters to the feed was able to increase the growth of carp significantly ($p < 0.05$). Guava leaves (T2) gave the best results in the growth of common carp, with an absolute weight of 69.67 g, specific growth rate of 2.9% day⁻¹, feed efficiency of 83%, and a FCR of 1.14. The lowest effect was found in the treatment without the addition of herbal growth promoter (T0), namely absolute weight, specific growth rate, feed efficiency, and FCR of 37.8 g, 2.12% day⁻¹, 62.98%, and 1.43, respectively (Figure 1).




Figure 1. Carp (*C. carpio*) weight growth during the study.

The addition of mango leaves, guava and noni leaves as growth promoters caused a positive effect on the growth of carp. Some researchers have reported that a number of plants increased fish growth. For example, mango leaves increased the growth of carp and tilapia (Achioye-Nzeh & Obaroh 2010; Obasa et al 2013; Fawole et al 2018; Amulejoye et al 2020), noni leaves improved tilapia development (Kristiana et al 2020), mango leaves improved growth in rainbow trout (Awad & Austin 2010). Noni also helped the growth of African catfish (Azizah et al 2020), and guava leaves the development of tilapia (Abdel-Tawwab & Hamed 2020; Ceballos-Francisco et al 2020; Omitoyin et al 2020), striped catfish (*Pangasianodon hypophthalmus*) (Nhu et al 2020) and common carp (Hoseinifar et al 2019).

The addition of guava in feed increased feed efficiency and decreased FCR for the 60 days of rearing. The content of secondary metabolite compounds from guava such as saponins, phenols, and tannins can increase the growth and appetite of fish, as well as the absorption of nutrients. According to Das et al (2012), saponins function as permeabilizing membranes and can affect growth and increase animal feeding responses. According to Omitoyin et al (2019), phenols, tannins, and saponins from guava stimulate the secretion of digestive enzymes, thereby affecting the growth and absorption of nutrients by fish. The impact obtained in this study is better than that reported by Omitoyin et al (2019). It was reported that tilapia reared for 84 days, fed a diet containing guava leaf herbal growth promoter (a dose of 0.75%), resulted in a growth of 4.76 g of tilapia.

The guava leaves were also able to increase the specific growth rate of carp by 2.9%. This result was better than the research of Omitoyin et al (2019), where the specific growth rate of tilapia fed a diet containing herbal medicine at a dose of 0.50% resulted in a specific growth rate of 1.56%. This is presumably due to several things, including the proximate levels of guava leaf protein.

Guava leaves also contain essential oils (Olusola & Olorunfemi 2017; Sobral-Souza et al 2019), which improve feed quality, productivity, and health. Essential oils mixed with feed can stimulate the central nervous system, which ultimately results in an increase in appetite and consumption of nutrients. The presence of essential oils can stimulate the production of digestive juices that produce a suitable pH for digestive enzymes, such as peptinase (Weli et al 2018). At the same time, there is an increase in the activity of digestive enzymes, the real effect of this mechanism being the improvement of energy conversion and digestion of food substances and a positive influence on the metabolism of nitrogen, amino acids and glucose (Lin et al 2006; Yu et al 2009).

The higher feed efficiency or the lower FCR indicate the more efficient use of feed. The values of feed efficiency and FCR are closely related to the growth rate, because a greater weight gain of fish comes with an increase of the feed conversion (Fry et al 2018; Abdel-Rahim et al 2019).

Conclusions. The results showed that the addition of the herbal growth promoters from mango, guava and noni leaves to the commercial fish feed had a positive effect on the growth of common carp. Guava leaves showed better results in terms of absolute growth rate, specific growth rate, feed efficiency and feed conversion ratio. Therefore, it can be recommended to add guava leaves to feeds up to 1.5 g 100 g⁻¹ feed.

Acknowledgements. The authors would like to thank the Institute for Research and Community Service, University of Riau for funding and supporting this research program through the International Research Collaboration Program in 2021. Thanks are also conveyed to the Center for Coastal and Aquatic Studies, LPPM University of Riau.

Conflict of Interest. The authors declare that there is no conflict of interest.

References

Abdel-Rahim M. M., Awad Y. M., Abdallah Y. A., Radwan S. M., 2019 Effects of four medicinal plants on the bioeconomic analysis and water-use efficiency of Nile tilapia,

- Oreochromis niloticus* fry under a small-scale aquaponics system. *AACL Bioflux* 12(3):851-866.
- Abdel-Tawwab M., 2012 The use of American ginseng (*Panax quinquefolium*) in practical diets for Nile tilapia (*Oreochromis niloticus*): Growth performance and challenge with *Aeromonas hydrophila*. *Journal of Applied Aquaculture* 24(4):366-376.
- Abdel-Tawwab M., Ahmad M. H., Sakr S. M. F., Seden M. E. A., 2010 Use of green tea, *Camellia sinensis* L., in practical diet for growth and protection of Nile tilapia, *Oreochromis niloticus* (L.), against *Aeromonas hydrophila* infection. *Journal of the World Aquaculture Society* 41(S2):203-213.
- Abdel-Tawwab M., Hamed H. S., 2020 Antagonistic effects of dietary guava (*Psidium guajava*) leaves extract on growth, hemato-biochemical, and immunity response of cypermethrin-intoxicated Nile tilapia, *Oreochromis niloticus*, fingerlings. *Aquaculture* 529:735668.
- Achioye-Nzeh G. C., Obaroh I. O., 2010 Effect of *Mangifera indica* leaves extract on growth response of *Oreochromis niloticus*. *Animal Production Research Advances* 6(2).
- Ahmad M. H., Abdel-Tawwab M., 2011 The use of caraway seed meal as a feed additive in fish diets: Growth performance, feed utilization, and whole-body composition of Nile tilapia, *Oreochromis niloticus* (L.) fingerlings. *Aquaculture* 314(1-4):110-114.
- Ahmed K., Shaheen M., Mirzaei F., Khan Z., Gondal S., Fardous A., Hussain A., Arshad F., Mehmood T., 2013 Proximate analysis: Relative feed values of various forage plants for ruminants investigated in a semi-arid region of Punjab, Pakistan. *Agricultural Sciences* 4(6):302-308.
- Amulejoye F. D., Adebayo O. T., Dada A. A., 2020 Reproductive impact of aqueous extract of *Mangifera indica* leaves on female African catfish *Clarias gariepinus* broodstock. *International Journal of Fisheries and Aquatic Studies* 8(3):657-662.
- Apines-Amar M. J. S., Amar E. C., Faisan Jr. J. P., 2013 Growth, plasma cortisol, liver and kidney histology, and resistance to vibriosis in brown-marbled grouper, *Epinephelus fuscoguttatus* fed onion and ginger. *AACL Bioflux* 6(6):530-538.
- Awad E., Austin B., 2010 Use of lupin, *Lupinus perennis*, mango, *Mangifera indica*, and stinging nettle, *Urtica dioica*, as feed additives to prevent *Aeromonas hydrophila* infection in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases* 33(5):413-420.
- Azizah F., Arief M., Lokaspirnasari W. P., 2020 Influence addition of noni (*M. citrifolia*) in the commercial feed on protein and lipid retentions of sangkuriang catfish (*Clarias* sp.). *IOP Conf. Series: Earth and Environmental Science* 441:012017, 7 p.
- Budijono, Tang U. M., Putra R. M., Nofrizal, 2021 Dynamic of water fertility in Koto Panjang Reservoir, Riau Province, Indonesia. *AACL Bioflux* 14(2):965-975.
- Ceballos-Francisco D., Castillo Y., De La Rosa F., Vásquez W., Reyes-Santiago R., Cuello A., Cuesta A., Esteban M. A., 2020 Bactericidal effect on skin mucosa of dietary guava (*Psidium guajava* L.) leaves in hybrid tilapia (*Oreochromis niloticus* x *O. mossambicus*). *Journal of Ethnopharmacology* 259:1128382.
- Chatzifotis S., Kokou F., Ampatzis K., Papadakis I. E., Divanach P., Dermon C.R., 2008 Effects of dietary caffeine on growth, body composition, somatic indexes, and cerebral distribution of acetyl-cholinesterase and nitric oxide synthase in gilthead sea bream (*Sparus aurata*), reared in winter temperature. *Aquaculture Nutrition* 14(5):405-415.
- Das T. T., Benerjee D., Chakraborty D., Pakhira M. C., Shirvastava B., Kuhad R. C., 2012 Saponin: Role in animal system. *Veterinary World* 5(4):248-254.
- Fawole F. J., Sahu N. P., Nazeemashalul S., Adeoye A. A., 2018 Effect of *Psidium guajava* and *Mangifera indica* leaves extracts on growth, antioxidant and metabolic enzymes activities of *Labeo rohita* fingerlings. *Journal of the Association of Nigerian Fisheries Scientists* 1:40-49.
- Francis G., Harinder P., Makkar, S., Becker, K., 2005 *Quillaja* saponins - a natural growth promoter for fish. *Animal Feed Science and Technology* 121(1-2):147-157.
- Fry J. P., Mailloux N. A., Love D. C., Milli M. C., Cao L., 2018 Feed conversion efficiency in aquaculture: do we measure it correctly? *Environmental Research Letters* 13(2):024017.

- Hoseinifar S. H., Sohrabi A., Paknejad, H., Jafari V., Paolucci M., Van Doan H., 2019 Enrichment of common carp (*Cyprinus carpio*) fingerlings diet with *Psidium guajava*: The effects on cutaneous mucosal and serum immune parameters and immune related genes expression. *Fish and Shellfish Immunology* 86:688-694.
- Ji S. C., Jeong G. S., IM G. S., Lee S. W., Yoo J. H., Takii K., 2007 Dietary medicinal herbs improve growth performance, fatty acid utilization, and stress recovery of Japanese flounder. *Fisheries Science* 73:70-76.
- Jian J., Wu Z., 2004 Influences of traditional Chinese medicine on non-specific immunity of Jian Carp (*Cyprinus carpio* var. Jian). *Fish and Shellfish Immunology* 16(2):185-191.
- Khan M. N., Shahzad K., Chatta A., Sohail M., Piria M., Treer T., 2016 A review of introduction of common carp *Cyprinus carpio* in Pakistan: origin, purpose, impact and management. *Croatian Journal of Fisheries* 74(2):71-80.
- Kristiana V., Mukti A. T., Agustono, 2020 Increasing growth performances of Nile tilapia (*Oreochromis niloticus*) by supplementation of noni *M. citrifolia* fruit extract via diet. *AAFL Bioflux* 13(1):159-166.
- Lin H. Z., Li Z. J., Chen Y. Q., Zheng W. H., Yang K., 2006 Effect of dietary traditional Chinese medicines on apparent digestibility coefficients of nutrients for white shrimp *Litopenaeus vannamei*, Boone. *Aquaculture* 253(1-4):495-501.
- Nhu T. Q., Dam N. P., Hang B. T. B., Bach L. T., Huong D. T. T., Hue B. T. B., Scippo M. L., Phuong N. T., Quetin-Leclercq J., Kestemont P., 2020 Immunomodulatory potential of extracts, fractions and pure compounds from *Phyllanthus amarus* and *Psidium guajava* on striped catfish (*Pangasianodon hypophthalmus*) head kidney leukocytes. *Fish and Shellfish Immunology* 104:289-303.
- Obasa S. O., Alatise S. P., Omoniyi I. T., Alegbeleye W. O., George F. A., 2013 Evaluation of fermented mango (*Mangifera indica*) seed meal in the practical diet of Nile tilapia, (*Oreochromis niloticus*) fingerlings. *Croatian Journal of Fisheries* 71:116-123.
- Olusola S. E., Olorunfemi B. V., 2017 Bioproductive effects of *Clarias gariepinus* fingerlings fed guava (*Psidium guajava*) leaves and drumstick (*Moringa oleifera*) leaves extract supplemented diets. *Applied Topical Agriculture* 22(2):156-165.
- Omitoyin B. O., Ajani E. K., Orisasona O., Bassey H. E., Kareem K. O., Osho F. E., 2019 Effect of guava *Psidium guajava* (L.) aqueous extract diet on growth performance, intestinal morphology, immune response and survival of *Oreochromis niloticus* challenged with *Aeromonas hydrophila*. *Aquaculture Research* 50(7):1851-1861.
- Rachmawati D., Hutabarat J., Samidjan I., Windarto S., 2019 The effects of papain enzyme-enriched diet on protease enzyme activities, feed efficiency, and growth of fingerlings of Sangkuriang catfish (*Clarias gariepinus*) reared in tarpaulin pool. *AAFL Bioflux* 12(6):2177-2187.
- Radhakrishnan S., Saravana B. P., Seenivasan C., Shanthi R., Poongodi R., 2014 Influence of medicinal herbs (*Alteranthera sessilis*, *Eclipta alba* and *Cissus quadrangularis*) on growth and biochemical parameters of the freshwater prawn *Macrobrachium rosenbergii*. *Aquaculture International* 22:551-572.
- Salati A. P., Baghbanzadeh A., Soltani M. Peyghan R., Riazi G., 2011 Effect of different levels of salinity on gill and kidney function in common carp *Cyprinus carpio* (Pisces: Cyprinidae). *Italian Journal of Zoology* 78(3):298-303.
- Salehi-Farsani A., Soltani M., Kamali A., Shamsaie M., 2014 Effect of immune motivator Macrogard and *Spirulina platensis* on some growth, carcass and biochemical indices of stellate sturgeon *Acipenser stellatus*. *AAFL Bioflux* 7(3):137-147.
- Singh G., Ansal M. D., Kaur V. I., 2018 Salinity tolerance and survival of freshwater carp, *Cyprinus carpio* Linn. in inland saline water. *Indian Journal of Ecology* 45(3):598-601.
- Sobral-Souza C. E., Silva A. R. P., Leite N. F., Rocha J. E., Sousa A. K., Costa J. G. M., Menezes I. R. A., Cunha F. A. B., Rolim L. A., Coutinho H. D. M., 2019 *Psidium guajava* bioactive product chemical analysis and heavy metal toxicity reduction. *Chemosphere* 216:785-793.

- Syawal H., Siregar A. R., Yuharmen Y., Effendi I., 2021 Phytochemical analysis and inhibitory effect of *Bruguiera gymnorrhiza* leaf extract on *Aeromonas salmonicida*, *Aeromonas hydrophila*, and *Edwardsiella ictaluri*. *AAFL Bioflux* 14(4):1926-1933.
- Weli A., Al-Kaabi A., Al-Sabahi J., Said S., Hossain A., Al-Riyami S., 2018 Chemical composition and biological activities of the essential oils of *Psidium guajava* leaf. *Journal of King Saud University - Science* 31(4):993-998.
- Yu M. C., Li Z. J., Lin H. Z., Wen G. L., Ma S., 2009 Effects of dietary medicinal herbs and *Bacillus* on survival, growth, body composition, and digestive enzyme activity of the white shrimp *Litopenaeus vannamei*. *Aquaculture International* 17:377-384.
- *** FAO, 2019 Fisheries and aquaculture information and statistics common carp, *Cyprinus carpio*. Food and Agriculture Organization, Rome, Italy.

Received: 09 November 2021. Accepted: 24 January 2022. Published online: 04 May 2022.

Authors:

Irwan Effendi, Faculty of Fisheries and Marine Sciences, University of Riau, Kampus Bina Widya, 28295 Kecamatan Tampan, Indonesia, e-mail: helpingirwan@gmail.com

Dessy Yoswaty, Faculty of Fisheries and Marine Sciences, University of Riau, Kampus Bina Widya, 28295 Kecamatan Tampan, Indonesia, e-mail: dyoswaty13@gmail.com

Henni Syawal, Faculty of Fisheries and Marine Sciences, University of Riau, Kampus Bina Widya, 28295 Kecamatan Tampan, Indonesia, e-mail: hennirizal@gmail.com

Brian Austin, Stirling University, Stirling FK9 4LA, United Kingdom, e-mail: baustin5851@gmail.com

Alastair Robert Lyndon, Heriot-Watt University, Edinburgh, EH 14 4AS, United Kingdom, e-mail: a.r.lyndon@hw.ac.uk

Ronal Kurniawan, Faculty of Fisheries and Marine Sciences, University of Riau, Kampus Bina Widya, 28295 Kecamatan Tampan, Indonesia, e-mail: kurniawanronal09@gmail.com

Sri Wahyuni, Faculty of Fisheries and Marine Sciences, University of Riau, Kampus Bina Widya, 28295 Kecamatan Tampan, Indonesia, e-mail: srayuni17@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Effendi I., Yoswaty D., Syawal H., Austin B., Lyndon A. R., Kurniawan R., Wahyuni S., 2022 Effects of herbal growth promoters on common carp (*Cyprinus carpio*). *AAFL Bioflux* 15(3):1096-1103.