



Heriot-Watt University  
Research Gateway

## Halogenation effects in Intramolecular Furan Diels-Alder reactions

### Citation for published version:

Bebbington, MWP, Rae, R, Zurek, JM & Paterson, M 2013, 'Halogenation effects in Intramolecular Furan Diels-Alder reactions: broad scope synthetic and computational studies', *Organic and Biomolecular Chemistry*, vol. 11, no. 45, pp. 7946 - 7952. <https://doi.org/10.1039/C3OB41616J>

### Digital Object Identifier (DOI):

[10.1039/C3OB41616J](https://doi.org/10.1039/C3OB41616J)

### Link:

[Link to publication record in Heriot-Watt Research Portal](#)

### Document Version:

Publisher's PDF, also known as Version of record

### Published In:

Organic and Biomolecular Chemistry

### Publisher Rights Statement:

CC-BY

### General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [open.access@hw.ac.uk](mailto:open.access@hw.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Halogenation effects in intramolecular furan Diels–Alder reactions: broad scope synthetic and computational studies†

Robert L. Rae, Justyna M. Żurek, Martin J. Paterson\* and Magnus W. P. Bebbington\*

Cite this: *Org. Biomol. Chem.*, 2013, **11**, 7946Received 7th August 2013,  
Accepted 14th October 2013

DOI: 10.1039/c3ob41616j


www.rsc.org/obc

For the first time a comprehensive synthetic and computational study of the effect of halogen substitution on both furan and dienophile for the intramolecular Furan Diels–Alder (IMDAF) reaction has been undertaken. Contrary to our initial expectations, halogen substitution on the dienophile was found to have a significant effect, making the reactions slower and less thermodynamically favourable. However, careful choice of the site of furan halogenation could be used to overcome dienophile halogen substitution, leading to highly functionalised cycloadducts. These reactions are thought to be controlled by the interplay of three factors: positive charge stabilisation in the transition state and product, steric effects and a dipolar interaction term identified by high level calculations. Frontier orbital effects *do not appear to make a major contribution* in determining the viability of these reactions, which is consistent with our analysis of calculated transition state structural data.

## Introduction

The intramolecular furan Diels–Alder (IMDAF) reaction has been the subject of numerous studies over the last 30 years.<sup>1</sup> The reaction is very sensitive to both furan and dienophile substitution,<sup>1b</sup> which has rendered it a useful probe of more general fundamental effects on pericyclic reactions. The reaction has also been employed as a key complexity-generating step in a number of target syntheses.<sup>2</sup>

Halogen substitution was first examined more than 25 years ago,<sup>3</sup> but not until a series of elegant experimental and computational studies by Padwa and Houk<sup>3d,4</sup> (Scheme 1) did the effect of halogen substitution on furan become clear – IMDAF reactions with halogenated furans were faster and more exergonic than those with non-halogenated furans. Although halogen substitution at the 3-position of 2-furanyl amides was found to promote the cycloaddition largely for conformational reasons, the electronic effect of halogen substitution was most profound at the 5-position (Scheme 1). This was attributed to greater stabilisation of the partial positive



**Scheme 1** Padwa's demonstration of activation by halogen substitution in the intramolecular furan Diels–Alder reaction.

charge at the 5-carbon in both transition state and cycloadduct than in the starting material, as a result of the change in hybridisation from  $sp^2$  to  $sp^3$ .<sup>4</sup> The effect was calculated to be greatest for fluorine, although this is yet to be demonstrated experimentally, perhaps reflecting current difficulty in the preparation of simple  $\alpha$ -fluorofurans.<sup>5</sup> Chlorine or bromine substitution is, however, sufficient to have a profound promoting effect in the cases where a true comparison is available with the corresponding non-halogenated substrates.<sup>3</sup>

Although considerable progress has been made in understanding these substituent effects, there remain some unresolved issues. For example, although the change in hybridisation at carbon has been identified as one of the

Institute of Chemical Science, Heriot-Watt University, Riccarton, Edinburgh, EH14 4AS, UK. E-mail: m.w.p.bebbington@hw.ac.uk; Fax: +44 (0)131 451 3180; Tel: +44 (0)131 451 8071

†Electronic supplementary information (ESI) available: Experimental procedures, characterisation data, <sup>1</sup>H and <sup>13</sup>C NMR spectra for all new compounds; further computational experiments on model systems and full discussion of the CBS methodologies used, results for reactions of 1a–z at 298 K, dipolar interaction, selected FMO and transition state structural data. See DOI: 10.1039/c3ob41616j



contributing factors in substrate activation by halogen substitution at the 5-position, significant activation is still predicted by halogen substitution at the 4-position,<sup>4</sup> where no change in hybridisation occurs during the reaction and no significant conformational effects would be anticipated. At the beginning of our studies we also noted that very little work<sup>6</sup> had been done on IMDAF substrates involving halogenation on the dienophile, where change in hybridisation from  $sp^2$  to  $sp^3$  at the halogen-bearing carbon also occurs. We thus undertook to investigate these fundamental issues through examination of suitable cycloaddition precursors.

## Results and discussion

We prepared various classes of IMDAF substrates that would allow us to probe the significance of halogenation with respect to variations in electronics that would also affect the viability of the cycloaddition. Thus, we first prepared substrates closely related to Padwa's original furoic acid-derived examples that would be used essentially for control experiments and to compare directly the effect of halogenation of the diene and the dienophile in turn.









Substrates **1a,b** and **1d** are directly analogous to those prepared by Padwa and co-workers<sup>3d,7</sup> and show similar reactivity, consistent with the early work by that group. Bromine substitution at any of the 3-, 4- and 5-positions increases the rate and yield of the reaction compared to the parent unsubstituted system **1a** (Table 1).<sup>8</sup>

All of these reactions, and subsequent ones described herein, are optimised for maximum isolated yield of cycloadduct – longer reaction times than those listed led to some decomposition. The effect of the *N*-substituent on the cycloaddition rates of furanyl amides has been studied and found to vary according to the steric bulk of the substituent, rather than the relative populations of the different amide rotamers.<sup>9</sup> Our results are consistent with this work, which suggests that the *N*-phenyl substituent would lead to a slightly slower forward reaction rate (by a factor of ~3) than the *N*-benzyl one.

In order to determine whether the halogenation effects showed any variation with a change in the electron-demand of the IMDAF reaction, we prepared substrates **1e–1h** (Table 2), bearing electron-deficient alkenes, from *N*-phenyl furfuryl-amines (see ESI† for details). In these cases, the difference between the 5-bromo and 4-bromo substrates **1f** and **1g** was marginal, although both showed higher reactivity than the non-halogenated substrate. The 3-bromo derivative **1h** was once again the most reactive of the series, with the cycloaddition proceeding at noticeable rate even at room temperature, suggesting that conformational effects play an important part in promoting this reaction whether or not the carbonyl group is attached directly to the furan ring.

We then elected to study substrates that were halogenated on the dienophile. Chlorine was chosen in the majority of cases because the three isomeric *N*-chloroallyl anilines are readily prepared from the commercially available chloroallyl chlorides.<sup>10</sup> The 2-bromoallyl derivative was also easily

**Table 1** Control experiments with furanyl amides

Precursor <b>1</b>	Cycloadduct <b>2</b>	Time/h	Ratio <b>2/1</b>	Yield <b>2</b> <sup>a</sup> /%
 <b>1a</b>	 <b>2a</b>	24	86 : 14	62
 <b>1b</b>	 <b>2b</b>	24	100 : 0	100
 <b>1c</b>	 <b>2c</b>	24	100 : 0	77
 <b>1d</b>	 <b>2d</b>	24	100 : 0	77

<sup>a</sup> All reactions carried out at 110 °C in toluene.

synthesised. Clearly, a change in hybridisation at the halogen-bearing carbon, from  $sp^2$  to  $sp^3$ , is a necessary feature of the IMDAF reaction for substrates **1i–1l** and **1n–1q** (Table 3),<sup>11</sup> and we initially hypothesised that this effect would promote the cycloaddition with respect to the non-halogenated substrates as was observed for the earlier work with halogenated furans.<sup>3</sup> However, our results indicate the opposite – in all cases the reaction is slower when the halogen is present, although this effect is greater for the amides **1i–1l** than for the *N*-Boc derivatives **1m–1q**. This suggests that steric effects are more important when the dienophile is halogenated than when the furan is, since transannular interactions are expected to be more significant in these cases. The mass balance was not completely recovered in these optimised reactions, so in order to verify that the retardation in rate was genuine, substrates **1a** and **1h** were heated for only 3 h. The conversion to **2a** was 37% and to **2h** was 15%. The crude reaction mixtures were analysed by <sup>1</sup>H NMR using a known quantity of cyclohexane as an internal standard and no significant mass loss was observed for these shorter reactions,<sup>12</sup> supporting our initial supposition of a decrease in rate for the halogenated system. In particular, the use of a *Z*-chloroalkene as dienophile (in **1k** and **1p**) resulted in only 11–12% conversion after 24 h at 110 °C. This is somewhat surprising as the chlorine is installed on the normally less hindered face of the product. The lower reactivity of the *Z*-dienophiles compared to the *E*-dienophiles is in contrast to



**Table 2** Cycloadditions with acylated furfuryl amines

Precursor 1	Cycloadduct 2	Time/h	Ratio 2/1	Yield 2 <sup>a</sup> /%
		24	93:7	62
		2	90:10	70
		12	>95:5	76
		6	100:0	83

<sup>a</sup> All reactions carried out at 110 °C in toluene.

that observed in some other IMDA reactions using acyclic dienes.<sup>13</sup> Indeed, reaction of the *E*-substrates **1l** and **1q** and also the 2-haloallyl systems **1i**, **1j**, **1n** and **1o** requires the halogen to be on the more hindered face of the [2.2.1]-bicyclic ring system.

We next decided to examine frontier orbital effects using DFT (B3LYP/6-31G(d))<sup>14</sup> calculations to attempt to explain these results. Furan is normally considered to be an electron-rich diene in Diels–Alder reactions, such that the important frontier molecular orbital interaction is HOMO(furan)–LUMO(dienophile).<sup>1b</sup> Many examples of IMDAF are consistent with this, but in a number of cases the cycloaddition occurs still occurs readily when, as here, the furan bears an electron-withdrawing carbonyl group, so there has been some debate regarding frontier orbital effects in IMDAF.<sup>3a,b,d,15</sup> The effect of halogen substitution on frontier orbital energies is less clear cut,<sup>16</sup> due to a mixture of electron-donating field/mesomeric and electron-withdrawing inductive effects. The similar trends observed for substrates **1i–1l** and **1n–1q**, which would be expected to have rather different FMO properties, suggest that frontier orbital effects are not overriding source of the observed differences in reactivity. Our calculations<sup>17</sup> predict that both furan and 2-chlorofuran react with ethylene in a

**Table 3** Cycloadditions with haloalkene dienophiles

Precursor 1	Cycloadduct 2	Time/h	Ratio 2/1	Yield 2 <sup>a,b</sup> /%
		24	40:60	27 (68)
		24	50:50	44 (88)
		24	12:88	7 (58)
		24	63:37	28 (44)
		24	54:46	46 (86)
		96	31:69	15 (51)
		24	48:52	28 (58)
		48	11:89	7 (63)
		24	53:47	31 (58)

<sup>a</sup> All reactions carried out at 110 °C in toluene. <sup>b</sup> Combined recovery (of **1** + **2**) in parentheses.



normal electron-demand sense. The presence of the Cl atom only results in a very minor change (<0.1 eV) in the HOMO–LUMO energy gap in these systems. Examination of the relevant orbitals in the more complex intramolecular reactions indicates that non-halogenated substrates **1e** and **1m** react with normal demand, whereas **1a**, with an electron-withdrawing group on furan, reacts with inverse demand. Halogenation does not appear to change the nature of the FMO interactions, and produces only minor changes in orbital energies (0.1–0.4 eV) across most of the substrates studied.<sup>17</sup> The exceptions to this are **1e–1h**, where the FMO energy gap *increases* by 1.1–1.3 eV upon furan halogenation, although the halogenated systems appear to react more rapidly. Similarly, comparison of the HOMO–LUMO energy gaps in inverse demand substrates **1a**, **1i** and **1j** suggests that halogenation *decreases* the energy gap (by 0.3–0.4 eV), even though **1a** undergoes cycloaddition fastest of these three substrates. Intriguingly, very similar experimental and FMO trends are found in normal demand substrates **1m**, **1n** and **1o**, where the FMO energy gap is predicted to decrease for halogenated systems **1n** and **1o**, even though the cycloaddition appears marginally faster in the non-halogenated substrate **1m**. Therefore, *dienophile halogenation decreases the FMO energy gap for both normal- and inverse-demand systems*. These results therefore support the hypothesis that *FMO interactions are not the most important factor* in controlling these reactions. This is consistent with a less exergonic reaction than is usual for a Diels–Alder cycloaddition (e.g. with a non-aromatic diene),<sup>18</sup> giving rise to a less reactant-like transition state that is not well approximated by examination of the orbital energies of the starting materials. An examination of the calculated transition state and product structures is also supportive of later transition states among the less exergonic reactions (*vide infra*).

Our results could instead be explained if a dipolar interaction between the C–X and furan C–O bonds served to raise the energy of the transition state for **1k** and **1p** and had the opposite effect in the other substrates. Indeed, such a possibility is supported by the calculations. Using a simple approximation for the interaction of two (weak) dipoles we have calculated the orientational dependence of the dipolar interactions of the C–O and the dienophile C–X dipoles at the transition state structures of certain reactions. Table S4 of the ESI† presents these values for reactions of **1i–1l** and **1p–1q**. The X–C–C–O dihedral angle differs significantly in the transition states for reactions of **1k** and **1l**, which leads to a significant difference in the dipolar interaction term. There is some repulsion present in the structure of the transition states for reactions of **1k** and **1p** thus leading to higher Diels–Alder activation energy. Transition states for reactions of **1i**, **1j**, **1l** and **1q** acquire more attraction interaction, leading to lower potential energy for the transition state, and thus lower activation energy for Diels–Alder reaction.

The 2-bromoallyl derivatives **1j** and **1o** undergo cycloaddition to a slightly greater extent than the corresponding 2-chloroallyl substrates **1i** and **1n**, and indeed any of the other chlorinated precursors, despite the fact that the bulky bromine

atom is attached to a crowded tertiary centre. This indicates that the increased steric requirements of the halogenated alkene, positive charge stabilisation at halogenated carbon in the transition state, or dipolar interactions do not account for these observations, since all of these factors should be more significant for Cl-substituted systems. It is presently unclear why the brominated systems react marginally more rapidly than the chlorinated ones.

In order to compare experimentally the beneficial effect of furan halogenations *versus* the detrimental effect of dienophile halogenation, we finally prepared dihalogenated examples **1r–1z** leading to adducts **2r–2z** (Table 4). In general, greater reactivity was observed than in substrates **1i–1l**. Interestingly, bromination at the 5-position of furan was not always more effective in promoting the cycloaddition than at the 4-position (substrates **1s** vs. **1v**),<sup>4</sup> and indeed the 5-bromo-Z-chloro substrate **1s** showed little more reactivity than the analogous compound **1k** lacking a brominated furan, presumably for steric reasons. The 3-bromo derivatives **1y** and **1z** show the highest reactivity, which is consistent with our earlier results. Substrate **1x** is anomalous given the reactivity of **1r** and **1u**, and illustrates the complexity of the interaction between the steric and electronic factors controlling these reactions. Careful adjustment of the bromination site is thus sufficient to override the detrimental effect of the dienophile halogenations, demonstrating the power of this substitution in facilitating previously difficult cycloadditions.

## Computation



















All calculations were performed with the Gaussian09 program (versions a and c).<sup>19</sup> The Complete Basis Set (CBS) set of methods of Petersson and co-workers was used to obtain accurate thermochemical data.<sup>20–23</sup> The specific extrapolation used here (Table 5) is the CBS-QB3 one. Details of others used are given in ESI (Table S1†) for full comparison. All calculations were performed in the gas phase. Free energies with CBS models were calculated for two temperatures: 298 K and 383 K (the temperature of the experimental studies). The results for CBS-QB3 model of chosen Diels–Alder reactions for 383 K are presented in Table 5.<sup>24</sup>

Comparison of the calculated thermodynamic and experimental data is somewhat complicated by decomposition of many of the adducts at the extended reaction times (typically a number of days) sometimes needed for the reactions to go to completion. All of the reactions studied are predicted to be exergonic in nature (Fig. 1 and Table 5), although there is significant variation in both activation and overall reaction free energies.

Those reactions with calculated higher activation energies predominantly show, as a general trend, lower conversions to product over the timescale of the reactions, where directly comparable. However, a quantitative correlation between activation barrier and yield was not observed. A plot of the calculated activation energies (Fig. 2 and Table 5) is colour-coded to show grouping of the different types of substrate. Thus, precursors with halogenated furans (**1b–d**, **1f–h**) possess lower



**Table 4** Dihalogenated systems



Precursor 1	Cycloadduct 2	Time/h	Ratio 2/1	Yield 2 <sup>a,b</sup> /%
		24	85 : 15	61 (72)
		24	14 : 86	12 (85)
		8	83 : 17	60 (72)
		24	78 : 22	50 (64)
		24	40 : 60	31 (78)
		24	83 : 17	54 (65)
		5	40 : 60	20 (50)
		16	87 : 13	67 (77)
		7	94 : 6	75 (80)

<sup>a</sup> All reactions conducted at 110 °C in toluene. <sup>b</sup> Combined recovery of 1 + 2 in parentheses.

**Table 5** CBS-QB3 calculated thermochemical data of IMDAF reactions at 383 K<sup>a</sup>

Reaction	$\Delta_r H^\circ$	$\Delta_r G^\circ$	$\Delta_{\text{activ}} H$	$\Delta_{\text{activ}} G$
1a → 2a	-10.6	-5.2	19.4	23.8
1b → 2b	-14.6	-9.1	17.9	22.4
1c → 2c	-14.0	-8.7	17.6	21.7
1d → 2d	-14.0	-8.1	16.4	21.3
1e → 2e	-17.2	-9.9	15.6	21.8
1f → 2f	-20.7	-14.5	15.0	19.9
1g → 2g	-19.8	-13.7	14.5	19.1
1h → 2h	-18.8	-12.8	14.7	19.5
1i → 2i	-9.6	-4.0	20.5	25.0
1j → 2j	-9.9	-4.3	20.4	25.0
1k → 2k	-11.6	-5.9	19.4	23.9
1l → 2l	-12.1	-5.3	18.1	23.9
1m → 2m	-8.2	-2.7	19.9	24.5
1n → 2n	-8.2	-2.4	20.4	25.2
1o → 2o	-8.6	-3.1	20.1	25.2
1p → 2p	-7.2	-0.1	21.5	27.6
1q → 2q	-8.4	-2.7	20.0	24.7
1r → 2r	-13.5	-7.7	18.8	23.4
1s → 2s	-13.7	-8.0	18.4	22.9
1t → 2t	-14.9	-9.2	20.6	21.6
1u → 2u	-13.0	-7.3	18.7	23.4
1v → 2v	-14.4	-8.6	17.7	22.4
1w → 2w	-13.8	-7.9	17.6	22.6
1x → 2x	-12.6	-6.2	18.7	24.0
1y → 2y	-15.3	-10.1	15.6	19.7
1z → 2z	-14.6	-8.5	16.2	21.4

<sup>a</sup> All energies reported in kcal mol<sup>-1</sup>.

**Fig. 1** Scatter plot of Gibbs free energies of reaction (in kcal mol<sup>-1</sup>) for systems a–z (black = unhalogenated; red = halofuran substrate; blue = haloalkene substrate; yellow = dihalogenated substrate).**Fig. 2** Scatter plot of Gibbs free energies of activation (in kcal mol<sup>-1</sup>) for systems a–z.






Fig. 3 Scatter plot of percentage change in C–C bond distances from transition state to product.

activation barriers for cycloaddition ( $19.1$ – $22.4$  kcal mol $^{-1}$  with mean value  $20.6$  and standard deviation  $\pm 1.3$ ) than those with halogenated dienophiles (**1i–l**, **1n–q**) which lie between  $23.9$  and  $27.5$  kcal mol $^{-1}$  (mean  $25.1 \pm 1.2$ ). In turn, those substrates with halogens on both furan and dienophile (**1r–z**) have activation barriers that are intermediate in value ( $21.4$  to  $23.4$  kcal mol $^{-1}$ , mean  $22.4 \pm 1.3$ ). Substrate **1y** is anomalous, with a low calculated activation energy ( $19.7$  kcal mol $^{-1}$ ), but experimentally it shows little reactivity.

In order to probe further the validity of our conclusions from our studies on possible frontier orbital effects, we elected to quantify the “product-likeness” of the transition states in terms of the fractional shortening<sup>25</sup> of the newly forming C–C  $\sigma$ -bonds (marked as *a* and *b* in Fig. 3; the corresponding calculated numerical data are in Table 6) in moving from transition state to product. A similar grouping to that seen in both activation energies and free energies of reaction is observed. Thus, the more exergonic the reactions (see Fig. 1 and 3), the earlier their transition states, so that the percentage contraction of the C–C distance in going from transition state to product is greater. Conversely, those reactions that exhibit a lower percentage contraction are predicted to be less exergonic, consistent with later transition states. These data are thus supportive of our earlier conclusions on FMO energies which are not found to correlate with the reactivities we observe. FMO energies are thus likely to be less important contributors to transition state energies for reactions with later transition states than is usual for Diels–Alder cycloadditions, such as those discussed in our work.

## Conclusions

We have investigated the effect of both furan and dienophile halogenations in IMDAF reactions. Halogenation of the furan is found to favour the cycloaddition, irrespective of the electronic effects of the other furan and dienophile substituents.

Table 6 Calculated transition state and product C–C bond distances<sup>a</sup> for systems a–z

System	Transition state		Product	
	C–C distance <i>a</i>	C–C distance <i>b</i>	C–C bond length <i>a</i>	C–C bond length <i>b</i>
a	2.029	2.178	1.558	1.571
b	2.044	2.199	1.558	1.565
c	2.053	2.179	1.556	1.568
d	2.004	2.227	1.555	1.570
e	2.035	2.223	1.556	1.569
f	2.056	2.241	1.556	1.563
g	2.061	2.221	1.553	1.566
h	2.022	2.259	1.553	1.567
i	2.064	2.153	1.563	1.570
j	2.067	2.152	1.562	1.570
k	1.995	2.225	1.557	1.569
l	1.987	2.231	1.555	1.570
m	2.095	2.111	1.576	1.572
n	2.137	2.076	1.581	1.570
o	2.139	2.077	1.579	1.571
p	2.061	2.137	1.574	1.570
q	2.067	2.146	1.574	1.570
r	2.079	2.165	1.563	1.565
s	1.950	2.332	1.553	1.583
t	2.031	2.235	1.554	1.572
u	2.095	2.146	1.562	1.566
v	2.016	2.222	1.554	1.567
w	1.986	2.253	1.552	1.570
x	2.086	2.150	1.565	1.566
y	1.962	2.278	1.554	1.569
z	1.968	2.275	1.553	1.570

<sup>a</sup> All distances (*a* and *b* are defined in Fig. 3) in angstroms.

Halogenation of the dienophile was found to lead to slower reactions than those with the corresponding non-halogenated dienophiles in all cases studied. Frontier orbital effects do not appear to have a huge influence on these reactions, and the results have been attributed to a combination of positive charge stabilisation, simple steric effects, and a contribution from a dipolar interaction between the furan C–O and the dienophile C–X dipoles, supported by computation. This varies with the dihedral angle between the two bonds in the transition states, suggesting a partial explanation for the difference in reactivity between analogous systems with differing alkene geometries. The use of dihalogenated substrates can provide a means of overriding the effect of dienophile halogenations, leading to synthetically useful yields of highly functionalised cycloadducts with considerable potential for further transformation.

The authors thank the ERC (for funding for M. J. P.) and the EPSRC (for a DTA for R. L. R.; for funding for J. M. Ž. and for use of the EPSRC National Mass Spectrometry Facility for mass spectra).

## Notes and references

- (a) For a review of the intramolecular Diels–Alder reaction, see: E. Ciganek, *Org. React.*, 2004, 1–371, Wiley Online Library; (b) For a comprehensive review of the furan Diels–



- Alder reaction, see: C. O. Kappe, S. S. Murphree and A. Padwa, *Tetrahedron*, 1997, **53**, 14179.
- For a review of the use of the product oxanorbornenes in natural product synthesis, see: (a) P. Vogel, J. Cossy, J. Plumet and O. Arjona, *Tetrahedron*, 1999, **55**, 13521; For selected more recent examples, see: (b) E. N. Pitsinos, N. Athinaios and V. P. Veroniki, *Org. Lett.*, 2012, **14**, 4666; (c) C.-C. Wang and W.-D. Z. Li, *J. Org. Chem.*, 2012, **77**, 4217; (d) P. Fischer, M. Gruner, A. Jaeger, O. Kataeva and P. Metz, *Chem.-Eur. J.*, 2011, **17**, 13334; (e) K. Tanino, M. Takahashi, Y. Tomata, H. Tokura, T. Uehara, N. Takashi and M. Miyashita, *Nat. Chem.*, 2011, **3**, 484; (f) F. R. Petronijevic and P. Wipf, *J. Am. Chem. Soc.*, 2011, **133**, 7704; Y.-P. Xue and W.-D. Z. Li, *J. Org. Chem.*, 2011, **76**, 57; (g) M. B. O'Keefe, D. M. Mans, D. E. Kaelin and S. F. Martin, *J. Am. Chem. Soc.*, 2010, **132**, 15528; (h) S. Mitsuru and Y. Hayashi, *Eur. J. Org. Chem.*, 2007, 3783; (i) S. M. Sparks, C.-L. Chen and S. F. Martin, *Tetrahedron*, 2007, **63**, 8619; (j) G. E. Morton and A. G. M. Barrett, *Org. Lett.*, 2006, **8**, 2859; (k) A. Padwa and J. D. Ginn, *J. Org. Chem.*, 2005, **70**, 5197; (l) M. Takadoi, T. Katoh, I. Tadashi and S. Terashima, *Tetrahedron*, 2002, **58**, 9903; (m) S. Mitsuru, J. Yamaguchi, H. Kakeya, H. Osada and Y. Hayashi, *Angew. Chem., Int. Ed.*, 2002, **41**, 3192.
  - For earlier studies on IMDAF reactions with halogenated furans, see: (a) Z. Klepo and K. Jakopcic, *J. Heterocycl. Chem.*, 1987, **24**, 1787; (b) A. D. Mance, M. Sindler-Kulyk, K. Jakopcic, A. Hergold-Brundic and A. Nagl, *J. Heterocycl. Chem.*, 1997, **34**, 1315; (c) K. R. Crawford, S. K. Bur, C. S. Straub and A. Padwa, *Org. Lett.*, 2003, **5**, 3337; (d) A. Padwa, K. R. Crawford, C. S. Straub, S. N. Pieniazek and K. N. Houk, *J. Org. Chem.*, 2006, **71**, 5432.
  - S. N. Pieniazek and K. N. Houk, *Angew. Chem., Int. Ed.*, 2006, **45**, 1442.
  - Procedures for the preparation of  $\alpha$ -fluorofurans tend to be low-yielding and lacking in generality – for a review of fluorofuran synthesis, see: O. Serdyuk, A. Butin and V. Abaev, *J. Fluorine Chem.*, 2010, **131**, 296.
  - For examples of IMDAF reactions using haloalkene dienophiles, see: (a) M. Karaarslan and A. Demircan, *Asian J. Chem.*, 2007, **19**, 2999; (b) A. Demircan, M. Karaarslan and E. Turac, *Heterocycl. Commun.*, 2006, **12**, 233. No comparison is made with non-halogenated systems herein and the halogen is always attached to a tertiary centre. For examples of IMDAF reactions with chloroacrylate dienophiles, see: (c) M. E. Jung and L. J. Street, *J. Am. Chem. Soc.*, 1984, **106**, 8327; (d) M. E. Jung and L. J. Street, *Heterocycles*, 1988, **27**, 45. A slight retardation of the cycloaddition is observed for chloroacrylates compared with acrylates.
  - No substrate analogous to **1c**, with a sole Br atom in the 4-position has ever been studied prior to this work.
  - The conversions of substrates **1a** and **1b** after only 1 h at 110 °C to the cycloadducts **2a** and **2b** were 19% and 63% respectively, as judged by <sup>1</sup>H NMR spectroscopy.
  - M. Nakamura, I. Takahashi, S. Yamada, Y. Dobashi and O. Kitagawa, *Tetrahedron Lett.*, 2011, **52**, 53.
  - J. Barluenga, F. Foubelo, F. J. Fañanás and M. Yus, *J. Chem. Soc., Perkin Trans. 1*, 1989, 553.
  - Somewhat surprisingly, non-halogenated control substrate **1m** has not previously been prepared.
  - Some decomposition was observed (NMR) in reactions heated for 24 h or longer. Some mass loss is also thought to have occurred in chromatography – separation was difficult and some mixed fractions were obtained in a number of cases, the contents of which obviously are not included in the isolated yields.
  - T. N. Cayzer, M. N. Paddon-Row, D. Moran, A. D. Payne, M. S. Sherburn and P. Turner, *J. Org. Chem.*, 2005, **70**, 5561.
  - Calibration calculations using larger bases on selected systems did not alter the conclusions from this part of the study.
  - T. Takebayashi, N. Iwasawa and T. Mukaiyama, *Chem. Lett.*, 1983, 1107.
  - R. Robiette, J. Marchand-Brynaert and D. Peeters, *J. Org. Chem.*, 2002, **67**, 6823.
  - See ESI.†
  - Calculations in ref. 4 indicate that Diels–Alder reactions with non-aromatic dienes are considerably more exergonic than those with furan. Hammond's postulate would hence suggest that FMO energies of the starting materials would be less predictive of reactivity in the case of furan cycloadditions, which would be expected to have more product-like transition states.
  - M. J. T. Frisch, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, N. J. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, *Gaussian 09, Revision A.02*, 2009.
  - M. R. Nyden and G. A. Petersson, *J. Chem. Phys.*, 1981, **75**, 1843.
  - J. W. Ochterski, G. A. Petersson and J. A. Montgomery, *J. Chem. Phys.*, 1996, **104**, 2598.
  - J. A. Montgomery, M. J. Frisch, J. W. Ochterski and G. A. Petersson, *J. Chem. Phys.*, 1999, **110**, 2822.
  - J. A. Montgomery, M. J. Frisch, J. W. Ochterski and G. A. Petersson, *J. Chem. Phys.*, 2000, **112**, 6532.
  - For analogous data calculated at 298.15 K, see the ESI.†
  - For a table detailing these values, see Table S5 in the ESI.†

