



Heriot-Watt University
Research Gateway

Analysing the polarisation of the CMB with spin scale-discretised wavelets

Citation for published version:

Leistedt, B, McEwen, JD, Buttner, M, Peiris, H, Vandergheynst, P & Wiaux, Y 2015, 'Analysing the polarisation of the CMB with spin scale-discretised wavelets', Paper presented at BASP Frontiers 2015, Lausanne, Switzerland, 25/01/15 - 30/01/15. <<http://www.baspfrontiers.org/proceedings.pdf>>

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Analysing the polarisation of the CMB with spin scale-discretised wavelets

Boris Leistedt^{*}, Jason D. McEwen[†], Martin Büttner^{*}, Hiranya V. Peiris^{*}, Pierre Vandergheynst[‡], and Yves Wiaux[§].

^{*} Department of Physics and Astronomy, University College London (UCL), London WC1E 6BT, UK

[†] Mullard Space Science Laboratory (MSSL), University College London (UCL), Surrey RH5 6NT, UK

[‡] Institute of Electrical Engineering, Ecole Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland

[§] Institute of Sensors, Signals & Systems, Heriot-Watt University, Edinburgh EH14 4AS, UK

Abstract—We discuss a new scale-discretised directional wavelet transform to analyse spin signals defined on the sphere, in particular the polarisation of the cosmic microwave background (CMB).

I. SPIN SCALE-DISCRETISED WAVELETS ON THE SPHERE

We design a directional scale-discretised wavelet transform to analyse the directional features of signals of arbitrary spin on the sphere. Following Refs. [1]–[4], the spin-discretised wavelets ${}_s\Psi^j$ on the sphere are defined in harmonic space as ${}_s\Psi_{\ell m}^j \equiv \kappa^j(\ell)\xi_{\ell m}$, where $\kappa^j(\ell)$ characterises the angular localisation of the wavelets (for the j th scale), while $\xi_{\ell m}$ controls their directionality. Fig. 1 shows examples of wavelets obtained with this construction.




Fig. 1. Spin 0 and 2 wavelets constructed with angular $L = 512$, azimuthal band-limit $N = 15$, and tilling parameters $B = 2$ and $J_{\min} = 2$.

The wavelet transform of a spin signal ${}_s f$ on the sphere is given by the directional convolution with the wavelets. Provided the wavelets satisfy an admissibility property, the original signal can be synthesised exactly from its wavelet coefficients. More details about this transform can be found in Ref. [4].

II. APPLICATION TO THE CMB

The polarisation of the CMB is currently an intense avenue of research, since it may reveal signatures of primordial gravitational waves and a glimpse on the initial conditions of the universe. However, the physical quantities necessary for these investigations – namely the E- and B-modes of the polarisation, or gradient and curl modes – are not directly available. They are obtained by reducing and transforming frame-dependent observables: the Q and U “local” polarisation measured on the sky. This Q-U to E-B transformation is unambiguous when dealing with data covering the entire sky. However, when CMB observations cover fractions of the sky only, E-B reconstruction is imperfect near the boundaries of the observation mask, causing leakage and potential biases in the recovered E- and B-mode maps. This issue is traditionally addressed by smoothing and extending the mask to remove boundary regions where the leakage is important. We develop a method exploiting the novel directional spin wavelet transform to decrease the leakage near the mask boundaries and obtain a more accurate E-B reconstruction.

First, we compute the wavelet coefficients of the observable $Q+iU$ using our *spin* wavelet transform. Second, we deal with the partial sky coverage by masking the data in spin wavelet space. Third, we apply to the real and imaginary parts of these wavelet coefficients a *scalar* inverse wavelet transform. It can be shown that this yields estimates of the E and B signals provided the wavelets of this inverse transform are spin-lowered versions of those used in the first transform Ref. [4]. This new method to estimate the E- and B-mode contributions from Q and U observations can significantly reduce the E-B leakage by exploiting improved masking in spin wavelet space.

We anticipate our novel spin wavelet transform will be of general use in analysing CMB polarisation data, beyond this first application to E-B separation. A more detailed description of spin scale-discretised wavelets, fast algorithms, and a rigorous evaluation of their performance will be given in a series of forthcoming articles.

REFERENCES

- [1] Y. Wiaux, J. D. McEwen, P. Vandergheynst, and O. Blanc, “Exact reconstruction with directional wavelets on the sphere,” *MNRAS*, vol. 388, no. 2, pp. 770–788, 2008.
- [2] B. Leistedt, J. D. McEwen, P. Vandergheynst, and Y. Wiaux, “S2LET: A code to perform fast wavelet analysis on the sphere,” *A&A*, vol. 558, no. A128, pp. 1–9, 2013.
- [3] J. D. McEwen, P. Vandergheynst, and Y. Wiaux, “On the computation of directional scale-discretized wavelet transforms on the sphere,” in *SPIE Wavelets and Sparsity XV*, 2013.
- [4] J. D. McEwen, M. Büttner, B. Leistedt, H. V. Peiris, P. Vandergheynst, and Y. Wiaux, “On spin scale-discretised wavelets on the sphere for the analysis of CMB polarisation,” in *Proceedings IAU Symposium 306*, 2014, pp. 119–126.