



Heriot-Watt University  
Research Gateway

## The impact of unilateral brain damage on weight perception, sensorimotor anticipation, and fingertip force adaptation

### Citation for published version:

Buckingham, G, Biekiewicz, M, Rohrbach, N & Hermsdörfer, J 2015, 'The impact of unilateral brain damage on weight perception, sensorimotor anticipation, and fingertip force adaptation', *Vision Research*, vol. 115, no. Part B, pp. 231-237. <https://doi.org/10.1016/j.visres.2015.02.005>

### Digital Object Identifier (DOI):

[10.1016/j.visres.2015.02.005](https://doi.org/10.1016/j.visres.2015.02.005)

### Link:

[Link to publication record in Heriot-Watt Research Portal](#)

### Document Version:

Early version, also known as pre-print

### Published In:

Vision Research

### General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [open.access@hw.ac.uk](mailto:open.access@hw.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

1 The impact of unilateral brain damage on weight perception, sensorimotor anticipation, and  
2 fingertip force adaptation

3

4 Gavin Buckingham<sup>1</sup>

5 Marta Bierkiewicz<sup>2</sup>

6 Nina Rohrbach<sup>2</sup>

7 Joachim Hermsdörfer<sup>2</sup>

8

9 1. Department of Psychology, Heriot-Watt University, Edinburgh, UK

10 2. Department of Sport and Health Sciences, Technische Universität München, Munich, Germany

11

12 Keywords: Brain damage; object lifting; grip force; load force; material weight illusion, size weight  
13 illusion

14

15 Corresponding author

16 Gavin Buckingham

17 Department of Psychology

18 School of Life Sciences

19 Heriot-Watt University

20 Riccarton Campus

21 Edinburgh, UK

22 EH14 4AS

23 Email: gav.buckingham@gmail.com

24 Phone: +441314514643

25

1 Abstract

2 Damage to the left parietal cortex can lead to apraxia – a selective deficit in tool use and action  
3 planning. There is conflicting evidence as to whether this disorder affects more fundamental motor  
4 parameters, such as applying the appropriate forces to lift objects based upon how heavy they look.  
5 Here we examined how individuals with left and right-lateralized brain damage lift and perceive the  
6 weight of objects of the same mass which vary in their size and material properties. No clear  
7 differences emerged between the groups in terms of how visual material properties affected their  
8 perceptions of object weight or their initial application of grip and load forces. There was, however,  
9 some evidence that unilateral brain injury impaired the use of size cues for the parameterization of  
10 grip forces.

11

## 1 Introduction

2 A large body of evidence has indicated that our conscious perception of an object's weight does not  
3 accurately reflect its physical mass. This discrepancy between experience and veridical reality is  
4 readily demonstrated through various weight illusions, where objects of identical mass feel different  
5 weights to one another due to variations in other physical properties. In the most famous of these,  
6 the size-weight illusion (SWI), small objects feel heavier than large objects which have been adjusted  
7 to have the same mass (Charpentier, 1891; Murray, Ellis, Bandomir, & Ross, 1999). Similarly, in the  
8 material-weight illusion (MWI), objects which appear to be made from low-density materials are  
9 judged as feeling heavier than objects of the same mass which appear to be made from high-density  
10 materials (Ellis & Lederman, 1999; Seashore, 1899). These illusions are persistent, cognitively  
11 impenetrable, and appear to be a consequence of how we integrate our prior expectations with  
12 sensory input (i.e., the heavy-feeling objects feel heavy because lifters expected them to be lighter  
13 than they actually are – for review see Buckingham, in press).

14 Expectations of object heaviness do not just affect perceptual judgements, but also influence the  
15 way that lifters interact with objects. Due to the predictive way in which objects are lifted, the rates  
16 at which the initial grip and load forces are applied during a lift will reflect how heavy and object  
17 looks rather than its actual mass. Thus an individual will initially lift a large object with a higher rate  
18 of force than they would use to lift an identically-weighted small object (Flanagan & Beltzner, 2000;  
19 Gordon, Forssberg, Johansson, & Westling, 1991), and they will initially lift a heavy-looking object  
20 with a higher rate of force than they would use to lift a light-looking object of the same mass (Baugh,  
21 Kao, Johansson, & Flanagan, 2012; Buckingham, Cant, & Goodale, 2009). In contrast to the  
22 persistence with which expectations influence a lifter's perception of heaviness, the gripping and  
23 lifting forces are rapidly modified with practice. Thus, only for the first few trials will individuals grip  
24 and lift the various illusion-inducing objects with different forces from one another, before adjusting  
25 their forces to reflect the stimuli's actual and identical masses (Buckingham et al., 2009; Flanagan &  
26 Beltzner, 2000). These effects are known as sensorimotor prediction and sensorimotor adaptation,  
27 respectively. This divergence between how illusion-inducing objects are perceived and acted upon  
28 has been taken to indicate that there are distinct neural representations maintained for conscious  
29 weight perception and predictive fingertip force control (Chouinard, Large, Chang, & Goodale, 2009;  
30 Flanagan, Bittner, & Johansson, 2008). However, there has been little support for this proposition  
31 from work with various neurological patient groups (Li, Randerath, Goldenberg, & Hermsdörfer,  
32 2011; Rabe et al., 2009).

33 The left cerebral hemisphere is an obvious candidate for the neural locus of predictive fingertip force  
34 control based on learned object properties such as size and material. The left hemisphere, in  
35 particular regions in the parietal cortex, has been associated with action planning and tool use in a  
36 number of studies (e.g., Valyear, Cavina-Pratesi, Stiglick, & Culham, 2007). Furthermore, damage to  
37 the left parietal cortex is thought to be the primary cause of deficits with tool use which are a  
38 particularly important manifestation of limb apraxia (Buxbaum, 2001; Goldenberg, 2014). Of  
39 particular relevance to the current work, it has been proposed that apraxia represents a specific  
40 impairment to the use of internal representations required for planning interactions with tools  
41 (Buxbaum, Johnson-Frey, & Bartlett-Williams, 2005). However, in spite of the clear conceptual link  
42 from this literature to the predictive way objects are lifted in weight-illusion paradigms, there has  
43 been no evidence for any such deficits in patients with left brain damage. The most comprehensive

1 study examined groups of patients with varying degrees of apraxia following unilateral damage to  
2 the left hemisphere and compared their fingertip forces and weight perception with non-apraxic  
3 patients with left and right brain lesions, as well as a control group (Li et al., 2011). The authors  
4 noted that individuals with apraxia showed a similar degree of size-based fingertip force scaling to  
5 the other groups, suggesting that the predictive application of fingertip force behaviour is neither  
6 lateralized to the motor-dominant left hemisphere (see also Buckingham, Ranger, & Goodale, 2012)  
7 nor related to the tool-used deficits seen in individuals with apraxia.

8 The findings outlined above would seem to indicate that the deficits in tool use/pantomime seen in  
9 apraxic patients are unrelated to sensorimotor prediction when lifting objects. A recent study by  
10 (Eidenmüller, Randerath, Goldenberg, Li, & Hermsdörfer, 2014) suggests that at least some forms of  
11 apraxia may disrupt the natural application of forces that individuals use to lift a range of familiar  
12 objects. In their study, individuals with left and right unilateral brain damage lifted a variety of  
13 everyday objects such as bottles of milk and packets of cigarettes while their gripping forces were  
14 measured with finger-mounted sensors. Typically, an unimpaired lifter's grip forces will show a  
15 precise anticipation of a familiar object's mass (Hermsdörfer, Li, Randerath, Goldenberg, &  
16 Eidenmüller, 2011). However, individuals with left brain damage (LBD) had clear impairments in their  
17 fingertip force scaling, showing far weaker associations between grip force rate and object heaviness  
18 than age-matched controls and patients with damage to their right hemisphere. Instead, the left  
19 brain damaged patients tended to grip the heaviest objects with approximately the same rate of  
20 force as the lightest objects. The deficit was most prominent in patients with visuo-motor apraxia as  
21 measured by the imitation of meaningless gestures. Similar findings have been reported in a group  
22 of six LBD patients when the vertical lifting forces were examined as a measure of weight  
23 anticipation during lifts of everyday objects (Dawson, Buxbaum, & Duff, 2010). Thus, in contrast to  
24 the research on how size affects anticipatory fingertip force control (Li et al., 2011), it appears that  
25 the fronto-parietal networks which have been implicated in successful praxis may be also critical for  
26 sensorimotor prediction based on object identity. These findings could be interpreted as an  
27 involvement of left hemisphere networks in independent processing of tool and object related  
28 information, which may in turn code the implicit knowledge of the forces required to lift a familiar  
29 object as well as one's understanding of how to use a particular tool.

30 The divergent findings of the object lifting studies in patients with LBD to date appears to indicate  
31 that object size influences fingertip force control in a way which is independent from how object  
32 identity influences how we grip and lift objects. Although there is no clear empirical evidence for this  
33 proposition, there are some findings consistent with this proposal in the object lifting literature. For  
34 one, the influence of size on fingertip forces appears to be particularly persistent, far more so than  
35 object identity or material properties. In most SWI studies, lifters will take several trials to adapt  
36 their fingertip forces to the objects' actual masses (e.g., Buckingham & Goodale, 2010; Chouinard et  
37 al., 2009; Flanagan & Beltzner, 2000; Grandy & Westwood, 2006), whereas studies where lifters who  
38 encounter unexpectedly-weighted objects without variations in size show extremely rapid, single-  
39 trial, adaptation (Buckingham et al., 2009; Gordon, Westling, Cole, & Johansson, 1993). If size cues  
40 are sufficiently salient and persistent to hinder the rapid adaptation to object mass, they may be  
41 resistant to the effects of LBD and apraxia which seem to hinder predictive grip force application  
42 with real-world objects.

1 To better determine if and how left brain damage influences the predictive application of fingertip  
2 forces, we examined object lifting behaviour in variety of contexts. Groups of patients with left and  
3 right unilateral brain injury lifted and judged the weight of (1) cubes with a normal size-weight  
4 relationship, (2) identically-weighted cubes with different volumes which typically induce the SWI,  
5 and (3) identically-weighted cubes of the same volume which varied in their apparent material  
6 properties to induce the MWI. In this latter case, when these objects are lifted, unimpaired lifters  
7 will lift the heavy-looking objects with more force than the light-looking objects, and judge the  
8 heavy-looking object as feeling lighter than the one which appear to be less heavy (Buckingham et  
9 al., 2009; Buckingham, Ranger, & Goodale, 2011). If the ability to use tools is related to the ability to  
10 the utilization of representations of object properties, lifters with LBD and apraxia are likely to show  
11 impaired predictive application of fingertip forces based on material properties, which should  
12 contrast their performance in the SWI condition.

13

#### 14 Materials and methods

15 In the current work, thirteen patients with left brain damage (LBD, 7 male, mean age = 56.7 years  $\pm$   
16 15.9), eleven with right brain damage (RBD, 6 male, mean age = 59.3 years  $\pm$  9.4), and fourteen  
17 neurologically-intact control subjects (CTR, 6 male, mean age = 57.3 years  $\pm$  8.4) were tested.  
18 Demographic and clinical information for the individuals in the LBD and RBD groups can be found in  
19 table 1. All participants were recruited from Munich, and testing was undertaken in the native  
20 language (German). All participants gave informed consent prior to testing, and all procedures were  
21 run in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki)  
22 and approved by the local ethics board at the Medical Faculty of the Technische Universität  
23 München.

24

---

25 Insert table 1 here

---

26

27 A variety of objects were lifted in three conditions – Natural, SWI and MWI. In the Natural condition,  
28 participants lifted naturally-weighted large and small wooden cubes. The large cube was  
29 10×10×10cm, and weighed 700g, whereas the small cube was 5×5×5cm and weighed 90g. In the SWI  
30 condition, participants lifted identically-weighted large and small metal cubes. In this condition, the  
31 large cube was 10×10×10cm, and weighed 700g, whereas the small cube was 5×5×5cm and also  
32 weighed 700g. In the MWI condition, participants lifted identically-weighted large cubes which  
33 appeared to be made from aluminium and expanded polystyrene. Both cubes in this condition were  
34 10×10×10cm and weighed 700g. In the SWI and MWI conditions, where adjustments were made to  
35 the natural weight of the stimuli, objects were hollowed out and filled with lead. Care was taken to  
36 ensure that all objects were centrally weighted. Further details about the construction of these  
37 stimuli can be found in Buckingham & Goodale (2013). Finally, a 540g grey plastic cylinder (diameter  
38 7.5cm, height 8cm) served as a neutral object for the washout trials. All objects were fitted with a  
39 mount in the centre of the top surface in order to facilitate the rapid attachment and removal of a  
40 140g handle. The contact surfaces of the handle were covered with sandpaper, allowing participants

1 to grip all the objects with a grip utilising the thumb, index finger, and middle finger. The custom-  
2 made handle contained three force sensors, two of which recorded the forces lateral (left and right  
3 grip forces, 0-80 N, accuracy  $\pm 0.1\text{N}$ ) to the handle's surface and one of which recorded the vertical  
4 load force (0-60N, accuracy  $\pm 0.1\text{N}$ ) each with a sampling frequency of 100 Hz (for details see Li et al.,  
5 2011).

6

---

7 Insert Figure 1 here

8

---

9 Participants were seated in front of a table and lifted objects in three sequential conditions,  
10 preceded and separated by washout trials with a neutral object. LBD and RBD patients lifted with the  
11 hand ipsilateral to their lesion, and all subjects in the CTR group lifted with their left hand. First,  
12 participants lifted the neutral object six times in a row. Then, participants lifted the large and small  
13 wooden cubes in alternation six times apiece (the Natural condition). Following another six lifts of  
14 the neutral objects, participants then lifted the large and small metal cubes in alternation six times  
15 apiece (the SWI condition). Finally, following another six lifts of the neutral object, participants lifted  
16 the aluminium and polystyrene cubes in alternation six times apiece (the MWI condition). The first  
17 object to be lifted in each condition was counterbalanced across participants. Prior to each  
18 condition, participants were asked to verbally indicate which object out of the pair they expected to  
19 be the heaviest. They were also asked to give a verbal indication of which object felt heaviest after  
20 the 3<sup>rd</sup> pair of lifts and after the final pair of lifts. Both of these post-lift ratings were averaged within  
21 subject to yield a proportional value representing how many perceptual reports on average stated  
22 they experienced the various weight illusions. In all cases, participants were permitted to report that  
23 the objects felt equally heavy if necessary. One patient in the LBD group was severely aphasic and,  
24 despite completing the lifting trials, was unable to give a verbal report of how heavy the objects felt.

25 The force transducers recorded the lateral grip forces and the vertical load forces for a total of 5.5  
26 seconds after participants were told to initiate their lift. The averaged grip force and the load force  
27 were differentiated to yield grip force rate (GFR) and load force rate (LFR). Non-parametric kernel  
28 estimates were used to smoothen the data as well as to calculate time derivatives with an effective  
29 cut-off frequency of 12 Hz (Marquardt & Mai, 1994). The difference between the peak GFR and peak  
30 LFR used to lift the objects in each pair (heavy-looking object – light-looking object) yielded the  
31  $\text{GFR}_{\text{diff}}$  and  $\text{LFR}_{\text{diff}}$  measures. The  $\text{GFR}_{\text{diff}}$  and  $\text{LFR}_{\text{diff}}$  on the first lift in each condition served as our  
32 measure of sensorimotor prediction based each object's appearance, and the  $\text{GFR}_{\text{diff}}$  and  $\text{LFR}_{\text{diff}}$   
33 averaged across the lifts 3-5 in each condition was taken to indicate the degree of fingertip force  
34 adaptation in each condition. Lifts 3-5 were chosen for this latter measure because, in multiple  
35 cases, the force data from final trial pair was lost due participant fatigue in the LBD and RBD groups.  
36 Other missing values (3.3% of the CTR data, 7.1% of the LBD data, and 7.2% of the RBD data) were  
37 filled with the series mean.

38

39

40

1

## 2 Results

### 3 Perceptual reports

4 There were no obvious differences between the groups in terms of their expectations of heaviness  
5 or their subsequent perceptions of heaviness in any of the conditions.

6 Prior to lifting the differently-sized, differently-weighted wooden cubes in the natural condition, 75%  
7 of those in the LBD group, 90.9% of those in the RBD group, and 100% of those in the CTR group  
8 showed intact expectations for size, stating that they predicted the large wood cube would be  
9 heavier than the small wood cube. The majority of participants reported accurate weight perception,  
10 with 79.2% of the LBD, 100% of the RBD and 100% of CTR reports indicating that they perceived the  
11 large wood cube to be heavier than its smaller and lighter counterpart.

12 Prior to lifting the differently-sized, identically-weighted cubes in the SWI condition, 83.3% of the  
13 LBD group, 70% of the RBD group and 100% of the control group again showed intact expectations  
14 for size, stating that they expected the large metal cube to be heavier than the small metal cube.  
15 After lifting, 83.3% of the LBD, 75.3% of the RBD, and 95.8% of the CTR perceptual reports given by  
16 the participants indicated that they experienced the SWI, stating that the small cube felt heavier  
17 than the large cube.

18 Prior to lifting the identically-sized and-weighted cubes with different material properties in the  
19 MWI condition, 83.3% of the LBD group, 100% of the RBD group, and 100% of the CTR group  
20 showed appropriate expectations for object material properties, stating that they expected the  
21 aluminium cube to be heavier than the polystyrene cube. After lifting, and in contrast to the SWI  
22 condition, relatively few individuals experienced the MWI, with only 37.5% of the LBD, 59.1% of the  
23 RBD, and 35.1% of the CTR perceptual reports stating that the polystyrene cube felt heavier than the  
24 aluminium cube.

25

### 26 Fingertip forces

27 The primary measures of sensorimotor prediction and adaptation reported in this study are the  
28  $GFR_{diff}$  and  $LFR_{diff}$  – the peak grip and load force rate used to lift the light-looking cube subtracted  
29 from the peak grip and load force rate used to lift the heavy-looking cube in each condition. This  
30 measure was examined on trial 1 and across the averages of trials 3-5. A positive  $GFR_{diff}$  and  $LFR_{diff}$   
31 which is significantly different from 0 during the initial trials in any of the conditions would indicate  
32 adequate sensorimotor prediction. By contrast, a  $GFR_{diff}$  and  $LFR_{diff}$  which significantly differed from 0  
33 on later trials would indicate good sensorimotor adaptation in the Natural condition (where the  
34 objects had different mass), but poor adaptation in the SWI and MWI conditions (where objects had  
35 the same mass).

36 First, the groups were examined with individual one-sample t-tests to determine if the  $GFR_{diff}$   
37 differed significantly from zero on the first trial, as an index of sensorimotor prediction (Figure 2A).  
38 As expected, on trial 1 the CTR group showed clear sensorimotor prediction in the Natural ( $t(13) =$



1 2.73,  $p < .05$ ), SWI ( $t(13) = 4.92$ ,  $p < .001$ ), and MWI conditions ( $t(13) = 2.63$ ,  $p < .05$ ). By contrast, the  
2 LBD group showed no evidence of sensorimotor prediction on trial 1 in the Natural ( $t(12) = 0.63$ ,  
3  $p = .54$ ) or SWI ( $t(12) = 1.75$ ,  $p = .11$ ) conditions. In the MWI condition, however, the LBD group did  
4 show significant sensorimotor prediction ( $t(12) = 2.22$ ,  $p < .05$ ). The RBD group also demonstrated  
5 intact sensorimotor prediction in the MWI condition ( $t(10) = 2.28$ ,  $p < .05$ ), and marginal evidence for  
6 sensorimotor prediction in the SWI condition ( $t(10) = 2.10$ ,  $p = .06$ ), but no evidence of sensorimotor  
7 prediction in the Natural condition ( $t(10) = 1.75$ ,  $p = .11$ ). A one-way ANOVA of the initial  
8 sensorimotor prediction scores revealed no significant difference between the groups in either the  
9 Natural ( $F(2,35) = 0.80$ ,  $p = .46$ ), MWI ( $F(2,35) = 0.08$ ,  $p = .99$ ) or SWI conditions ( $F(2,35) = 0.23$ ,  $p = .79$ ).

10 In terms of  $LFR_{diff}$  on the first lifts (Figure 2B), the CTR group showed evidence of sensorimotor  
11 prediction in the Natural ( $t(13) = 3.74$ ,  $p < .005$ ), SWI ( $t(13) = 9.19$ ,  $p < .001$ ), and MWI ( $t(13) = 2.56$ ,  
12  $p < .05$ ) conditions. In contrast to their  $GFR_{diff}$  scores, the LBD group also showed normal sensorimotor  
13 prediction in the Natural ( $t(7) = 6.59$ ,  $p < .001$ ), SWI ( $t(7) = 3.46$ ,  $p < .05$ ), and MWI ( $t(7) = 2.54$ ,  $p < .05$ )  
14 conditions. The RBD group's data followed a broadly similar pattern, showing normal sensorimotor  
15 prediction in the Natural ( $t(9) = 3.69$ ,  $p < .01$ ) and SWI ( $t(9) = 2.42$ ,  $p < .05$ ) conditions, but no  
16 sensorimotor prediction in the MWI condition ( $t(9) = 2.03$ ,  $p = .07$ ). A one-way ANOVA of the initial  
17 sensorimotor prediction scores revealed no significant difference between the groups in either the  
18 Natural ( $F(2,29) = 0.11$ ,  $p = .90$ ), MWI ( $F(2,29) = 0.39$ ,  $p = .68$ ) or SWI conditions ( $F(2,29) = 0.78$ ,  $p = .78$ ).

19 \_\_\_\_\_  
20 Insert figure 2 here

21 \_\_\_\_\_  
22

23 Next, each of the groups were examined with individual one-sample t-tests to determine if the  
24  $GFR_{diff}$  differed significantly from 0 on the later trials, as an index of sensorimotor adaptation (Figure  
25 3A). Although the  $GFR_{diff}$  in the CTR group did not differ from 0 in the SWI condition ( $t(13) = 1.44$ ,  
26  $p = .17$ ), and showed a robust difference in the Natural condition ( $t(13) = 5.14$ ,  $p < .001$ ), indicating  
27 adequate sensorimotor adaptation, they did not appear to adapt their forces in the MWI condition  
28 ( $t(13) = 2.70$ ,  $p < .05$ ). By contrast, the LBD group showed good evidence of sensorimotor adaptation,  
29 lifting on later trials with a  $GFR_{diff}$  which did not differ from zero on both the SWI ( $t(12) = 1.14$ ,  $p = .27$ )  
30 and MWI conditions ( $t(12) = 1.80$ ,  $p = .10$ ), and a  $GFR_{diff}$  which was significantly larger than zero in the  
31 Natural condition ( $t(12) = 3.64$ ,  $p < .005$ ). Finally, the RBD group showed sensorimotor adaptation in  
32 the Natural condition ( $t(10) = 3.87$ ,  $p < .005$ ) and the MWI condition ( $t(10) = 1.9$ ,  $p = .09$ ), but poor  
33 adaptation in the SWI condition ( $t(10) = 2.33$ ,  $p < .05$ ). A one way ANOVA for the sensorimotor  
34 adaptation scores revealed no differences between the groups in the SWI ( $F(2,35) = 0.58$ ,  $p = .56$ ),  
35 MWI ( $F(2,35) = 0.21$ ,  $p = .82$ ), or Natural ( $F(2,35) = 2.3$ ,  $p = .11$ ) conditions for the  $GFR_{diff}$  scores.

36 In terms of  $LFR_{diff}$  on the later trials (Figure 3B), the CTR group showed normal adaptation to the  
37 objects' different weights in the Natural condition ( $t(13) = 7.71$ ,  $p < .001$ ), and the objects' identical  
38 weights in the SWI condition ( $t(13) = 1.97$ ,  $p = .07$ ), but poor adaptation in the MWI condition ( $t(13) =$   
39  $2.59$ ,  $p < .05$ ). The LBD group showed good sensorimotor adaptation in the Natural ( $t(7) = 6.19$ ,  
40  $p < .001$ ) and MWI ( $t(7) = 1.29$ ,  $p = .24$ ) conditions, but poor adaptation in the SWI ( $t(7) = 2.97$ ,  $p < .05$ ),.

1 Finally, the RBD group showed good adaptation in the Natural ( $t(9) = 3.56, p < .01$ ) and MWI ( $t(9) =$   
2  $1.12, p = .29$ ) and SWI ( $t(9) = 1.88, p = .09$ ) conditions. A one way ANOVA for the sensorimotor  
3 adaptation scores revealed no differences between the groups in the SWI ( $F(2,29) = 1.22, p = .31$ ),  
4 MWI ( $F(2,29) = 0.21, p = .98$ ), or Natural ( $F(2,29) = 0.23, p = .80$ ) conditions for the  $LFR_{diff}$  scores.

---

5 Insert figure 3 here

---

6

## 7 Discussion

8 The goal of this study was to determine if damage to the motor-dominant left hemisphere affects  
9 fingertip force control and weight perception. To this end, we examined how individuals with  
10 unilateral brain injury interacted with object pairs which varied in either their mass, volume or  
11 apparent material properties. In each of these three conditions, participants lifted pairs of objects,  
12 one of which appeared to be heavier than the other. Prior to lifting, participants reported which of  
13 the objects they expected to be heavier, and twice during each condition they reported which object  
14 felt heavier. The rate at which grip and load forces were applied to each object was measured on  
15 every trial, and the difference between the forces used to lift the heavy-looking and light-looking  
16 object in each pair served as a metric of different aspects of sensorimotor performance.

17 In the first condition (Natural), participants lifted large and a small wooden cubes with the same  
18 density and, as a consequence, different weights from one another. The majority of participants in  
19 all groups expected the large cube to weigh more than the small cube, and (unsurprisingly)  
20 subsequently judged the large cube as feeling heavier than the small cube. This condition, while of  
21 little theoretical interest to the experimental question, does serve to show that participants were,  
22 on the whole, able to understand the concept of heaviness and accurately report real weight  
23 differences. However, in terms of the grip forces applied to the objects, there were indications that  
24 the LBD and RBD groups were impaired in their initial sensorimotor prediction, failing to lift the  
25 heavy-looking cube with more force than the light-looking cube. This contrasted with the behaviour  
26 of the control subjects, who applied more force to the large cube than the small cube, as indicated  
27 by  $GFR_{diff}$  which was significantly greater than zero. Interestingly, neither the LBD nor RBD groups  
28 showed any difficulties scaling their load force rates to object size, suggesting that any deficit in  
29 force scaling may be restricted to the gripping forces. In both grip and load forces, all groups in this  
30 condition showed similar patterns of sensorimotor adaptation, learning to apply grip force at a  
31 greater rate to the heavy cube than to the light cube.

32 In the SWI condition, participants lifted large and small metal cubes which had the same mass as one  
33 another. As in the first condition, the majority of participants in all groups expected the large cube to  
34 outweigh the small cube, and in the majority of cases reported experiencing a SWI (i.e., that the  
35 small cube felt heavier than the large cube). It is worth noting that the clear presence of the size  
36 weight illusion in the LBD group compliments earlier work showing an intact SWI in individuals with  
37 left brain damage and apraxia (Li et al., 2011). Similarly, and also in agreement with Li and colleagues  
38 (2011), the LBD and RBD groups showed accurate sensorimotor prediction of load forces based on  
39 object size, indicating that unilateral brain damage does not impair the use of size as a cue to weight.  
40 By contrast, there was some evidence that unilateral brain injury selectively impairs the scaling of  
41 grip forces to size cues, as neither the LBD nor RBD group showed significant sensorimotor

1 prediction in the SWI condition. However, it is worth highlighting that none of the groups differed  
2 from one another when directly compared on either measure. In terms of fingertip force adaptation,  
3 the LBD group showed difficulties adapting their load forces, whereas the RBD group showed  
4 difficulties adapting their grip forces.

5 In the MWI condition, participants lifted identically-sized and –weighted cubes which appeared to be  
6 made from metal and polystyrene. The majority of participants in all groups expected the metal  
7 cube to outweigh the polystyrene cube. However, despite the fact that these cubes have been  
8 shown to induce a MWI in normal populations (Buckingham et al., 2009, 2011; Buckingham &  
9 Goodale, 2013), there was little evidence of such an effect here. Only the RBD group reported  
10 experiencing the MWI a substantial proportion of the time, whereas other groups tended to report  
11 either that the objects felt the same weight as one another or, less frequently, that the metal cube  
12 felt heavier than the polystyrene cube. In terms of lifting behaviour, however, all of the groups  
13 showed robust sensorimotor prediction of grip forces to objects' material properties, and only the  
14 RBD group showed an impairment in prediction with their loading forces. In this case, it is unclear  
15 whether this lack of prediction in the RBD group stems from a particular deficit in utilization of  
16 object properties, or merely from the high variability seen in this measure.

17 The main aim of this study was to examine how visual material cues influenced the application of  
18 fingertip forces when lifting objects in patients with damage to the left hemisphere. Previous work  
19 has demonstrated that, while these patients have no deficits in using size cues to scale their gripping  
20 (Li et al., 2011), they are impaired at using object identity to guide the application of fingertip forces  
21 (Eidenmüller et al., 2014). Conceptually, the apparent material of an object appears to bridge the  
22 gap between these two paradigms, and examining object lifting performance in this patient group  
23 would have enhanced our understanding of the extent and root cause of the difficulties that  
24 individuals with apraxia experience with tool use. If patients in the LBD group showed a familiar  
25 object-like impairment in their sensorimotor anticipation of material properties, it would suggest  
26 that size cues to object weight are a unique and special property which is largely conserved in the  
27 sensorimotor system following this type of brain injury. If, however, patients with LBD were not  
28 impaired in using materials properties to guide their initial forces, it would suggest that object  
29 identity is processed separately from more general size and material cues to weight. Our data  
30 appear to support this latter view, as both of our patient groups lifted the metal cube with a higher  
31 rate of force than the polystyrene cube on the first trial. Thus, our findings provide tentative  
32 evidence that unilateral brain injury does not affect how visual material cues are used to guide  
33 gripping and loading forces, suggesting that object identity is either less robust to brain injury than  
34 material cues (perhaps due to differences in how difficult it is to identify an individual object as  
35 opposed to identifying a block of homogenous material), or processed by alternate regions of the  
36 sensorimotor system. Further work examining how sensorimotor memory for object identity and  
37 material properties interact with one another in patients with unilateral brain damage is necessary  
38 to disentangle these possibilities.

39 Beyond the main aim of the study, two findings from the current work were sufficiently unexpected  
40 to require further discussion. The first, the lack of a robust MWI effect, may stem from the method  
41 of reporting employed in the current work, as it is likely that our simple method of perceptual  
42 reports is substantially less sensitive than the numerical ratings used in previous works (Buckingham  
43 et al., 2009, 2011; Buckingham & Goodale, 2013). Furthermore, the lack of a MWI, especially in our

1 control group, may be a consequence of the age of the sample in our current work. The MWI studies  
2 using these stimuli listed above have focussed on university-aged participants. To our knowledge, no  
3 body of work has investigated how weight illusions alter over the course of one’s lifespan, and our  
4 findings provide some indications that this might be an interesting topic for further study.

5 The second point worth examining from the current body of work is the lack of sensorimotor  
6 prediction for grip force rates based on object size. The previous work on this topic (Li et al., 2011)  
7 suggested that it was likely that both the LBD and RBD groups would show intact sensorimotor  
8 prediction. Indeed, both groups lifted the large cube with a higher rate of load force than the small  
9 cube. However, this pattern failed to emerge with the grip force rate – both groups lifted the  
10 differently-sized objects with similar rates of grip force on the first trial. Similar patterns of data  
11 were seen in the natural condition – the LBD and RBD groups lifted the large and small cubes with  
12 similar grip forces, but disparate load forces. It seems unlikely that this lack of grip force scaling to  
13 object size cues is a consequence of a default grip force, as this would not explain why these  
14 individual scaled their grip forces to object material properties just as well as the control subjects  
15 did. Instead, it seems that unilateral brain damage may lead to a selective impairment in the  
16 parameterization of grip forces for object size cues. Li et al. (2011) report similar conclusions in a  
17 sub-sample of their LBD individuals, suggesting that those with the largest occipito-parietal lesions  
18 showed the greatest impairments of grip force scaling. This finding is also consistent with an earlier  
19 single case study reporting a patient with a left parieto-occipital lesion who also showed no scaling  
20 of grip forces according to size (Li, Randerath, Goldenberg, & Hermsdörfer, 2007). Interestingly, the  
21 opposite pattern was found in a study of lifting SWI-inducing objects in patients with cerebellar  
22 diseases, who showed normal anticipation of size for the grip forces, but reduced anticipation for  
23 the load forces (Rabe et al., 2009).

24 To sum up, the current work aimed to investigate whether individuals with unilateral brain damage  
25 were impaired in using apparent material properties to anticipate the mass of objects when lifting  
26 them. We found no evidence that unilateral brain damage affected the anticipatory control of  
27 objects based upon their material properties. We did, however, find deficits in how individual with  
28 brain injury scale their grip forces, but not load forces, to size cues Future work will build on these  
29 findings to examine how different types of brain injury affects the integration of long-term prior  
30 expectations of object weight derived from various visual cues with short-term feedback from recent  
31 object interactions.

32

### 33 Acknowledgements

34 This project was funded with a grant to support the initiation of international collaboration from the  
35 Deutsche Forschungsgemeinschaft (DFG). We thank Prof. G. Goldenberg, head of the Clinic for  
36 Neuropsychology at Hospital München-Bogenhausen for his support providing us with access to the  
37 patients tested in this study.

38

1 References

- 2 Baugh, L. A., Kao, M., Johansson, R. S., & Flanagan, J. R. (2012). Material Evidence- Interaction of  
3 Well-Learned Priors and Sensorimotor Memory When Lifting Objects. *Journal of*  
4 *Neurophysiology*. doi:10.1152/jn.00263.2012
- 5 Buckingham, G. (2014). Getting a grip on heaviness perception: a review of weight illusions and their  
6 probable causes. *Experimental Brain Research*, 232(6), 1623–1629. doi:10.1007/s00221-014-  
7 3926-9
- 8 Buckingham, G., Cant, J. S., & Goodale, M. A. (2009). Living in A Material World: How Visual Cues to  
9 Material Properties Affect the Way That We Lift Objects and Perceive Their Weight. *Journal*  
10 *of Neurophysiology*, 102(6), 3111–3118. doi:10.1152/jn.00515.2009
- 11 Buckingham, G., & Goodale, M. A. (2010). The influence of competing perceptual and motor priors in  
12 the context of the size–weight illusion. *Experimental Brain Research*, 205(2), 283–288.  
13 doi:10.1007/s00221-010-2353-9
- 14 Buckingham, G., & Goodale, M. A. (2013). Size Matters: A Single Representation Underlies Our  
15 Perceptions of Heaviness in the Size-Weight Illusion. *PLoS ONE*, 8(1), e54709.  
16 doi:10.1371/journal.pone.0054709
- 17 Buckingham, G., Ranger, N. S., & Goodale, M. A. (2011). The material–weight illusion induced by  
18 expectations alone. *Attention, Perception, & Psychophysics*, 73(1), 36–41.  
19 doi:10.3758/s13414-010-0007-4
- 20 Buckingham, G., Ranger, N. S., & Goodale, M. A. (2012). Handedness, laterality and the size-weight  
21 illusion. *Cortex*.
- 22 Buxbaum, L. J. (2001). Ideomotor apraxia: a call to action. *Neurocase*, 7(6), 445–458.  
23 doi:10.1093/neucas/7.6.445
- 24 Buxbaum, L. J., Johnson-Frey, S. H., & Bartlett-Williams, M. (2005). Deficient internal models for  
25 planning hand-object interactions in apraxia. *Neuropsychologia*, 43(6), 917–929.  
26 doi:10.1016/j.neuropsychologia.2004.09.006

- 1 Charpentier, A. (1891). Analyse expérimentale quelques éléments de la sensation de opoids.  
2 *Archives de Physiologie Normales et Pathologiques*, 3, 122–135.
- 3 Chouinard, P. A., Large, M., Chang, E., & Goodale, M. (2009). Dissociable neural mechanisms for  
4 determining the perceived heaviness of objects and the predicted weight of objects during  
5 lifting: An fMRI investigation of the size–weight illusion. *NeuroImage*, 44(1), 200–212.  
6 doi:10.1016/j.neuroimage.2008.08.023
- 7 Dawson, A. M., Buxbaum, L. J., & Duff, S. V. (2010). The impact of left hemisphere stroke on force  
8 control with familiar and novel objects: neuroanatomic substrates and relationship to  
9 apraxia. *Brain Research*, 1317, 124–136. doi:10.1016/j.brainres.2009.11.034
- 10 Eidenmüller, S., Randerath, J., Goldenberg, G., Li, Y., & Hermsdörfer, J. (2014). The impact of  
11 unilateral brain damage on anticipatory grip force scaling when lifting everyday objects.  
12 *Neuropsychologia*, 61, 222–234. doi:10.1016/j.neuropsychologia.2014.06.026
- 13 Ellis, R. R., & Lederman, S. J. (1999). The material-weight illusion revisited. *Perception &*  
14 *Psychophysics*, 61(8), 1564–1576.
- 15 Flanagan, J. R., & Beltzner, M. A. (2000). Independence of perceptual and sensorimotor predictions  
16 in the size-weight illusion. *Nature Neuroscience*, 3(7), 737–741. doi:10.1038/76701
- 17 Flanagan, J. R., Bittner, J. P., & Johansson, R. S. (2008). Experience can change distinct size-weight  
18 priors engaged in lifting objects and judging their weights. *Current Biology*, 18(22), 1742–  
19 1747. doi:10.1016/j.cub.2008.09.042
- 20 Goldenberg, G. (2014). Apraxia - the cognitive side of motor control. *Cortex; a Journal Devoted to the*  
21 *Study of the Nervous System and Behavior*, 57, 270–274. doi:10.1016/j.cortex.2013.07.016
- 22 Goldenberg, G., & Hagmann, S. (1997). The meaning of meaningless gestures: a study of visuo-  
23 imitative apraxia. *Neuropsychologia*, 35(3), 333–341.
- 24 Gordon, A. M., Forssberg, H., Johansson, R. S., & Westling, G. (1991). Visual size cues in the  
25 programming of manipulative forces during precision grip. *Experimental Brain Research*,  
26 83(3), 477–482.

- 1 Gordon, A. M., Westling, G., Cole, K. J., & Johansson, R. S. (1993). Memory Representations  
2 Underlying Motor Commands Used During Manipulation of Common and Novel Objects.  
3 *Journal of Neurophysiology*, 69(6), 1789–1796.
- 4 Grandy, M. S., & Westwood, D. A. (2006). Opposite Perceptual and Sensorimotor Responses to a  
5 Size-Weight Illusion. *Journal of Neurophysiology*, 95(6), 3887–3892.  
6 doi:10.1152/jn.00851.2005
- 7 Hermsdörfer, J., Li, Y., Randerath, J., Goldenberg, G., & Eidenmüller, S. (2011). Anticipatory scaling of  
8 grip forces when lifting objects of everyday life. *Experimental Brain Research*, 212(1), 19–31.  
9 doi:10.1007/s00221-011-2695-y
- 10 Huber, W., Poeck, K., & Willmes, K. (1984). The Aachen Aphasia Test. *Advances in Neurology*, 42,  
11 291–303.
- 12 Li, Y., Randerath, J., Goldenberg, G., & Hermsdörfer, J. (2007). Grip forces isolated from knowledge  
13 about object properties following a left parietal lesion. *Neuroscience Letters*, 426(3), 187–  
14 191. doi:10.1016/j.neulet.2007.09.008
- 15 Li, Y., Randerath, J., Goldenberg, G., & Hermsdörfer, J. (2011). Size–weight illusion and anticipatory  
16 grip force scaling following unilateral cortical brain lesion. *Neuropsychologia*, 49(5), 914–  
17 923. doi:10.1016/j.neuropsychologia.2011.02.018
- 18 Marquardt, C., & Mai, N. (1994). A computational procedure for movement analysis in handwriting.  
19 *Journal of Neuroscience Methods*, 52(1), 39–45.
- 20 Murray, D. J., Ellis, R. R., Bandomir, C. A., & Ross, H. E. (1999). Charpentier (1891) on the size-weight  
21 illusion. *Perception & Psychophysics*, 61(8), 1681–1685.
- 22 Rabe, K., Brandauer, B., Li, Y., Gizewski, E. R., Timmann, D., & Hermsdörfer, J. (2009). Size–Weight  
23 Illusion, Anticipation, and Adaptation of Fingertip Forces in Patients With Cerebellar  
24 Degeneration. *Journal of Neurophysiology*, 101(2), 569–579. doi:10.1152/jn.91068.2008
- 25 Seashore, C. E. (1899). Some psychological statistics II. The material weight illusion. *University of*  
26 *Iowa Studies in Psychology*, (2), 36–46.

1 Valyear, K. F., Cavina-Pratesi, C., Stiglick, A. J., & Culham, J. C. (2007). Does tool-related fMRI activity  
2 within the intraparietal sulcus reflect the plan to grasp? *NeuroImage*, *36 Suppl 2*, T94–T108.  
3 doi:10.1016/j.neuroimage.2007.03.031  
4  
5



1 Figure captions

2 **Table 1.** Demographic for all groups, and available clinical information for the LBD and RBD groups.  
3 Aphasia tested with the Aachen Aphasia test (Huber, Poeck, & Willmes, 1984). Hand and finger  
4 imitation tests developed by G. Goldenberg (G Goldenberg & Hagmann, 1997).

5 **Figure 1.** The stimuli used across the conditions (left panels), and the handle containing the force  
6 transducers attached to the large aluminium cube (right panel).

7 **Figure 2.** The (A)  $GFR_{diff}$  and (B)  $LFR_{diff}$  used for the first lift the object pairs in each condition This  
8 measure is an index of effective sensorimotor prediction based on the visual properties of the  
9 objects, with significant differences suggesting good sensorimotor prediction. Error bars indicate  
10 standard error of the means. \* indicates a significant difference from zero at  $p \leq .05$ .

11 **Figure 3.** The (A)  $GFR_{diff}$  and (B)  $LFR_{diff}$  used in the later lifts of the object pairs in each condition. This  
12 measure provides an indication of the efficiency of sensorimotor adaptation, with significant  
13 differences in the natural condition indication good adaptation, but significant differences in the  
14 illusion conditions indicating poor adaptation. Error bars indicate standard error of the means. \*  
15 indicates a significant difference from zero at  $p \leq .05$ .

16