



Heriot-Watt University
Research Gateway

Flows and stochastic Taylor series in Ito calculus

Citation for published version:

Wiese, A, Malham, SJ, Ebrahimi-Fard, K & Patras, F 2015, 'Flows and stochastic Taylor series in Ito calculus', *Journal of Physics A: Mathematical and Theoretical*, vol. 48, no. 49, 495202.
<https://doi.org/10.1088/1751-8113/48/49/495202>

Digital Object Identifier (DOI):

[10.1088/1751-8113/48/49/495202](https://doi.org/10.1088/1751-8113/48/49/495202)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Peer reviewed version

Published In:

Journal of Physics A: Mathematical and Theoretical

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

FLOWS AND STOCHASTIC TAYLOR SERIES IN ITÔ CALCULUS

KURUSCH EBRAHIMI-FARD^a, SIMON J.A. MALHAM^b,
FRÉDÉRIC PATRAS^c AND ANKE WIESE^b

ABSTRACT. For general stochastic systems driven by continuous semimartingales an explicit formula for the logarithm of the Itô flow map is given. The computation relies on the lift to quasi-shuffle algebras of formulas involving products of Itô integrals of semimartingales. Whereas the Chen–Strichartz formula computing the logarithm of the Stratonovich flow map is classically expanded as a formal sum indexed by permutations, the analogous formula in Itô calculus is naturally indexed by surjections. This reflects the change of algebraic background involved in the transition between the two integration theories. Lastly we extend our formula for the quasi-shuffle Chen–Strichartz series for the logarithm of the flow map to the non-commutative case. For linear matrix-valued SDEs driven by arbitrary semimartingales we obtain a similar formula.

1. INTRODUCTION

The setting of our work is classical stochastic calculus, as exposed, e.g., in the textbooks [24, 32]. Its aim is to obtain an explicit formula for the logarithm of the Itô flow map associated to a (general, nonlinear) stochastic differential system (Theorem 2.1), generalizing the classical work of Ben Arous [3] on Stratonovich flows and stochastic Taylor series.

As observed by Ben Arous, his results show “a *miraculous* interaction between purely algebraic formulas on the Campbell–Hausdorff series and probabilistic identities between Itô and Stratonovich formulas” [3, p. 31]. In the same passage he also raises the question of whether these phenomena are intrinsically probabilistic or whether they result from more formal and algebraic phenomena. Our work aims in the same direction by addressing the general interplay between stochastic integration on the one hand and purely algebraic structures pertaining to the modern combinatorial theory of surjections on the other hand. It shows that the same phenomena hold, when Stratonovich integrals are replaced by Itô ones, except for the fact that a more involved Campbell–Hausdorff-type series has to be introduced to take into account extra terms arising from Itô’s formula (Theorem 7.2).

In view to applications we develop the theory directly for differential systems driven by arbitrary continuous semimartingales, and thereby go beyond the case of systems driven by Brownian motion. This is motivated by the very definition of semimartingales, which is devised precisely so as to extend the domain of applications of classical stochastic integration methods.

We recall that examples for continuous martingales include integrable stochastic integrals over Brownian motion and continuous time-changed Brownian motion. The Martingale Representation Theorem for Brownian Motion shows in turn that a continuous martingale that is adapted to the filtration generated by a Brownian motion can be represented as a stochastic integral with respect to this Brownian motion, see Theorem IV.42 in Protter [32]. Further, the representation results of Dambis, of Dubins and Schwarz, and of Knight show that a continuous martingale with a quadratic variation process that is increasing to infinity is a time-changed Brownian motion, see Theorems 3.4.6 and 3.4.13 in Karatzas & Shreve [25]. Stochastic integration and time-change of Brownian motion thus enable the construction of a rich class of new processes. Examples for semimartingales that admit

jumps include Lévy processes; some of these can be constructed as time-changed Brownian motion. Lévy processes have become increasingly important and have proven to be useful as models in financial and physical applications in recent years, see the monograph by Barndorff–Nielsen, Mikosch and Resnick [1].

Having computed the logarithm of the flow map, the next step is to show that it is indeed a Lie series. This fact (which is not immediate, but can be proven to hold true due, among others, to general properties of quasi-shuffle Hopf algebras) will be the object of further studies. Let us mention here that the existence of such a Lie series is important for the construction of Lie group integration schemes for stochastic differential systems whose solutions evolve on a homogeneous manifold; see Malham and Wiese [28]. In principle, for example using a method analogous to that developed by Castell and Gaines [5], one could use our results to construct strong approximation schemes for which the approximate solution lies on the homogeneous manifold (to a high accuracy). This would be for general systems, for example driven by continuous semimartingales. In [28], Malham and Wiese considered for example the dynamics of a rigid body such as a satellite perturbed by two independent multiplicative Wiener processes, as well as the dynamics of an ellipsoidal autonomous underwater vehicle perturbed in the same way.

The techniques developed in the article to study Itô-type formulas are of interest on their own. In fact, the present results also connect to rough paths theory (as well as to Hairer’s theory of regularity structure) – by going in the same direction, namely, by formalising integral stochastic calculus with the help of Hopf algebraic tools. Developments of a series of similar methods, accompanied by Hopf algebra techniques, to handle, e.g., stochastic differential equations driven by Brownian motions and continuous semimartingales; rough differential equations driven by geometric and non-geometric rough paths [20]; or the more general equations driven by highly irregular paths of the theory of regularity structures [21], suggest indeed the emergence of a common group theoretical framework beyond the “classical” integral calculus governed by Chen’s calculus of iterated integrals and shuffle algebras.

This remark applies in particular when comparing the present article with reference [20], which develops an algebraic approach to non-geometric rough paths and their links with geometric ones that can be thought of as a generalisation of the Itô–Stratonovich correction formula. However, whereas in [20] these links are understood by means of the decorated Butcher–Connes–Kreimer Hopf algebra of rooted trees, our approach follows a different path by deducing the principles governing Itô’s calculus from the (Hopf algebraic) structure underlying quasi-shuffle algebras.

Acknowledgements: The first author is supported by a Ramón y Cajal research grant from the Spanish government. Support from the Spanish government under the project MTM2013-46553-C3-2-P is acknowledged. The third author acknowledges support from the grant ANR-12-BS01-0017, Combinatoire Algébrique, Résurgence, Moules et Applications.

2. THE LOGARITHM OF THE ITÔ FLOW MAP

In this section, we state the main result of this work. A description of the historical background and details on the underlying definitions and objects of study are postponed to the next section.

Let $\{X^1, X^2, \dots, X^N\}$ be scalar continuous semimartingales. We assume, without loss of generality, that $X_0^i = 0$ and that their quadratic covariation, or square bracket operation, is such that $[X^i, X^j] \equiv 0$ for $i \neq j$. We consider the general stochastic differential system (in what follows the notation $\int_0^t \dots dX_s^i$ refers to Itô integrals)

$$(1) \quad Y_t = Y_0 + \sum_{i=1}^N \int_0^t V_i(Y_s) dX_s^i,$$

where $V_i : \mathbb{R}^d \rightarrow \mathbb{R}^d$ are smooth vector fields. In the following, we will identify V_i with the partial differential operator $V_i \partial_y := \sum_{j=1}^d V_i^j \partial_{y_j}$. For $i = N + 1, \dots, 2N$, let X^i denote the quadratic variation of X^{i-N} , that is $X^i = [X^{i-N}, X^{i-N}]$, and define the second order differential operator V_i by

$$(2) \quad V_i := \frac{1}{2} \sum_{j, k=1}^d V_i^j V_i^k \partial_{y_j y_k}.$$

By analogy to the Stratonovich stochastic system driven by Wiener processes in [2], we define the stochastic partial differential operator S as

$$S := \sum_{i=1}^{2N} V_i X^i.$$

Since the differential operators V_i are time-homogeneous, we also have $dS_t := \sum_{i=1}^{2N} V_i dX_t^i$. For $n \geq 1$, let $\int S^n$ denote the n -times repeated integral of S and set $\int S^0 = \text{Id}$, so that

$$\int S^n = \int \cdots \int dS_{t_1} \cdots dS_{t_n} = \sum_{j_1, j_2, \dots, j_n=1}^{2N} V_{j_1} \circ \cdots \circ V_{j_n} \int \cdots \int dX_{t_1}^{j_1} \cdots dX_{t_n}^{j_n}.$$

The Itô-Taylor series expansion for the flow map $\varphi_t : Y_0 \mapsto Y_t$ corresponding to the stochastic system is given by

$$\varphi_t = \sum_{n \geq 0} \int S^n \circ \text{Id}.$$

We set $\mathbb{S} := \sum_{n \geq 0} \int S^n = \text{Id} + \int \mathbb{S} dS$, which describes the action of the flow map on smooth functions. The central aim of this work is to calculate $\log \mathbb{S}$ in the Itô framework.

The computation of the logarithm of this action may be considered as the stochastic analog of a well-known problem in the theory of classical differential equations (motivated, e.g., by numerical considerations, and referred often to as the continuous Baker–Campbell–Hausdorff problem). And it turns out that, provided one uses Stratonovich integrals, the solution for stochastic differential equations is essentially the same as in the classical case. We refer the reader to the next section for more details.

Displaying now the first few terms of $\log \mathbb{S}$ in the Itô framework may give an idea of the complexity of its expression. We adopt the notation $I_{[i,j]} = [X^i, X^j]$, and for a repeated Itô integral $I_{j_1, \dots, j_n} := \int \cdots \int dX^{j_1} \cdots dX^{j_n}$. Then the first three terms are as follows

$$(3) \quad \begin{aligned} \log \mathbb{S} = & \sum_i V_i I_i + \sum_{i,j} V_i V_j \left(\frac{1}{2} I_{ij} - \frac{1}{2} (I_{ji} + I_{[i,j]}) \right) \\ & + \sum_{i,j,k} V_i V_j V_k \left(\frac{1}{3} I_{ijk} - \frac{1}{6} (I_{jik} + I_{kij} + I_{[i,j]k} + I_{j[i,k]}) \right. \\ & \left. - \frac{1}{6} (I_{ikj} + I_{jki} + I_{i[j,k]}) + \frac{1}{3} (I_{kji} + I_{[j,k]i} + I_{k[i,j]}) \right) + \cdots \end{aligned}$$

To write the general expression for this expansion, let us introduce some notation. For f a surjection from the set $[n] := \{1, \dots, n\}$ to the set $[k]$ (written $f \in \text{Sj}_{n,k}$), we set

$$d(f) := |\{i < n, f(i) \geq f(i+1)\}|.$$

The set of surjections f from $[n]$ to $[k]$ such that $\forall i \leq k, |f^{-1}\{i\}| \leq 2$ is written $\text{Sj}_{n,k}^{(2)}$. For a sequence $J = (j_1, \dots, j_n)$ of elements of $[2N]$ and $\mathcal{A} = A_1 \amalg \cdots \amalg A_k = [n]$ an ordered partition of the set $[n]$ into disjoint subsets, we write $I_{\mathcal{A}}^J$ for the iterated integral $\int \cdots \int dX_J^{A_1} \cdots dX_J^{A_k}$, where, for $A_i = \{a_1, \dots, a_l\}$, $X_J^{A_i}$ stands

for the iterated quadratic covariation $[X^{j_{a_1}}, \dots, X^{j_{a_l}}] := [X^{j_{a_1}}, [X^{j_{a_2}}, \dots, X^{j_{a_l}}]]$. For f as above, we set $\mathcal{A}(f) := f^{-1}(1) \coprod \dots \coprod f^{-1}(k)$ and

$$S_f := \sum_{i_1, \dots, i_n=1}^{2N} V_{i_1} \cdots V_{i_n} I_{\mathcal{A}(f)}^{(i_1, \dots, i_n)}.$$

Our main result reads

Theorem 2.1. *We have:*

$$\log(\mathbb{S}) = \sum_{n>0} \sum_{n \geq k \geq 1} \sum_{f \in S_{j_{n,k}}^{(2)}} \frac{(-1)^{d(f)}}{n} \cdot \binom{n-1}{d(f)}^{-1} S_f.$$

This statement follows from Theorem 7.2. The restriction of the indexing set to $S_{j_{n,k}}^{(2)}$ follows from the fact that, for continuous semimartingales, iterated brackets such as $[[X_i, X_j], X_k]$ vanish. When allowing semimartingales with jumps (the article will develop the combinatorial theory of iterated integrals of semimartingales in this more general setting), this restriction disappears.

Example 2.2. With f the surjection from the set [3] to the set [2], defined by

$$f(1) = 2, \quad f(2) = 1, \quad f(3) = 2,$$

we obtain $d(f) = 1$, and

$$\frac{(-1)^{d(f)}}{n} \cdot \binom{n-1}{d(f)}^{-1} = \frac{(-1)}{3} \cdot \binom{2}{1}^{-1} = -\frac{1}{6}$$

and $S_f = \sum_{i,j,k} V_i V_j V_k I_{j[ik]}$, as expected from the low order direct computation given previously in (3).

The rest of the paper develops the formalism necessary to prove the above theorem. Several of the tools that enter our approach are of general interest, and allow to handle the algebraic structures of iterated Itô integrals. As a supplement, following the same ideas, a formula similar to the one for the logarithm of the Itô flow map is obtained for solutions of linear matrix-valued SDEs driven by arbitrary semimartingales (Theorem 8.5).

3. THE STRICHARTZ FORMULA

Let us recall first the historical as well as technical background of Theorem 2.1. Its knowledge will help to enlighten our forthcoming constructions, and make clear to what extent Itô calculus differs from the Riemann or Stratonovich calculus.

From the seminal 1957 work by K.T. Chen on the algebraic structures underlying products of iterated integrals [8] followed the existence of an exponential solution of the classical nonautonomous initial value problem

$$\dot{Y}(t) = F(t, Y(t)), \quad Y(0) = Y_0,$$

where $F(t) : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a vector field depending continuously on time.

Over the decades following Chen's work, the explicit formula for this exponential solution was obtained independently by several authors. Mielnik and Plebanski [29] as well as Strichartz [35] calculated the function $\Omega(t)$ such that $Y(t) = \exp(\Omega(t))Y_0$. Using the fact that Stratonovich integrals obey Chen's rules of calculus for iterated integrals, Ben Arous showed soon after the work of Strichartz, that the application domain of this formula extends to stochastic realm [2, 3].

The explicit expression of the series $\Omega(t)$ is rather intricate. Its most classical formulation involves permutations and iterated integrals of iterated Lie brackets

$$\Omega(t) = \sum_{n>0} \sum_{\sigma \in S_n} \frac{(-1)^{d(\sigma)}}{n^2 \binom{n-1}{d(\sigma)}} \int_{\Delta_{[0,t]}^n} [\cdots [F(s_{\sigma(1)}), F(s_{\sigma(2)}), \cdots], F(s_{\sigma(n)})] ds_1 \cdots ds_n,$$

where the bracket of vector fields follows from their interpretation as differential operators, and the integration domain is the n -dimensional simplex

$$\Delta_{[0,t]}^n := \{(s_1, \dots, s_n), 0 \leq s_1 \leq \cdots \leq s_n \leq t\}.$$

In the formula for $\Omega(t)$, S_n denotes the set of permutations of the set $[n] := \{1, \dots, n\}$. The quantity $d(\sigma)$ (already introduced in the more general case of surjections) is called the number of descents of the permutation $\sigma \in S_n$, that is the number of positions in the permutation $(\sigma(1), \dots, \sigma(n))$, where $\sigma(i) > \sigma(i+1)$, for $i = 1, \dots, n-1$. The formula is known in the literature as Strichartz, Chen–Strichartz or continuous Baker–Campbell–Hausdorff formula.

It can also be stated by replacing iterated Lie brackets by standard operator products. The formula is then essentially the same except for the scalar coefficient, which becomes the one displayed in Theorem 2.1 with surjections replaced by permutations, i.e., $(-1)^{d(\sigma)} n^{-1} \binom{n-1}{d(\sigma)}^{-1}$. We refer the reader to [35] for details on the analytic background, and to [33, 19] for the underlying combinatorics of free Lie algebras and Lie idempotents.

The derivation of the formula for $\Omega(t)$ relies on a precise understanding of the calculus of iterated integrals, which in turn is based on integration by parts. For two scalar valued indefinite integrals $F(t) := \int_0^t f(s) ds$ and $G(t) := \int_0^t g(s) ds$, recall that

$$F(t) \cdot G(t) = \int_0^t f(s)G(s) ds + \int_0^t F(s)g(s) ds.$$

For general iterated integrals

$$F_n(t) := \int_{\Delta_{[0,t]}^n} f_1(s_1) f_2(s_2) \cdots f_n(s_n) ds_1 \cdots ds_n,$$

the above product generalizes to Chen’s shuffle product formula

$$(4) \quad F_n(t) \cdot F_m(t) = \sum_{\sigma \in \text{Sh}_{n,m}} \int_{\Delta_{[0,t]}^{n+m}} f_{\sigma^{-1}(1)}(s_1) \cdots f_{\sigma^{-1}(n+m)}(s_{n+m}) ds_1 \cdots ds_{n+m},$$

where $\text{Sh}_{n,m} \subset S_{n+m}$ is the set of so-called (n, m) -shuffles, i.e., permutations σ on the set $[n+m]$, such that $\sigma(1) < \cdots < \sigma(n)$ and $\sigma(n+1) < \cdots < \sigma(n+m)$.

The above product is conveniently abstracted into an algebraically defined shuffle product on words. Let $Y := \{y_1, y_2, \dots\}$ be an alphabet, and Y^* the corresponding free monoid of words $\omega = y_{i_1} \cdots y_{i_n}$. The vector space $\mathbb{K}\langle Y \rangle$, which is freely generated by Y^* , is a commutative algebra for the shuffle product:

$$(5) \quad v_1 \cdots v_n \text{ III } v_{n+1} \cdots v_{n+m} := \sum_{\sigma \in \text{Sh}_{n,m}} v_{\sigma_1^{-1}} \cdots v_{\sigma_{n+m}^{-1}}$$

with $v_j \in Y$, $j \in \{1, \dots, n+m\}$. We define the empty word $\mathbf{1}$ as unit: $\mathbf{1} \text{ III } v = v \text{ III } \mathbf{1} = v$ for $v \in Y^*$. This product is homogenous with respect to the length of words and can be defined recursively. Indeed, one can show that:

$$(6) \quad v_1 \cdots v_n \text{ III } v_{n+1} \cdots v_{n+m} = v_1 (v_2 \cdots v_n \text{ III } v_{n+1} \cdots v_{n+m}) \\ + v_{n+1} (v_1 \cdots v_n \text{ III } v_{n+2} \cdots v_{n+m}).$$

The shuffle product was axiomatized in the 1950’s in the seminal works of Eilenberg–MacLane and Schützenberger [11, 34], and has proven to be essential in many fields of pure and applied mathematics. In [9, 10] Chen studied fundamental groups and loop spaces. In control theory, Chen’s abstract shuffle product plays a central role in Fliess’ work [16], which combines iterated integrals and formal power series in noncommutative variables into an algebraic approach to nonlinear functional expansions. Reutenauer’s monograph on free Lie algebras [33] embeds Chen’s work into an abstract Hopf algebra theoretic setting. More recently, Chen’s formalism came to prominence in Lyons’ seminal theory of rough paths [27].

When dealing with iterated Itô integrals of semimartingales, this machinery of shuffle products does not apply any more. One has instead to use the quasi-shuffle product [7]. Its definition will be recalled further below. We simply mention for the time being that, although it was discovered and investigated much more recently, it appears to be as important as the shuffle product, both from a theoretical as well as applied point of view. Indeed, it encodes the algebraic structure of discrete

sums, as the shuffle product encodes the one of integration maps, and appears in various domains, e.g., multiple zeta values, Rota–Baxter algebras, and Ecalle’s mould calculus. The latter is related to the computation of normal forms in the theory of dynamical systems. We refer to [13, 14, 17, 23] for further historical and technical details on quasi-shuffle algebras.

4. SEMIMARTINGALES

A *semimartingale* consists of a process X , which has a decomposition $X_t = X_0 + M_t + A_t$ for $t \geq 0$, where $M_0 = A_0 = 0$. The process M is a local martingale and A is an adapted process that is right-continuous with left limits and has finite variation on each finite interval $[0, t]$, see for example Jacod & Shiryaev [24]. For semimartingales X and Y the *quadratic covariation* or *square bracket process* $[X, Y]$ is defined as

$$(7) \quad [X, Y] = X \cdot Y - X_0 Y_0 - \int X_- dY - \int Y_- dX.$$

The quadratic covariation of a process X with itself is known as its *quadratic variation*. The properties of the stochastic integral imply that the quadratic covariation process defines a commutative and associative product on the space of semimartingales, i.e. $[X, Y] = [Y, X]$ and $[X, [Y, Z]] = [[X, Y], Z]$ for any semimartingales X, Y and Z , see Protter [32] and Jacod & Shiryaev [24] for details.

For notational convenience, we will write from now on iterated integrals of semimartingales as follows:

$$\int XY := \int X_- dY, \quad \text{and} \quad \int X_1 \cdots X_n := \int \left(\int X_1 \cdots X_{n-1} \right)_- dX_n.$$

Iterated brackets are denoted by:

$$(8) \quad X \star Y := [X, Y], \quad \text{and} \quad X_1 \star \cdots \star X_n := [\cdots [X_1, X_2], \cdots, X_n].$$

From now on we will assume that all processes are normalized so that $X_0 = 0$. Terms such as $X_0 Y_0$, can therefore be ignored in products of stochastic integrals, e.g., as in equation (7).

Let us briefly illustrate the combinatorial nature of iterated Itô integrals, before turning to the general, and more abstract picture. Recall first that, for arbitrary semimartingales A, B, C, D and $X := \int A_- dB, Y := \int C_- dD$, we have

$$(9) \quad X \cdot Y = \int (XC)_- dD + \int (AY)_- dB + \int (AC)_- d[B, D].$$

Example 4.1. For $B := \int dB, Y := \int C_- dD$

$$\begin{aligned} B \cdot Y &= \int (CB)_- dD + \int Y_- dB + \int C_- d[B, D] \\ &= \int CDB + \int \left(\int C_- dB + \int B_- dC + [B, C]_- \right) dD + \int C_- d[B, D] \\ &= \int BCD + \int CBD + \int CDB + \int [B, C]D + \int C[B, D]. \end{aligned}$$

Whereas the first three terms of the expansion, i.e., $\int BCD + \int CBD + \int CDB$, are those that would also appear in the shuffle product expansion, the last two terms arise from the bracket operation. These extra terms are typical outcomes of what distinguishes quasi-shuffle and shuffle computations. The algebra underlying the quasi-shuffle calculus has been explored in [17, 30]. In a nutshell, the bracket terms that appear in the above product require the replacement of permutations in the calculation of the Chen–Strichartz formula by the larger class of surjections.

5. QUASI-SHUFFLE ALGEBRA

We recall now the definition and basic properties of quasi-shuffle algebras. In spite of the fact that such algebras encode naturally Itô integral calculus, they appeared only in a few papers in the context of stochastic integration, see e.g. the works by Gaines, and Liu and Li [18, 26] and [7]. As the name indicates, quasi-shuffle algebras are obtained as a deformation of classical shuffle algebras. It is generally agreed that the idea can be traced back to the work of P. Cartier on free commutative Rota–Baxter algebras [4]. However, it was formalized only recently by M. Hoffman [23]. The link between Hoffman’s ideas and commutative Rota–Baxter algebras was explored in [13].

Abstractly, a quasi-shuffle algebra is defined as a commutative algebra B with product \bullet , which is equipped with two extra bilinear products denoted \uparrow and \downarrow (known as “half-shuffles”), such that for $x, y, z \in B$ one has

$$(10) \quad x \uparrow y = y \downarrow x,$$

$$(11) \quad (x \bullet y) \uparrow z = x \bullet (y \uparrow z)$$

$$(12) \quad (x \uparrow y) \uparrow z = x \uparrow (y \uparrow z + y \downarrow z + y \bullet z).$$

One writes usually

$$x \sqcup\sqcup y := x \uparrow y + x \downarrow y + x \bullet y,$$

and calls $\sqcup\sqcup$ the quasi-shuffle product. In particular the last axiom then reads $(x \uparrow y) \uparrow z = x \uparrow (y \sqcup\sqcup z)$. Note that when the product \bullet on B is the null product, then one recovers the usual axioms for shuffle algebras [11, 34].

Remark 5.1. Let us mention that the “deformation” induced by the \bullet product can be understood in terms of $-$ weight-one $-$ commutative Rota–Baxter algebras. The term $y \uparrow z + y \downarrow z + y \bullet z$ can then be interpreted as the so-called double Rota–Baxter product. We refrain however from developing these ideas here, since they are only indirectly relevant to our present purposes. The interested reader is referred to the survey paper [14] for an overview of the links between the theories of Rota–Baxter algebras, integral calculus and (quasi-)shuffle algebras.

The standard example of a quasi-shuffle algebra, studied in detail in [23], is provided by the tensor algebra $T(A) := \bigoplus_{n \in \mathbb{N}} A^{\otimes n}$ over a commutative algebra $(A, *)$. The three products $\uparrow, \downarrow, \bullet$ are defined inductively by: $a \bullet b := a * b$, $a \uparrow b := ba$, $a \downarrow b := ab$, and

$$a_1 \cdots a_n \uparrow b_1 \cdots b_m := (a_1 \cdots a_{n-1} \sqcup\sqcup b_1 \cdots b_m) a_n,$$

$$a_1 \cdots a_n \downarrow b_1 \cdots b_m := (a_1 \cdots a_n \sqcup\sqcup b_1 \cdots b_{m-1}) b_m,$$

$$a_1 \cdots a_n \bullet b_1 \cdots b_m := (a_1 \cdots a_{n-1} \sqcup\sqcup b_1 \cdots b_{m-1}) (a_n * b_m),$$

where we used the common word notation $a_1 \cdots a_n$ for $a_1 \otimes \cdots \otimes a_n \in A^{\otimes n}$.

Example 5.2. The product of two words of length two gives explicitly

$$\begin{aligned} a_1 a_2 \sqcup\sqcup b_1 b_2 &= (a_1 \sqcup\sqcup b_1 b_2) a_2 + (a_1 a_2 \sqcup\sqcup b_1) b_2 + (a_1 \sqcup\sqcup b_1) (a_2 * b_2) \\ &= b_1 b_2 a_1 a_2 + a_1 b_1 b_2 a_2 + b_1 a_1 b_2 a_2 + a_1 a_2 b_1 b_2 + b_1 a_1 a_2 b_2 + a_1 b_1 a_2 b_2 \\ &\quad + b_1 (a_1 * b_2) a_2 + (a_1 * b_1) b_2 a_2 + a_1 (a_2 * b_1) b_2 + (a_1 * b_1) a_2 b_2 \\ &\quad + a_1 b_1 (a_2 * b_2) + b_1 a_1 (a_2 * b_2) + (a_1 * b_1) (a_2 * b_2). \end{aligned}$$

Recall from Section 3 that the recursive description of shuffle product is complemented by its definition in terms of permutations. Similarly, the above recursive definition of the quasi-shuffle product has an explicit presentation in terms of surjections. Concretely, let f be a surjective map from $[n] = \{1, \dots, n\}$ to $[p]$. We set:

$$f(a_1 \cdots a_n) := \left(\prod_{j \in f^{-1}(1)}^* a_j \right) \otimes \cdots \otimes \left(\prod_{j \in f^{-1}(p)}^* a_j \right) \in A^{\otimes p},$$

so that for f from, say, the set $[4]$ to the set $[2]$ given by $f(1) = 1$, $f(2) = 2$, $f(3) = 1$, $f(4) = 2$, we find $f(a_1 \cdots a_4) = (a_1 * a_3) \otimes (a_2 * a_4)$.

Then, we obtain:

$$(13) \quad a_1 \cdots a_n \sqcup b_1 \cdots b_m := \sum_f f(a_1 \cdots a_n b_1 \cdots b_m),$$

where f runs over all surjections from the set $[n+m]$ to the set $[k]$, for $\max(n, m) \leq k \leq m+n$, and such that $f(1) < \cdots < f(n)$, $f(n+1) < \cdots < f(n+m)$.

Let us write now \mathcal{T} for the tensor algebra over the algebra \mathcal{S} of semimartingales, equipped with the \star product defined in (8), so that from now on $X_1 \cdots X_n$ denotes a tensor product of semimartingales in $\mathcal{S}^{\otimes n} \subset \mathcal{T}$, and $\int X_1 \cdots X_n$ the corresponding iterated stochastic integral. We finally obtain the analog for iterated integrals of semimartingales of the usual Chen formulas of iterated integrals. Recall that the latter hold for either Stratonovich or indefinite Riemann integrals.

Proposition 5.3. *The product of two iterated stochastic integrals is given by:*

$$\begin{aligned} \int X_1 \cdots X_n \cdot \int Y_1 \cdots Y_m &= \int (X_1 \cdots X_n \sqcup Y_1 \cdots Y_m) \\ &= \sum_f \int f(X_1 \cdots X_n Y_1 \cdots Y_m), \end{aligned}$$

where, as above, f runs over all surjections from the set $[n+m]$ to the set $[k]$, for $\max(n, m) \leq k \leq m+n$, and such that $f(1) < \cdots < f(n)$, $f(n+1) < \cdots < f(n+m)$.

Example 5.4. The product of two iterated stochastic integrals gives

$$\int X_1 X_2 \cdot \int X_3 = \int (X_1 X_2 X_3 + X_1 X_3 X_2 + X_3 X_1 X_2 + (X_1 \star X_3) X_2 + X_1 (X_2 \star X_3)).$$

The proposition follows from the observation that the inductive rules for the quasi-shuffle product in the tensor algebra give the pattern obeyed by products of iterated integrals of semimartingales. Namely, setting for $X := \int A_- dB$ and $Y := \int C_- dD$, we have

$$\begin{aligned} X \uparrow Y &:= \int (A(\int C_- dD))_- dB, \\ X \downarrow Y &:= \int ((\int A_- dB)C)_- dD, \\ X \bullet Y &:= \int (AC)_- d[B, D]. \end{aligned}$$

6. SURJECTIONS

This section presents a concise and mostly self-contained account on the modern algebraic theory of surjections, originating independently from F. Hivert's PhD-thesis [22] and from the work by Chapoton on the permutohedron [6]. We refer the reader to the works [17, 30] for more details on the subject.

Let us write $\mathbf{Sj}_{n,p}$ for the set of surjective maps from the set $[n] := \{1, \dots, n\}$ to the set $[p]$, $\mathbf{Sj}_{n,p}$ for its linear span, \mathbf{Sj}_n for the union of the $\mathbf{Sj}_{n,p}$, $p \leq n$, and

$$\mathbf{Sj}_n := \bigoplus_{1 \leq p \leq n} \mathbf{Sj}_{n,p}.$$

The linear span of all surjections is denoted

$$\mathbf{Sj} := \bigoplus_{n,p} \mathbf{Sj}_{n,p},$$

and will be called from now on the set of *surjective functions*.

Let us mention for the sake of completeness that surjective functions in this sense are often referred to as *word quasisymmetric functions* in the literature on algebraic combinatorics. This is because they can be encoded by formal sums of words over

an ordered alphabet. The set \mathbf{Sj} is then written \mathbf{WQSym} (this is the notation used in the articles we quoted for further details on the underlying theory). This interpretation, which we will not use in this work, permits to deduce automatically certain properties for \mathbf{Sj} from general properties of words. See the references [30, 17] for more details.

The vector space \mathbf{Sj} is naturally equipped with a Hopf algebra structure, through its action on quasi-shuffle algebras [30]. However, we will make use here only of the algebra structure. As we just saw, quasi-shuffle algebras are closely related to stochastic integration, and this is the reason why \mathbf{Sj} will prove to provide an appropriate algebraic framework for what is going to be presented in the next sections.

Let us consider a word w over the integers (a sequence of integers) whose set of letters is $I = \{i_1, \dots, i_n\}$ (e.g. $w = 35731$, $I = \{1, 3, 5, 7\}$). Let us write f for the unique increasing bijection that sends $I = \{i_1, \dots, i_n\}$ to $\{1, \dots, n\}$ (e.g. $f(1) = 1, f(3) = 2, f(5) = 3, f(7) = 4$). The *packing map* pack is the induced map on words (for example $\text{pack}(35731) := f(3)f(5)f(7)f(3)f(1) = 23421$).

Definition 6.1. *The product in \mathbf{Sj} of $f \in \mathbf{Sj}_{n,k}$ with $g \in \mathbf{Sj}_{m,l}$ is defined by $f \diamond g := \sum_h h$, where h runs over the elements in $\mathbf{Sj}_{n+m, i+j}$, $\max(l, k) \leq i + j \leq l + k$ such that*

$$\text{pack}(h(1) \cdots h(n)) = f(1) \cdots f(n), \quad \text{pack}(h(n+1) \cdots h(n+m)) = g(1) \cdots g(m).$$

This product is associative and unital (the unit can be understood as the unique surjection from the empty set $\emptyset =: [0]$ to itself), but it is not commutative.

Proposition 6.2. *The linear span of all surjection, \mathbf{Sj} , equipped with the \diamond product, is an associative, unital, non-commutative algebra.*

Associativity follows by noticing that the product $f \diamond g \diamond j$ of three surjections, where $j \in \mathbf{Sj}_{p,q}$ is obtained as the sum of all surjections h with $\text{pack}(h(1) \cdots h(n)) = f(1) \cdots f(n)$, $\text{pack}(h(n+1) \cdots h(n+m)) = g(1) \cdots g(m)$, $\text{pack}(h(n+m+1) \cdots h(n+m+p)) = j(1) \cdots j(p)$.

7. DESCENTS AND A ITÔ-TYPE BCH FORMULA

It turns out that, similar to the classical theory of iterated integrals, the most interesting computations that will take place later in this article on iterated stochastic integrals do not involve the full algebra \mathbf{Sj} , but only a small subalgebra, known as the *descent algebra* or *algebra of noncommutative symmetric functions* \mathbf{Sym} . For more details the reader is referred to the standard references [19, 33].

As an algebra, \mathbf{Sym} is the free graded associative unital algebra over generators 1_n , indexed by non-negative integers. We denote the product $*$ and set $1_{n,m} := 1_n * 1_m$. In general, for a sequence $\bar{n} := n_1, \dots, n_k$ of integers, we write $1_{\bar{n}} := 1_{n_1} * \cdots * 1_{n_k}$. As a vector space, \mathbf{Sym} is simply the linear span of the $1_{\bar{n}}$.

A surjection f in \mathbf{Sj}_n is said to have a descent in position $i < n$ if and only if $f(i) \geq f(i+1)$. The set of all descents of f is written $\text{Desc}(f)$ and

$$\text{Desc}(f) := \{i < n, f(i) \geq f(i+1)\}.$$

The number $d(f)$ that appears in Theorem 2.1 is the number of descents of f .

We also set, for $I \subset [n-1]$,

$$\begin{aligned} \text{Desc}_I^n &:= \{f \in \mathbf{Sj}_n, \text{Desc}(f) = I\}, \\ D_I^n &:= \sum_{f \in \text{Desc}_I} f \in \mathbf{Sj}, \end{aligned}$$

and

$$\begin{aligned} \text{Desc}_{\subseteq I}^n &:= \{f \in \mathbf{Sj}_n, \text{Desc}(f) \subseteq I\}, \\ D_{\subseteq I}^n &:= \sum_{f \in \text{Desc}_{\subseteq I}} f \in \mathbf{Sj}. \end{aligned}$$

When the value of n is obvious from the context, we will abbreviate D_I^n by D_I , and similarly for other symbols. Notice that $\text{Desc}_{\subseteq I} = \sum_{J \subseteq I} \text{Desc}_J$, so that, by Möbius inversion in the poset of subsets of the set $[n-1]$,

$$(14) \quad \text{Desc}_I = \sum_{J \subseteq I} (-1)^{|I|-|J|} \text{Desc}_{\subseteq J}.$$

The subsets Desc_I form a decomposition of Sj_n into a family of disjoint subsets, from which it follows that D_I and (by a triangularity argument) $D_{\subseteq I}$ form two linearly independent families in \mathbf{Sj}_n .

Lemma 7.1. *The map ι from \mathbf{Sym} to \mathbf{Sj} defined by:*

$$\iota(1_{\bar{n}}) := D_{\subseteq \{n_1, n_1+n_2, \dots, n_1+\dots+n_{k-1}\}}$$

is an injective algebra map from \mathbf{Sym} into \mathbf{Sj} .

Proof. Injectivity follows immediately from the linear independency of the $D_{\subseteq I}$. Let us show that ι is an algebra map. For arbitrary $\bar{n} = n_1, \dots, n_k$, $\bar{m} = m_1, \dots, m_l$, we have

$$\iota(1_{\bar{n}} * 1_{\bar{m}}) = D_{\subseteq \{n_1, n_1+n_2, \dots, n_1+\dots+n_k, n_1+\dots+n_k+m_1, \dots, n_1+\dots+n_k+m_1+\dots+m_{l-1}\}}.$$

On the other hand, $\iota(1_{\bar{n}}) \diamond \iota(1_{\bar{m}})$ is, by definition of the \diamond product in \mathbf{Sj} , the sum of all surjections $f \in \text{Sj}_{n_1+\dots+n_k+m_1+\dots+m_l}$ such that $\text{pack}(f(1) \cdots f(n_1 + \dots + n_k))$ lies in

$$\text{Desc}_{\subseteq \{n_1, n_1+n_2, \dots, n_1+\dots+n_{k-1}\}}$$

and $\text{pack}(f(n_1 + \dots + n_k + 1) \cdots f(n_1 + \dots + n_k + m_1 + \dots + m_l))$ lies in

$$\text{Desc}_{\subseteq \{m_1, m_1+m_2, \dots, m_1+\dots+m_{l-1}\}}.$$

Since there is no constraint on the relative values of $f(n_1 + \dots + n_k)$ and $f(n_1 + \dots + n_k + 1)$, the statement of the lemma follows. \blacksquare

The following theorem is the equivalent, in the quasi-shuffle framework, of the classical continuous Baker–Campbell–Hausdorff theorem, which computes, among others, the logarithm of the solution of a \mathfrak{g} -matrix-valued \mathfrak{g} -linear differential equation. Theorem 7.2 will appear to play the same role for matrix stochastic linear differential equations. It was first stated in [30], but the proof given in that article is indirect and relies on structure arguments from the theory of noncommutative symmetric functions. Stating those results, which are scattered in the literature on algebraic combinatorics, would go beyond the scope of this work. We propose therefore a simple and self-contained proof, which is reminiscent of the solution to the classical Baker–Campbell–Hausdorff problem stated in [33].

Let us set $I := \sum_{n=0}^{\infty} \iota(1_n) =: \sum_{n=0}^{\infty} p_n$, where we write p_n for the identity map of the set $[n]$ viewed as an element of Sj_n .

Theorem 7.2. *We have, in \mathbf{Sj} ,*

$$\begin{aligned} \log(I) &= \sum_{n=1}^{\infty} \sum_{I \subseteq [n-1]} \frac{(-1)^{|I|}}{|I|+1} \cdot D_{\subseteq I}^n \\ &= \sum_{n=1}^{\infty} \sum_{I \subseteq [n-1]} \frac{(-1)^{|I|}}{n} \cdot \binom{n-1}{|I|}^{-1} D_I^n. \end{aligned}$$

Proof. The first part of the statement follows from the computation of the logarithm of $\sum_{n=0}^{\infty} 1_n$ in \mathbf{Sym} , and from the previous Lemma. Indeed

$$\log\left(\sum_{n=0}^{\infty} 1_n\right) = \sum_{i=1}^{\infty} \frac{(-1)^{i-1}}{i} \left(\sum_{i=1}^{\infty} 1_i\right)^i = \sum_{\bar{n}=(i_1, \dots, i_k)} \frac{(-1)^{k-1}}{k} \cdot 1_{\bar{n}}.$$

Let us now expand the second term of the identity in the theorem in terms of the D_I^n . This yields the coefficient of D_I^n , $I = \{i_1, \dots, i_k\}$ given by:

$$\begin{aligned} \sum_{I \subseteq J \subseteq [n-1]} \frac{(-1)^{|J|}}{|J|+1} &= \sum_{j=0}^{n-k-1} \frac{(-1)^{j+k}}{j+k+1} \binom{n-k-1}{j} \\ &= (-1)^k \sum_{j=0}^{n-k-1} \int_0^1 (-1)^j \binom{n-k-1}{j} x^{k+j} dx \\ &= (-1)^k \int_0^1 (1-x)^{n-k-1} x^k dx = (-1)^k \frac{1}{n} \binom{n-1}{k}^{-1}, \end{aligned}$$

from which the theorem follows. \blacksquare

In view of Proposition 5.3, Theorem 2.1 also follows.

The elements $\sum_{I \subseteq [n-1]} \frac{(-1)^{|I|}}{n} \cdot \binom{n-1}{|S|}^{-1} D_I^n$ are analogs in \mathbf{Sj} of the celebrated Solomon idempotents, see [30] for further details.

8. NONCOMMUTATIVE STOCHASTIC CALCULUS

In the previous sections of the article, we investigated quasi-shuffle-type properties of iterated integrals of semimartingales. We also showed the strong relationship between them and properties of surjections. Now Proposition 5.3 is restated and generalized:

Proposition 8.1. *The product of k iterated stochastic integrals of semimartingales is given by:*

$$\begin{aligned} &\left(\int X_1^1 \cdots X_{n_1}^1 \right) \cdot \cdots \cdot \left(\int X_1^k \cdots X_{n_k}^k \right) \\ &= \sum_{f \in \text{Desc}_{\subseteq \{n_1, \dots, n_1 + \dots + n_{k-1}\}}} \int f(X_1^1 \cdots X_{n_1}^1 \cdots X_1^k \cdots X_{n_k}^k). \end{aligned}$$

More generally, for $f_1 \in \mathbf{Sj}_{n_1}, \dots, f_k \in \mathbf{Sj}_{n_k}$:

$$\begin{aligned} &f_1 \left(\int X_1^1 \cdots X_{n_1}^1 \right) \cdot \cdots \cdot f_k \left(\int X_1^k \cdots X_{n_k}^k \right) \\ &= (f_1 \diamond \cdots \diamond f_k) \left(\int X_1^1 \cdots X_{n_1}^1 \cdots X_1^k \cdots X_{n_k}^k \right). \end{aligned}$$

In the last formula, from which the first follows, it is implicitly assumed that the action of \mathbf{Sj}_k on $\int X_1 \cdots X_k$ is extended linearly to the linear span of \mathbf{Sj}_k , that is, the action of a linear combination of surjections f_i is the linear combination of the actions of the f_i .

We invite the reader to check that the $k = 2$ case of the last formula follows from the definition of $f(\int X_1^1 \cdots X_{n_1}^1)$ and from Proposition 5.3. The general case follows by induction.

Example 8.2. The expansion of the triple product $(\int X) \cdot (\int Y_1 Y_2) \cdot (\int Z_1 Z_2)$ includes terms such as:

$$\int XY_1(Y_2 \star Z_1)Z_2, \int (X \star Y_1)(Y_2 \star Z_1)Z_2 \text{ and } \int Z_1(X \star Y_1 \star Z_2)Y_2.$$

The purpose of the present section is to extend this picture to the operator setting, that is, to iterated stochastic integrals of, say, $n \times n$ square matrices $M = M(X^{i,j})_{1 \leq i, j \leq n}$ whose entries $M^{i,j} = X^{i,j}$ are semimartingales. Note that we write the indices of the entries as exponents for notational convenience in forthcoming computations. The set of such matrices is denoted \mathfrak{M} .

For matrices $M_1, \dots, M_k \in \mathfrak{M}$, we set:

$$\int M_1 \cdots M_k := \left(\sum_{i_1, \dots, i_k} \int M_1^{i_1, i_1} M_2^{i_1, i_2} \cdots M_k^{i_k, j} \right)_{1 \leq i, j \leq n}^{i, j},$$

and for $f \in \mathbf{Sj}_{k, l}$,

$$f\left(\int M_1 \cdots M_k\right) := \left(\sum_{i_1, \dots, i_k} \int f\left(M_1^{i_1, i_1} M_2^{i_1, i_2} \cdots M_k^{i_k, j}\right) \right)_{1 \leq i, j \leq n}^{i, j}.$$

Last, if $F = \sum_i \lambda_i f_i$ is a linear combination of surjections in \mathbf{Sj}_k , then we set:

$$F\left(\int M_1 \cdots M_k\right) := \sum_i \lambda_i f_i\left(\int M_1 \cdots M_k\right).$$

The product rule of Proposition 8.1 applies (entry-wise), and we obtain, for $M_1, \dots, M_k, N_1, \dots, N_m \in \mathfrak{M}$:

$$\begin{aligned} & \int M_1 \cdots M_k \cdot \int N_1 \cdots N_m \\ &= \left(\sum_{f \in \text{Desc}_{\subseteq \{k\}}} \sum_{i_1, \dots, i_{k+m}} \int f\left(M_1^{i_1, i_1} \cdots M_k^{i_k, i_{k+1}} N_1^{i_{k+1}, i_{k+2}} \cdots N_m^{i_{k+m}, i_j}\right) \right)_{1 \leq i, j \leq n}^{i, j} \\ &= \sum_{f \in \text{Desc}_{\subseteq \{k\}}} f\left(\int M_1 \cdots M_k N_1 \cdots N_m\right). \end{aligned}$$

Example 8.3. Let $k = 2, m = 1$, and consider 2×2 matrices, this gives for the first entry of the product:

$$\begin{aligned} \left(\int M_1 M_2 \cdot \int N_1\right)^{1,1} &= \sum_{i, j \leq 2} \int \left(M_1^{1, i} M_2^{i, j} N_1^{j, 1} + M_1^{1, i} N_1^{j, 1} M_2^{i, j} + N_1^{j, 1} M_1^{1, i} M_2^{i, j} \right. \\ &\quad \left. + M_1^{1, i} (M_2^{i, j} \star N_1^{j, 1}) + (M_1^{1, i} \star N_1^{j, 1}) M_2^{i, j} \right). \end{aligned}$$

For higher products we obtain similarly:

Proposition 8.4. For $M_1^1, \dots, M_{n_1}^1, M_1^k, \dots, M_{n_k}^k \in \mathfrak{M}$, we have:

$$\begin{aligned} & \left(\int M_1^1 \cdots M_{n_1}^1\right) \cdots \left(\int M_1^k \cdots M_{n_k}^k\right) \\ &= \sum_{f \in \text{Desc}_{\subseteq \{n_1, \dots, n_1 + \dots + n_{k-1}\}}} f\left(\int M_1^1 \cdots M_{n_1}^1 \cdots M_1^k \cdots M_{n_k}^k\right) \end{aligned}$$

and more generally, for $f_1 \in \mathbf{Sj}_{n_1}, \dots, f_k \in \mathbf{Sj}_{n_k}$ we obtain:

$$\begin{aligned} & f_1\left(\int M_1^1 \cdots M_{n_1}^1\right) \cdots f_k\left(\int M_1^k \cdots M_{n_k}^k\right) \\ &= (f_1 \diamond \cdots \diamond f_k)\left(\int M_1^1 \cdots M_{n_1}^1 \cdots M_1^k \cdots M_{n_k}^k\right). \end{aligned}$$

The proposition follows from the linear case (Proposition 8.1) by expanding entry-wise the products of matrices.

We are now in the position to calculate the logarithm of the Itô-Taylor series.

Theorem 8.5. For an arbitrary matrix $M \in \mathfrak{M}$ and $X = \sum_{n=0}^{\infty} \int M^n$ the (formal) solution of the stochastic differential equation $dX = X \cdot dM$, $X_0 := Id$, we have:

$$\log(X) = \sum_{n=1}^{\infty} \sum_{I \subseteq [n-1]} \frac{(-1)^{|I|}}{n} \cdot \binom{n-1}{|I|}^{-1} D_I^n \int M^n.$$

This formula may provide the basis for interesting numerical properties. For instance, truncating the expansion of $\log(X)$ at order k , that is, looking at

$$\sum_{n=1}^k \sum_{I \subseteq [n-1]} \frac{(-1)^{|I|}}{n} \cdot \binom{n-1}{|S|}^{-1} D_I^n \int M^n,$$

and applying the exponential map, can be expected (in view of similar phenomena in the deterministic case) to provide a better approximation to X , than the truncation of the original expansion $\sum_{i=0}^k \int M^i$.

The first few terms of the expansion of $\log(X)$ read:

$$\begin{aligned} \log(X) = & \int M + \left(\frac{1}{2}(12) - \frac{1}{2}((21) + (11)) \right) \int M^2 \\ & + \left(\frac{1}{3}(123) - \frac{1}{6}((213) + (312) + (112) + (212)) \right. \\ & - \frac{1}{6}((132) + (231) + (122) + (121)) \\ & \left. + \frac{1}{3}((321) + (211) + (111) + (221)) \right) \int M^3 + \dots \end{aligned}$$

We remind the reader that we represent a surjection $f \in \text{Sj}_k$ by the sequence of its values $(f(1) \cdots f(k))$.

Example 8.6. For 2×2 matrices, the $(1, 2)$ -entry of the term $(11) \int M^2$ is given by $\sum_{i \leq 2} \int M^{1,i} \star M^{i,2}$; the one of $(231) \int M^3$ reads $\sum_{i,j \leq 2} \int M^{j,2} M^{1,i} M^{i,j}$; the one of $(221) \int M^3$ reads $\sum_{i,j \leq 2} \int M^{j,2} (M^{1,i} \star M^{i,j})$.

REFERENCES

- [1] O. E. Barndorff-Nielsen, T. Mikosch, S. I. Resnick, *Lévy Processes: Theory and Applications*, New York: Springer 2001.
- [2] F. Baudoin, *An Introduction To The Geometry Of Stochastic Flows*, World Scientific Publishing Company, Singapore, 2005.
- [3] Gérard Ben Arous, *Flots et séries de Taylor stochastiques*, Probab. Th. Rel. Fields **81**, no. 1, 29 (1989).
- [4] P. Cartier, *On the structure of free Baxter algebras*, Adv. Mathematics **9**, 253 (1972).
- [5] F. Castell, J. Gaines, *An efficient approximation method for stochastic differential equations by means of the exponential Lie series*, Math. Comp. Simulation **38**, 13 (1995).
- [6] F. Chapoton, *Algèbres de Hopf des permutahédres, associahédres et hypercubes*, Adv. in Mathematics **150**, 264 (2000).
- [7] C. Curry, K. Ebrahimi-Fard, S.J. Malham, A. Wiese, *Lévy Processes and Quasi-Shuffle Algebras*, Stochastics **86**, 632 (2014).
- [8] K.T. Chen, *Integration of paths, geometric invariants and a generalized Baker–Hausdorff formula*, Ann. Math. **65**, 163 (1957).
- [9] K.T. Chen, *Algebras of iterated path integrals and fundamental groups*, Transactions of the American Mathematical Society **156**, 359 (1971).
- [10] K.T. Chen, *Iterated path integrals*, Bulletin of the American Mathematical Society **83**(5), 831 (1977).
- [11] S. Eilenberg, S. Mac Lane, *On the Groups $H(\pi, n)$* , Annals of Mathematics, Second Series, **58**, no. 1, 55 (1953).
- [12] K. Ebrahimi-Fard, *Loday-type algebras and the Rota–Baxter relation* Letters in Mathematical Physics **61**(2), 139 (2002).
- [13] K. Ebrahimi-Fard, L. Guo, *Quasi-shuffles, Mixable Shuffles, and Hopf Algebras*, Journal of Algebraic Combinatorics **24**(1), 83 (2006).
- [14] K. Ebrahimi-Fard, F. Patras, *La structure combinatoire du calcul intégral*, Gazette des Mathématiciens **138**, (2013).
- [15] K. Ebrahimi-Fard, F. Patras, *The pre-Lie structure of the time-ordered exponential* Letters in Mathematical Physics **104**, no. 10, 1281 (2014).
- [16] M. Fliess, *Fonctionnelles causales non linéaires et indéterminées non commutatives* Bull. Soc. Math. France **109**, 3 (1981).
- [17] L. Foissy, F. Patras, J.-Y. Thibon, *Deformations of shuffles and quasi-shuffles*, Ann. Institut Fourier, to appear.
- [18] J. Gaines, *The algebra of iterated stochastic integrals*, Stochastics and Stoch. Reports **49**, 169 (1994).
- [19] I. M. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V. S. Retakh, J.-Y. Thibon, *Noncommutative symmetric functions*, Adv. in Mathematics **112**, 218 (1995).
- [20] M. Hairer, D. Kelly, *Geometric versus non-geometric rough paths*, Ann. Inst. H. Poincaré Probab. Statist. **51**, No. 1, 207 (2015).
- [21] M. Hairer, *A theory of regularity structures*, Inventiones Mathematicae **198**(2), 269 (2014).
- [22] F. Hivert, *Combinatoire des fonctions quasi-symétriques*, Thèse de doctorat (Doctoral thesis), 1999.
- [23] M. E. Hoffman, *Quasi-shuffle products* Journal of Algebraic Combinatorics **11**, 49 (2000).
- [24] J. Jacod, A.N. Shiryaev, *Limit Theorems for Stochastic Processes*. Berlin, etc: Springer 2002.

- [25] I. Karatzas, S. Shreve, *Brownian Motion and Stochastic Calculus*, Graduate Texts in Mathematics Vol. 113, 2nd Edition, Berlin, etc: Springer 2000.
- [26] C.W Li, X.Q. Liu, *Algebraic structure of multiple stochastic integrals with respect to Brownian motions and Poisson processes*, Stochastics and Stoch. Reports **61**, 107 (1997).
- [27] T. Lyons, M. Caruana, T. Lévy, *Differential Equations Driven by Rough Paths*, École d'été des probabilités de Saint-Flour XXXIV 2004 (J. Picard, ed.). Lecture Notes in Mathematics **1908**.
- [28] S.J.A. Malham, A. Wiese, *Stochastic Lie group integrators*, SIAM J. Sci. Comput. **30**(2), 597 (2008).
- [29] B. Mielnik, J. Plebański, *Combinatorial approach to Baker–Campbell–Hausdorff exponents* Ann. Inst. Henri Poincaré A **XII**, 215 (1970).
- [30] J.C. Novelli, F. Patras, J.-Y. Thibon, *Natural endomorphisms of quasi-shuffle Hopf algebras* Bull. Soc. Math. France **141** (1), 107 (2013).
- [31] F. Patras, *L'algèbre des descentes d'une bigèbre graduée*, J. Algebra **170**, 2, 547 (1994).
- [32] P. Protter, *Stochastic Integration and Differential Equations*, Berlin, Springer 1992.
- [33] Ch. Reutenauer, *Free Lie algebras*. London Mathematical Society Monographs New Series 7. Oxford Science Publications 1993.
- [34] M. P. Schützenberger, *Sur une propriété combinatoire des algèbres de Lie libres pouvant être utilisée dans un problème de mathématiques appliquées*, Séminaire Dubreil–Jacotin Pisot (Algèbre et théorie des nombres), Paris, Année 1958/59.
- [35] R. S. Strichartz, *The Campbell–Baker–Hausdorff–Dynkin formula and solutions of differential equations*, J. Func. Anal. **72**, 320 (1987).

(a) ICMAT, CSIC-UAM-UCM-UC3M, C/ NICOLÁS CABRERA 13-15, 28049 MADRID, SPAIN.
ON LEAVE FROM UHA, MULHOUSE, FRANCE.

(b) MAXWELL INSTITUTE FOR MATHEMATICAL SCIENCES AND SCHOOL OF MATHEMATICAL AND COMPUTER SCIENCES, HERIOT-WATT UNIVERSITY, EDINBURGH EH14 4AS, UK.

(c) LABO. DE MATHÉMATIQUES J.A. DIEUDONNÉ, UNIVERSITÉ DE NICE - SOPHIA ANTIPOLIS, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE.