



Heriot-Watt University
Research Gateway

Some examples of separable convex-cocompact subgroups

Citation for published version:

Hagen, M & Sisto, A 2023, 'Some examples of separable convex-cocompact subgroups', *Bulletin of the London Mathematical Society*, vol. 55, no. 5, pp. 2242-2257. <https://doi.org/10.1112/blms.12848>

Digital Object Identifier (DOI):

[10.1112/blms.12848](https://doi.org/10.1112/blms.12848)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Bulletin of the London Mathematical Society

Publisher Rights Statement:

© 2023 The Authors. Bulletin of the London Mathematical Society is copyright © London Mathematical Society.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

Some examples of separable convex-cocompact subgroups

Mark Hagen¹ | Alessandro Sisto²¹School of Mathematics, University of Bristol, Bristol, UK²Maxwell Institute and Department of Mathematics, Heriot-Watt University, Edinburgh, UK**Correspondence**

Mark Hagen, School of Mathematics, University of Bristol, Woodland Road, Bristol BS8 1UG, UK.

Email: markfhagen@posteo.net**Abstract**

Reid asked whether all convex-cocompact subgroups of mapping class groups are separable. Using a construction of Manning–Mj–Sageev, we give examples of separable convex-cocompact subgroups that are free of arbitrary finite rank, while prior examples seem to all be virtually cyclic.

MSC 2020

20F65, 57K20 (primary)

1 | INTRODUCTION

A subgroup H of a group G is *separable (in G)* if, for all $g \notin H$, there exists a finite group F and a homomorphism $\phi : G \rightarrow F$ such that $\phi(g) \notin \phi(H)$. Separability arises naturally in topology when, given an immersed submanifold N of a manifold M , one tries to find a finite cover of M where N embeds; see [31]. Notably, separability, and more specifically separability of quasiconvex subgroups of certain hyperbolic groups, played a crucial role in the resolution of the Virtual Haken and Virtual Fibring Conjectures [2, 32].

In the context of mapping class groups, the closest analogues of quasiconvex subgroups of hyperbolic groups are *convex-cocompact* subgroups, as defined in [14]; see Subsection 2.1 for the characterisation that we will use. In analogy with the case of hyperbolic manifolds, Reid asked whether convex-cocompact subgroups of mapping class groups are separable [30, Question 3.5]. As far as we are aware, the only examples of convex-cocompact subgroups known to be separable are virtually cyclic, as covered by [20]. However, it is shown in [6] that if ‘enough’ hyperbolic groups are residually finite then all convex-cocompact subgroups are separable. (Note that it is known that not all finitely generated subgroups of mapping class groups are separable [20].)

The main goal of this paper is to provide the first examples of separable convex-cocompact subgroups that are not virtually cyclic, using the criterion in Theorem 1.1.

Write Σ_g to mean the closed connected orientable surface of genus g and $MCG(\Sigma_g)$ its mapping class group. Given $H \leq MCG(\Sigma_g)$, denote by Γ_H the corresponding extension of H . This can be seen as the preimage of H under the natural map $MCG(\Sigma_g - \{p\}) \rightarrow MCG(\Sigma_g)$ occurring in the Birman exact sequence (see, for example, [14, Section 1.2] for detailed description of Γ_H). When H is convex-cocompact, Γ_H is hyperbolic [17] (if in addition H is free, as it will be in our examples, hyperbolicity of Γ_H is also [14, Theorem 1.3]).

The criterion is:

Theorem 1.1. *Let $g \geq 1$ and let H be a torsion-free malnormal convex-cocompact subgroup of $MCG(\Sigma_g)$. If Γ_H is conjugacy separable, then H is separable in $MCG(\Sigma_g)$.*

To apply the criterion, we need examples of convex-cocompact subgroups H with Γ_H conjugacy separable. These come from work of Manning–Mj–Sageev [23] — who gave examples where Γ_H is virtually compact special — combined with Minasyan–Zalesskii’s result on conjugacy separability of hyperbolic virtually special groups [27]. So, the examples provided by the following theorem come from [23]; we just observe that their construction gives rise to malnormal subgroups in many cases.

Theorem 1.2. *For every $g \geq 3$ and $n \geq 1$, there exist infinitely many conjugacy classes of malnormal convex-cocompact subgroups $H < MCG(\Sigma_g)$ such that H is isomorphic to a free group of rank n whose corresponding extension Γ_H is virtually compact special.*

Recalling that convex-cocompactness of H implies hyperbolicity of Γ_H , combining the above theorem with [27, Theorem 1.1] and Theorem 1.1 gives:

Corollary 1.3. *For every $g \geq 3$ and $n \geq 1$ there exist infinitely many conjugacy classes of separable convex-cocompact subgroups H of $MCG(\Sigma_g)$, with each $H \cong F_n$.*

We note that Theorem 1.1 does not apply to arbitrary cyclic convex-cocompact subgroups, even though, in this case, Γ_H is the fundamental group of a fibred hyperbolic 3-manifold, which is virtually compact special [2, 12]. Indeed, such cyclic subgroups may not be malnormal, and the maximal elementary subgroups containing them may not be torsion free (in case the maximal elementary subgroup containing the cyclic group is torsion free, the criterion does apply, though). We wonder:

Question 1.4. To what extent can malnormality and/or torsion freeness be relaxed in Theorem 1.1?

Outline of proofs

In this paper, we did not pursue the most general versions of our intermediate results, instead leaving questions that we believe to be of independent interest; we highlight the main ones below.

To prove Theorem 1.1, we use a version of Grossman’s criterion for residual finiteness of an outer automorphism group [15] (Grossman’s criterion is one way to prove residual finiteness of $MCG(\Sigma_g)$). This is Proposition 2.5 and has two hypotheses. The first one, in the case

of cocompact subgroups, is conjugacy separability of the extension group. The second one translates to a question about automorphisms of surface groups. Roughly, given a surface group G , a group K of automorphisms of G , and automorphism ϕ not in K , one can ask whether there exists $x \in G$ such that $\phi(x)$ is not conjugate to $k(x)$ for any $k \in K$. In our case, we are interested in $K = \Gamma_H$ for some convex-cocompact subgroup H (where the mapping class group of the punctured surface can be identified with an index 2 subgroup of the automorphism group of G).

We check this second condition for suitable H in two steps. First, Proposition 3.1 reduces the problem above to an analogous problem where K is replaced by finitely many automorphisms. This is done by showing that left cosets of H contain finitely many elements that are not pseudo-Anosov. This is where we use the hypotheses that H is torsion free and malnormal.

The second step is to construct a suitable x , given finitely many automorphisms. We do so specifically for surface groups in Proposition 4.1, but, inspired by the fact that Grossman's criterion can be applied to acylindrically hyperbolic groups [3], we ask:

Question 1.5. Let G be a torsion-free acylindrically hyperbolic group, and let ϕ_1, \dots, ϕ_n be non-inner automorphisms of G . Does there exist $x \in G$ with x and $\phi_i(x)$ non-conjugate for all i ?

In fact, we do not know the answer to this question even in the special case where G is hyperbolic. One way of answering the question affirmatively would be to show that a 'generic' x has the required property, which would be interesting in its own right. That is:

Question 1.6. Let G be a torsion-free acylindrically hyperbolic group, let ϕ be a non-inner automorphism of G , and let (w_n) be a simple random walk on G . Is it true that, with probability going to 1 as n goes to infinity, w_n is not conjugate to $\phi(w_n)$?

This question is already of interest for G hyperbolic and the techniques of [21] seem relevant.

2 | PRELIMINARIES

2.1 | Curve graphs and convex-cocompactness

Given a finite-type surface Σ , we denote by $C(\Sigma)$ its curve graph. For all finite-type surfaces Σ , the curve graph $C(\Sigma)$ is hyperbolic [24] and, in fact, say, 100-hyperbolic [18], see also [5, 9, 29]. Also, the action of $MCG(\Sigma)$ on $C(\Sigma)$ is acylindrical [8]. A finitely generated subgroup of $MCG(\Sigma)$ is *convex-cocompact* if the orbit maps to $C(\Sigma)$ are quasi-isometric embeddings [19, Theorem 1.3].

2.2 | Elementary subgroups

Throughout the paper, we use the following notation. If X is a hyperbolic space and Y is a quasiconvex subspace, then the closest point projection π_Y is the bounded set defined by

$$\pi_Y(x) = \{y \in Y : d(x, y) \leq d(x, Y) + 1\}.$$

If a group A acts by isometries on X , then π_Y is $\text{Stab}_A(Y)$ -equivariant. Next, we summarise properties of *elementary closures* of loxodromic elements used later, as first considered by [7].

Proposition 2.1. *Let the group G act acylindrically on the hyperbolic space X . There exists a constant Q such that the following holds. Let $g \in G$ be a loxodromic element. Then:*

- (1) *there exists a unique maximal elementary subgroup $E(g)$ containing g , and a Q -quasiconvex $E(g)$ -invariant Q -quasiline $Y = Y_g$;*
- (2) *for every $D \geq 0$ there exists $L \geq 0$ (depending on D and g), such that the following holds. If $h \in G$ has the property that there exist $x, y \in Y$ with $d(x, y) \geq L$ and $d(x, h \cdot x), d(y, h \cdot y) \leq D$, then $h \in E(g)$.*

Proof. Item (1) is given by [11, Lemma 6.5]. Item (2) is [7, Proposition 6(2)]. □

2.3 | Intersection of conjugates versus coarse intersections of coset orbits

We also need a certain correspondence between the behaviour of orbits in $C(\Sigma)$ of cosets of convex-cocompact subgroups and intersections of conjugates. The next lemma, which is used in the proof of Proposition 3.1, follows from arguments in [4]. We note that [1, Proposition 4.1] is related to the lemma, though neither implies the other.

Lemma 2.2. *Let $g \geq 2$ and let H be a convex-compact subgroup of $MCG(\Sigma_g)$. Let $o \in C(\Sigma_g)$. Suppose that for some $x \in MCG(\Sigma_g)$ and some $C \geq 0$, the diameter of $xH \cdot o \cap N_C(H \cdot o) \subseteq C(\Sigma_g)$ is infinite. Then the subgroup $H \cap xHx^{-1}$ is infinite.*

Proof. The convex-cocompact subgroup H in the statement, acting on $C(\Sigma_g)$, satisfies an appropriately modified version of the *WPD condition* from [4, Definition 2.12]. More specifically, said definition is stated for the action on a Cayley graph with respect to a possibly infinite generating sets. The equivariant quasi-isometry classes of said Cayley graphs are in natural bijection with geodesic spaces with a cobounded action, so dealing with Cayley graphs or cobounded actions is equivalent.

The proof of [4, Lemma 5.2], and hence of [4, Proposition 5.3], only uses the WPD property and hence it goes through in our setting (replacing subgroups/cosets with their orbits in $C(\Sigma_g)$ etc.). In particular, we conclude that $xH \cdot o \cap N_C(H \cdot o) \subseteq C(\Sigma_g)$ lies within finite Hausdorff distance of $(H \cap xHx^{-1}) \cdot o$. Since the former has infinite diameter, we have that $H \cap xHx^{-1}$ is infinite, as required. □

2.4 | Concatenating geodesics in hyperbolic spaces

Later, we will verify that certain elements are pseudo-Anosov by constructing quasigeodesic axes using the following lemma, which we prove by reducing it to [16, Lemma 4.2].

Lemma 2.3. *For every $\delta \geq 0$ there exists a constant C_δ such that the following holds. Let $(\alpha_i)_{i \in \mathbb{Z}}$ be geodesics in a δ -hyperbolic geodesic space, of length bounded independently of i , with the terminal point of α_i being the initial point of α_{i+1} . Let*

$$d_{i,j} = \text{diam}(N_{3\delta}(\alpha_i) \cap \alpha_j)$$




FIGURE 1 An illustration of the hypotheses in Lemma 2.3. The long horizontal segment is α_i and the central subsegment is α'_i .

and assume that for all even i ,

$$\ell(\alpha_i) \geq d_{i,i+1} + d_{i,i+2} + d_{i,i-1} + d_{i,i-2} + C_\delta.$$

Then the concatenation of the α_i is a quasigeodesic (Figure 1).

Proof. We can assume $\delta > 0$ for convenience. For i even, let α'_i be the subgeodesic of α_i obtained by removing the initial open segment of length $d_{i,i-1} + d_{i,i-2}$ and the final open segment of length $d_{i,i+1} + d_{i,i+2}$.

For i odd, let α'_i be a geodesic joining the terminal point of α'_{i-1} to the initial point of α'_{i+1} . So, the geodesics α'_i concatenate to form a bi-infinite path.

Considering thin quadrangles, we now show that that [16, Lemma 4.2] applies, that is, we show that for i even, the intersections

$$N_{3\delta}(\alpha'_i) \cap \alpha'_{i\pm 1}$$

and

$$N_{3\delta}(\alpha'_i) \cap \alpha'_{i\pm 2}$$

each have diameter at most 10δ .

We will do the argument for α'_{i+1} and α'_{i+2} , the other arguments being analogous. Denoting by α''_i the terminal segment of α_i that starts at the terminal point of α'_i , note that α'_{i+1} is contained in the 2δ -neighbourhood of $\alpha''_i \cup \alpha_{i+1} \cup \alpha_{i+2}$, and clearly so is α'_{i+2} . Hence, it suffices to show that there is no point p on α'_i further than 20δ from the terminal point of α'_i but within 5δ of $\alpha_{i+1} \cup \alpha_{i+2}$ (there is clearly no point p on α'_i further than 20δ from the terminal point and within 5δ of α''_i).

Suppose by contradiction that such p exists, and in fact that it lies within 5δ of α_{i+2} , the other case being easier. The following argument is illustrated in Figure 2. Let a_i be the terminal point of α'_i , and let β_i be the union of the terminal segment of α'_i of length $d(a_i, p) - 8\delta$, and α''_i .

Consider the 2δ -thin quadrilateral formed by the following four geodesics:

- a geodesic σ of length at most 5δ from p to a point on α_{i+2} ;
- the subpath of α_{i+2} subtended by the terminal point of σ and $\alpha_{i+1} \cap \alpha_{i+2}$;




FIGURE 2 Proof of the non-existence of p in Lemma 2.3. Note that $d(a_i, p) \geq 20\delta$.

- α_{i+1} ;
- the terminal subpath of α_i from p to the initial point of α_{i+1} .

By 2δ -thinness of the quadrilateral, the fourth path on the list is contained in the union of the 2δ -neighbourhoods of the other three paths. Since the initial point of β_i is 8δ -far from p , and $|\sigma| \leq 5\delta$, we see that no point of β_i is 2δ -close to σ , for otherwise it would be 7δ -close to p , a contradiction. Hence any point on β_i lies 2δ -close to either α_{i+1} or α_{i+2} .

Note that $\beta_i \cap N_{2\delta}(\alpha_{i+1})$ has diameter at most $d_{i,i+1} + 4\delta$. Indeed, if $p, q \in \beta_i$ are, respectively, 2δ -close to points $p', q' \in \alpha_{i+1}$, then p', q' are in $N_{3\delta}(\alpha_i) \cap \alpha_{i+1}$, so $d(p', q') \leq d_{i,i+1}$, and hence, by the triangle inequality, $d(p, q) \leq d_{i,i+1} + 4\delta$. Similarly, $\beta_i \cap N_{2\delta}(\alpha_{i+2})$ has diameter at most $d_{i,i+2} + 4\delta$. Since these two intersections cover β_i , we conclude that $|\beta_i| \leq d_{i,i+1} + d_{i,i+2} + 8\delta$.

On the other hand, β_i is the concatenation of a terminal segment of α'_i of length at least 12δ (since p is assumed to lie 20δ -far from the terminal point of α'_i) with the path α''_i (whose length is $d_{i,i+1} + d_{i,i+2}$). Hence $|\beta_i| > d_{i,i+1} + d_{i,i+2} + 8\delta$. This is a contradiction, so the point p cannot exist.

Hence, for sufficiently large C_δ , [16, Lemma 4.2] applies, yielding that the concatenation of the α'_i is a quasigeodesic. Hence, so is the concatenation of the α_i , since all geodesics involved have uniformly bounded length. □

2.5 | Grossman criterion

We now give the generalisation of Grossman’s residual finiteness criterion mentioned in the introduction. Assume that the group G has trivial centre, and let $i : G \rightarrow \text{Inn}(G)$ be the natural isomorphism defined by $i(g)(x) = gxg^{-1}$.

Remark 2.4. For any $x \in G$ and $\gamma \in \text{Aut}(G)$,

$$\gamma i(x) \gamma^{-1} = i(\gamma(x)),$$

just by expanding the definition of i .

We also need the usual exact sequence

$$1 \rightarrow \text{Inn}(G) \rightarrow \text{Aut}(G) \xrightarrow{\Psi} \text{Out}(G) \rightarrow 1.$$

The separability version of Grossman’s criterion is:

Proposition 2.5. *Let G be a finitely generated group with trivial centre and let $H < \text{Out}(G)$. Let $\Gamma_H = \Psi^{-1}(H)$ and let $\phi \in \text{Aut}(G)$. Suppose that*

- (1) *there exists $x \in G$ such that $\phi(x) \neq h(x)$ for all $h \in \Gamma_H$;*
- (2) *Γ_H is conjugacy separable.*

Then there exists a finite group F and a homomorphism $q : \text{Out}(G) \rightarrow F$ with $q(\Psi(\phi)) \notin q(H)$.

Remark 2.6. Taking $H = \{1\}$ in the above proposition yields [15, Theorem 1] in the case where G has trivial centre.

Proof of Proposition 2.5. Item (1) provides $x \in G$ such that $\phi(x) \notin \Gamma_H \cdot x$.

First observe that $i(\phi(x))$, which is contained in Γ_H , cannot be conjugate in Γ_H to $i(x)$. Indeed, if it were, then for some $\gamma \in \Gamma_H$, we would have $i(\phi(x)) = \gamma i(x) \gamma^{-1} = i(\gamma(x))$, with the second equality following from Remark 2.4. Injectivity of i would then imply $\gamma(x) = \phi(x)$, a contradiction.

Hence, by item (2), there is a finite quotient $\pi : \Gamma_H \rightarrow F_0$ such that $\pi(i(\phi(x)))$ is not conjugate in F_0 to $\pi(i(x))$. It follows that $\pi(i(\phi(x))) \notin \pi(i(\Gamma_H \cdot x))$. Indeed, by Remark 2.4, for all $\gamma \in \Gamma_H$, we have that $\pi(i(\gamma(x)))$ is conjugate in F_0 to $\pi(i(x))$, which is not conjugate to $\pi(i(\phi(x)))$.

Consider the quotient $\pi \circ i : G \rightarrow \bar{G} \leq F_0$. Let K be the intersection of all subgroups of G with the same index as $\text{Ker}(\pi \circ i)$. This is a characteristic subgroup, and furthermore, since G is finitely generated, K has finite index in G . Denote the corresponding quotient map $q'' : G \rightarrow G/K$. Since $\pi \circ i$ factors through q'' we have $q''(\phi(x)) \notin q''(\Gamma_H \cdot x)$.

Since K is characteristic, q'' induces a homomorphism $q' : \text{Aut}(G) \rightarrow \text{Aut}(G/K)$ defined by $q'(\gamma)(gK) = q''(\gamma(g))$. This has the property that $q'(\phi) \notin q'(\Gamma_H)$, because $q'(\phi)(xK) \notin q'(\Gamma_H)(xK)$, since the former is $q''(\phi(x))$ and the latter is $q''(\Gamma_H \cdot x)$.

The homomorphism q' descends to a homomorphism $q : \text{Out}(G) \rightarrow \text{Out}(G/K)$ fitting into the following commutative diagram:

$$\begin{array}{ccc} \text{Aut}(G) & \xrightarrow{\Psi} & \text{Out}(G) \\ q' \downarrow & & \downarrow q \\ \text{Aut}(G/K) & \xrightarrow{\Psi_K} & \text{Out}(G/K) \end{array}$$

Since $q'(\phi) \notin q'(\Gamma_H)$ and $\text{Inn}(G/K) = q'(\text{Inn}(G)) < q'(\Gamma_H)$, we have $\Psi_K(q'(\phi)) \notin \Psi_K(q'(\Gamma_H))$. In view of the commutative diagram, the former is $q(\Psi(\phi))$ and the latter is $q(H)$, so that we found a finite quotient of $\text{Out}(G)$ that separates $\Psi(\phi)$ from H , namely $F = \text{Out}(G/K)$, which is finite since K has finite index in G . □

3 | MAPPING CLASS GROUP LEMMAS

In Section 3.1, we reduce condition (1) from Proposition 2.5 to a statement about finitely many automorphisms of a surface group; this is used in the proof of Theorem 1.1. In Section 3.2, we prove a lemma needed for verifying malnormality of the examples from Theorem 1.2.

3.1 | Finitely many non-pseudo-Anosovs

The goal of this subsection is to prove:

Proposition 3.1. *Let Σ be a finite-type surface and let $H < MCG(\Sigma)$ be a torsion-free malnormal convex-cocompact subgroup. Let $g \in MCG(\Sigma) - H$ have infinite order. Then the coset gH contains finitely many elements that are not pseudo-Anosov.*

The proof is postponed until after some preparatory lemmas. We will use the following lemma to check that specific elements are pseudo-Anosov. The lemma gives two alternative conditions, the first one is designed to show that the product of two elements one of which acts elliptically and has ‘large rotation angle’ is loxodromic. The second property, instead, captures a product where one element is loxodromic with sufficiently large translation length.

Lemma 3.2 (Loxodromic products). *For every $\delta, Q \geq 0$ there exists a constant $C_{Q,\delta}$ with the following property. Let A be a group acting on a δ -hyperbolic space X . Let $g_0, g_1 \in A$ be elements that, respectively, stabilise Q -quasiconvex sets $Y_0, Y_1 \subset X$. Let π_i be closest point projection to Y_i . Assume that one of the following holds:*

- (1) *let $y_i \in \pi_i(Y_{i+1}) \subseteq Y_i$ and suppose that $d(Y_0, Y_1) \geq C_{Q,\delta}$ and*

$$\text{diam}_X(N_{3\delta}([y_0, y_1]) \cap g_i[y_0, y_1]) \leq \frac{C_{Q,\delta}}{3}$$

for $i = 0, 1$; or

- (2) *$\text{diam}(\pi_1(g_0 Y_1)) \leq C_{Q,\delta}/3$ and $d(g_1 y, y) \geq C_{Q,\delta} + 2d(y, g_0 y)$ for some $y \in Y_1$.*

Then $g_0 g_1$ is loxodromic.

Proof. In case (1) we can form a bi-infinite quasigeodesic on which $g_0 g_1$ acts by translation by concatenating suitable translates of $U = [y_1, y_0]$, $V = [y_0, g_0 y_0]$, and $W = [y_1, g_1 y_1]$, with every other geodesic being a translate of $[y_0, y_1]$. Specifically, the $\langle g_0 g_1 \rangle$ -translates of the concatenation

$$U \cdot V \cdot (g_0 \bar{U}) \cdot (g_0 W),$$

which joins y_1 to $g_0 g_1 y_1$, concatenate to form a bi-infinite path, shown in Figure 3. Here the overline denotes the reverse path. In the notation of Lemma 2.3, by our assumption we have $d_{i,i\pm 2} \leq C_{Q,\delta}/3$, while the $d_{i,i\pm 1}$ are bounded independently of $C_{Q,\delta}$ since the $[y_0, y_1]$ are coarsely shortest geodesics connecting quasiconvex sets. So Lemma 2.3 implies $g_0 g_1$ is loxodromic provided $C_{Q,\delta}$ is sufficiently large in terms of Q and the constant C_δ from the lemma, which depends only on δ .

Case (2) is similar: we concatenate translates of $U = [y, g_1 y]$ and $V = [y, g_0 y]$. Specifically, the $\langle g_0 g_1 \rangle$ -translates of the concatenation $V \cdot g_0 U$ concatenate to form a bi-infinite path. Again, our assumptions allow us to apply Lemma 2.3. Indeed, successive translates of U have $d_{i\pm 2}$ bounded since intersections of neighbourhoods of the relevant translates of Y_1 have diameter bounded in terms of the diameter of $\pi_1(g_0 Y_1)$, so the assumption on $d(y, g_1 y)$, which controls $d_{i\pm 1}$, shows that Lemma 2.3 applies (with the translates of U as the even-indexed α_i) provided $C_{Q,\delta}$ is large enough in terms of Q and δ . \square

We will also need the following lemmas to check condition 1 in Lemma 3.2 when using it. Recall that for curve graphs we take $\delta = 100$.




FIGURE 3 Proof of Lemma 3.2(1). The paths concatenated to form the g_0g_1 -axis are coloured according to their $\langle g_0g_1 \rangle$ -orbits.

The following lemma can be formulated more generally for a group acting acylindrically on a hyperbolic space and an infinite-order element whose orbits are bounded, but we prefer to state just the special case we need to avoid discussing coarse fixed point sets.

Lemma 3.3. *Let Σ be a finite-type surface. Then there exists $D \geq 0$ with the following property. Let $g \in MCG(\Sigma)$ be an infinite-order reducible element, fixing the clique Y in $C(\Sigma)$. Then for any geodesic γ starting in Y we have that $N_{300}(\gamma) \cap g\gamma$ has diameter at most D .*

Proof. By hyperbolicity, for every $L \geq 0$ there exists $D \geq 0$ such that the following holds for g, γ as in the statement: if $p \in \gamma$ is moved distance at most L by a power of g , then γ has an initial subgeodesic of length $d(p, Y) - D$ that gets moved distance at most 300 by the same power of g . Therefore, if the lemma did not hold for a sufficiently large D , we would get a contradiction with acylindricity of the $MCG(\Sigma)$ -action on $C(\Sigma)$. □

Lemma 3.4. *Let the group A act acylindrically on the hyperbolic space X . Then there exists a constant D such that the following holds. Let $g \in A$ be loxodromic and let Y be its quasi-axis as in Proposition 2.1(1). Then for any $x \in X$ and geodesic $\gamma = [x, \pi_Y(x)]$ we have that $N_{3\delta}(\gamma) \cap g\gamma$ has diameter at most D .*

Proof. This is similar to Lemma 3.3, except we ‘shorten’ γ on both sides. □

We are ready for:

Proof of Proposition 3.1. Let Σ, H, g be as in the statement and fix an H -orbit $H \cdot o$ in $C(S)$. Let Q be the constant from Proposition 2.1. For each non-trivial $h \in H$, let Y_h be the Q -quasi-convex quasi-axis for h , from Proposition 2.1(1) (any non-trivial $h \in H$ is pseudo-Anosov). Let $Y_g \subset C(\Sigma)$ be the following Q -quasi-convex subspace:

- if g is reducible, then Y_g is a clique stabilised by g ;
- if g is pseudo-Anosov, then Y_g is the quasi-axis from Proposition 2.1(1).

Let D be the maximum of the constants from Lemmas 3.3 and 3.4. Let C_0 be such that $Y_h \subset N_{C_0}(H \cdot o)$ for all $h \in H$ (which exists since Y_h and $H \cdot o$ are uniformly quasiconvex and Y_h lies in some neighbourhood of $H \cdot o$). By enlarging Q once, we can assume $d(y', gy') \leq Q$ for all $y' \in Y_g$.

We will need the following claim:

Claim 3.5. For any $L \geq 0$, the set $Y_g \cap N_L(H \cdot o)$ is bounded. Moreover, there exists $C_1 \geq 0$ such that $\pi_{Y_h}(gY_h)$ has diameter at most C_1 for all $h \in H$.

Proof. If $Y_g \cap N_L(H \cdot o)$ had unbounded diameter then $gH \cdot o$ would have unbounded intersection with the $2L + Q$ -neighbourhood of $H \cdot o$, since $Q \geq d(y', gy')$ for all $y' \in Y_g$. Lemma 2.2 then gives that $gHg^{-1} \cap H$ is infinite, contradicting malnormality of H .

Similarly, malnormality of H and Lemma 2.2 imply that for all $L \geq 0$ there exists $B(L)$ such that $Y_h \cap N_L(gY_h)$ has diameter at most $B(L)$ (since $Y_h \subset N_{C_0}(H \cdot o)$ and $gY_h \subset N_{C_0}(gH \cdot o)$ and the lemma bounds the diameter of coarse intersections of the two coset orbits). It is well known that quasiconvex sets in a hyperbolic space that have bounded intersection of neighbourhoods also have bounded projection onto each other; we sketch this argument in our notation.

Consider two points $a, b \in gY_h$ and $a' \in \pi_{Y_h}(a), b' \in \pi_{Y_h}(b)$. Any point on $[a', b']$ which is further away than $500 + 2Q$ from $\{a', b'\}$ cannot be 200-close to $[a, a']$ or $[b, b']$, otherwise a' would not be nearly closest to a (that is, $a' \notin \pi_{Y_h}(a)$) or b' would not be nearly closest to b . Hence, any such point is within 200 of $[a, b]$ by thinness of quadrilaterals. Hence such point is within $200 + Q$ from gY_h , as well as within Q of Y_h . Hence, the diameter of $Y_h \cap N_L(gY_h)$ for $L = 200 + 2Q$ is at least $d(a', b') - 10^3 - 10Q$. Since a', b' were arbitrary, the diameter of $\pi_{Y_h}(gY_h)$ is bounded in terms of $B(L)$. \square

Finally, let $C_{Q,100}$ be the constant from Lemma 3.2. We can assume that $C_{Q,100} \geq 10D + 10^3 + 3C_1$.

Fix $h \in H$. First suppose that $d(Y_g, Y_h) > C_{Q,100}$. Then Lemma 3.2(1) applies in view of Lemma 3.4 (if g is pseudo-Anosov), or Lemma 3.3 (if g is infinite-order reducible). Hence Lemma 3.2(1) shows that gh is pseudo-Anosov.

Next, suppose that the translation length of h satisfies $\tau_h > 2Q + 10C_{Q,100}$ and that $d(Y_g, Y_h) \leq C_{Q,100}$. Recall that $\text{diam}(\pi_{Y_h}(gY_h)) \leq C_1 \leq C_{Q,100}/3$. Choose $y \in Y_h$ within distance $C_{Q,100}$ of some $y' \in Y_g$. Hence

$$2d(y, gy) + C_{Q,100} \leq 2d(y', gy') + 5C_{Q,100} < \tau_h \leq d(y, hy).$$

So we can apply Lemma 3.2(2) (with $g_1 = h$ and $g_0 = g$) and see that gh is pseudo-Anosov.

To conclude, it suffices to prove the following:

Claim 3.6. There exist finitely many h such that $d(Y_g, Y_h) \leq C_{Q,100}$ and $\tau_h \leq 2Q + 10C_{Q,100}$.

Proof. By Claim 3.5 (which says that Y_g can only come close to $H \cdot o$ at a bounded set) and since Y_h lies within C_0 of $H \cdot o$, we can only have $d(Y_g, Y_h) \leq C_{Q,100}$ if $d(Y_h, o) \leq C'$ for some C' (depending on g also).

Since H is quasi-isometrically embedded in $C(\Sigma)$, it suffices to show that in (a fixed Cayley graph of) a hyperbolic group G , for any $B \geq 0$ there are only finitely many $h \in G$ with translation distance bounded by B and such that Y_h contains a point within B of the identity (note that quiaxes in G get mapped within bounded distance of quiaxes in $C(\Sigma)$, and translation lengths get distorted in

a way controlled by the quasi-isometry constants). This is true since any such h moves the identity a uniformly bounded amount. □

We have shown that for all $h \in H$, either gh is pseudo-Anosov or h is one of the finitely many elements of H from Claim 3.6, which proves the proposition. □

3.2 | Pseudo-Anosovs without hidden symmetries

Lemma 3.7. *Let Σ be a finite-type hyperbolic surface with genus g and p punctures, with $(g, p) \notin \{(1, 1), (1, 2), (2, 0)\}$. Then there exists a pseudo-Anosov $\phi \in MCG(\Sigma)$ and $n_0 \geq 0$ such that the following holds. For any curve c on Σ and $n \geq n_0$ we have $Stab(c) \cap Stab(\phi^n(c)) = \{1\}$.*

Proof. Proposition 2.1(2) yields, for a pseudo-Anosov g , a unique maximal elementary subgroup $E(g)$ containing g and an invariant quasigeodesic γ_g in the curve graph $C(\Sigma)$, with the following property. For every $D \geq 0$ there exists $L \geq 0$ such that if $h \in G$ has the property that there exist $x, y \in \gamma_g$ with $d_{C(\Sigma)}(x, y) \geq L$ and $d_{C(\Sigma)}(x, h(x)), d_{C(\Sigma)}(y, h(y)) \leq D$, then $h \in E(g)$.

Let ϕ be a pseudo-Anosov for which $E(\phi) = \langle \phi \rangle$. Such a ϕ exists by [11, Theorem 6.14], since the maximal finite normal subgroup of a mapping class group coincides with its centre (see the proof of [22, Proposition 11.15]) and the centre is trivial under our assumptions on (g, p) (see [13, Section 3.4]).

Given any curve c and any sufficiently large n , if $h \in Stab(c) \cap Stab(\phi^n(c))$, then h coarsely stabilises the closest point projections of c and $\phi^n(c)$ to γ_ϕ , so that $h \in E(\phi) = \langle \phi \rangle$ by the above. No non-trivial power of ϕ fixes a curve, so $h = 1$. □

4 | COMMON WITNESSES FOR NON-INNER AUTOMORPHISMS

Proposition 4.1. *Let G be the fundamental group of a closed connected orientable surface of finite genus, and let ϕ_1, \dots, ϕ_n be non-inner automorphisms of G . Then there exists $x \in G$ such that for all i , the elements $\phi_i(x)$ and x are not conjugate.*

We need the following version of [26, Lemma 4.4] (which we use in the proof).

Lemma 4.2. *Let G be a torsion-free hyperbolic group. Let $l \geq 2$ and let $\{g_1, \dots, g_l\}, \{h_1, \dots, h_l\}$ be sets of non-commensurable, primitive, infinite-order elements. Assume that if $g_1^{m_1} \dots g_l^{m_l}$ is the identity of G for some m_i then all m_i are 0.*

There exists $N \in \mathbb{N}$ such that the following holds. Suppose that

$$g = g_1^{n_1} g_2^{n_2} \dots g_l^{n_l}$$

is conjugate to

$$h = h_1^{n_1} \dots h_l^{n_l}$$

in G , and for all $1 \leq i \leq l$ we have $n_i \geq N$. Then there exists $x \in G$ and $k \in \mathbb{N}$ with $h_i = xg_{i+k}x^{-1}$ for all i (indices mod l).




FIGURE 4 Edges in the bottom axis α_h are labelled by the h_i , with a translate of the piece $\{1, h_i, \dots, h_i^5\}$ shown in the middle. This has a long subpath D -close to a subpath of the piece $\{1, g_{\sigma(i)}, \dots, g_{\sigma(i)}^5\}$ of the translate of α_g shown at the top. (Large dots represent concatenation points of successive pieces.)

Proof. We conflate G with a fixed Cayley graph.

There exists a constant C_0 such that for all sufficiently large N we have that g stabilises a (C_0, C_0) -quasigeodesic α_g obtained by concatenating translates of the discrete paths $\{1, g_i, g_i^2, \dots, g_i^{n_i}\}$, with $n_i \geq N$, that we call *pieces*, and similarly for h . This is because the $\langle g_i \rangle$ are quasigeodesic lines, and moreover any two translates of two distinct such quasigeodesic lines cannot stay within bounded distance of each other for arbitrarily long time in view of Proposition 2.1(2) (g_i being not commensurable to g_j is equivalent to $g_i \notin E(g_j)$).

Let $D \geq 0$ be such that any two (C_0, C_0) -quasigeodesics in G with the same endpoints at infinity lie within distance D of each other.

If g and h are conjugate, we have that two translates of α_g and α_h have the same endpoints at infinity. Hence, for all M the following holds for all large N : each piece for h has a subpath of length at least M that has a translate that lies within D of a piece for α_g . See Figure 4.

For M large enough, this implies that each $h_i \in x_i E(g_{\sigma(i)}) x_i^{-1}$ for some $\sigma(i) \in \{1, \dots, l\}$ and some $x_i \in G$ by Proposition 2.1(2). Note that $i \mapsto \sigma(i)$ is injective since $\{h_1, \dots, h_l\}$ are pairwise non-commensurable.

To sum up, either there is no bijection σ and elements x_i such that h_i lies in $x_i E(g_{\sigma(i)}) x_i^{-1}$, in which case g and h cannot be conjugate for any large N and the conclusion holds vacuously, or there exist σ and x_i , in which case we can fix them and they depend only on the choice of g_i and h_i .

We now fix σ and x_i as above. We further let F be the (finite) collection of all elements of the form $x_i^{-1} x_{i+1}$ (indices mod l). Therefore, h (whence g) is conjugate to $(h'_1)^{n_1} f_1 \dots (h'_l)^{n_l} f_l$, for some $h'_i \in E(g_{\sigma(i)})$, where $f_i = x_i^{-1} x_{i+1}$ (indices mod l). We have now reduced to the setup of [26, Lemma 4.4], from which we can conclude that σ is a cyclic shift, say $\sigma(i) = i + k$ for some fixed k , and $f_i \in E(g_{\sigma(i)}) E(g_{\sigma(i+1)})$.

Now, we have $E(g_j) = \langle g_j \rangle$ for all j by the primitivity hypothesis and the assumption that G is torsion free, so $f_i = g_{\sigma(i)}^{l_i} g_{\sigma(i+1)}^{m_i}$ for some l_i, m_i . Up to multiplying each x_i on the right by $g_{\sigma(i)}^{l_i}$ (which does not affect the equation $h_i = x_i h'_i x_i^{-1}$), we then get $x_{i+1} = x_i g_{\sigma(i+1)}^{m'_i}$, where $m'_i = m_i + l_{i+1}$. This holds with indices mod l , so that $x_1 = x_1 g_{\sigma(2)}^{m'_1} \dots g_{\sigma(l)}^{m'_{l-1}} g_{\sigma(1)}^{m'_l}$. By our hypothesis on the g_i (applied to a cyclic conjugate) we get that all m_i are 0, and therefore we get that all x_i are equal, giving us the required x . \square

With the lemma in hand, we proceed to:

Proof of Proposition 4.1. Fix some $\phi = \phi_i$, and let g_1, \dots, g_l be the standard generators of the surface group G . Observe that the condition from Lemma 4.2 about products of the form $g_1^{m_1} \dots g_l^{m_l}$ holds

in this setup. Indeed, the g_i are generators in a $C'(1/6)$ small cancellation presentation where the (only) relator R , as a cyclic word, has the following property. Any subword of a cyclic permutation of R which has length more than half the length of R is not a subword of a cyclic permutation of a product $g_1^{m_1} \dots g_l^{m_l}$. We can conclude in view of Greendlinger's lemma.

Let $h_j = \phi(g_j)$. Consider the elements $g = g_1^T g_2^{T+1} \dots g_l^{T+l-1}$ and $h = h_1^T h_2^{T+1} \dots h_l^{T+l-1}$, for T larger than the N from Lemma 4.2. We claim that g and h cannot be conjugate, which implies that we can take $x = g$ for some sufficiently large T which works simultaneously for all ϕ_i .

If g and h were conjugate, then we would have $h_i = x g_{i+k} x^{-1}$ for some fixed $x \in G$ and natural number $k < l$. We claim that $k = 0$. This holds since there is a homomorphism ψ to \mathbb{Z} such that $\psi(g_1) = 1$ and $\psi(g_j) = 0$ for all $j \neq 1$, so that $\psi(g) = T$ and $\psi(h) = T + k$, so $k = 0$. Hence $\phi(g_i)$ acts as conjugation by x on all generators, implying that it is inner, contradicting the hypotheses. \square

Now we can prove the separability criterion.

Proof of Theorem 1.1. Throughout the proof we think of the mapping class group as an index 2 subgroup of $\text{Out}(\pi_1(\Sigma_g))$. Let $\Psi : \text{Aut}(\pi_1(\Sigma_g)) \rightarrow \text{Out}(\pi_1(\Sigma_g))$ be as in Section 2.5.

Since $MCG(\Sigma_g)$ is residually finite [15] and has finitely many conjugacy classes of finite-order elements [10], there is a finite-index torsion-free subgroup $M_0 \leq MCG(\Sigma_g)$.

Claim 4.3. Suppose that $\phi \in \text{Aut}(\pi_1(\Sigma_g))$ satisfies $\bar{\phi} = \Psi(\phi) \in M_0 - (H \cap M_0)$. Then there is a finite group F and a quotient $q : \text{Out}(\pi_1(\Sigma_g)) \rightarrow F$ such that $q(\bar{\phi}) \notin q(H)$.

Proof of Claim 4.3. We will use Proposition 2.5. First, by hypothesis, $\Gamma_H = \Psi^{-1}(H)$ is conjugacy separable, so condition (2) of Proposition 2.5 is satisfied. So, we are left to verify condition (1) of Proposition 2.5 for the given ϕ .

Since $\bar{\phi} \in M_0 - (H \cap M_0)$, the element $\bar{\phi}$ has infinite order in $MCG(\Sigma_g)$, so Proposition 3.1 applies. The only properties of ϕ used in the following argument are that $\bar{\phi} \in MCG(\Sigma)$, and $\bar{\phi}$ has infinite order, and $\bar{\phi} \notin H$.

We have to find $x \in \pi_1(\Sigma_g)$ such that $\phi(x) \neq h(x)$ for $h \in \Gamma_H$. This is equivalent to $\phi^{-1}h(x) \neq x$.

If $\Psi(\phi^{-1}h)$ (which is necessarily in $MCG(\Sigma_g)$ since $\Psi(h) \in H$ and $\bar{\phi} \in MCG(\Sigma_g)$) is pseudo-Anosov, then any non-trivial x satisfies the required condition for the h in question. Hence, we are concerned only with the non-pseudo-Anosov elements of $\bar{\phi}^{-1}H$, of which there are only finitely many by Proposition 3.1. Taking arbitrary preimages of these elements, we obtain finitely many $h_i \in \phi^{-1}\Gamma_H$ such that if $x \in G$ has the property that $h_i(x)$ is not conjugate to x for all i , then x satisfies the required property, that is, $\phi(x) \neq h(x)$ for all $h \in \Gamma_H$.

The existence of such x is guaranteed by Proposition 4.1. Hence Proposition 2.5 supplies a finite quotient $q : \text{Out}(\pi_1(\Sigma_g)) \rightarrow F$ such that $q(\bar{\phi}) \notin q(H)$. \square

From Claim 4.3, we get that $H \cap M_0$ is separable in M_0 , since for each $\phi \in M_0 - H \cap M_0$, we can restrict the quotient q from the claim to $M_0 \leq \text{Out}(\pi_1(\Sigma_g))$ to get a finite quotient $q : M_0 \rightarrow \bar{M}_0 \leq F$ where $q(\phi) \notin q(H \cap M_0)$. In summary, we have a finite-index subgroup $M_0 \leq MCG(\Sigma_g)$ and a subgroup $H \leq MCG(\Sigma_g)$ such that $H \cap M_0$ is separable in M_0 . Applying [28, Proposition 2.2(ii)] (taking $K = K_0 = \{1\}$ in the statement of that proposition) shows that H is separable in $MCG(\Sigma_g)$, as required. \square

5 | MMS SUBGROUPS

We finally construct examples to which Theorem 1.1 applies. We will use the following form of the Bounded Geodesic Image Theorem. We denote the subsurface projection to the subsurface Y by π_Y and abbreviate $d_{C(Y)}(\pi_Y(\cdot), \pi_Y(\cdot)) = d_{C(Y)}(\cdot, \cdot)$.

Theorem 5.1 [25, Theorem 3.1]. *For each $g \geq 2$ there exists $B \geq 0$ such that the following holds. If the vertices $x, y \in C(\Sigma)$ satisfy $d_{C(\Sigma_g - v)}(x, y) \geq B$ for some non-separating curve v , then any geodesic from x to y contains v .*

We now prove Theorem 1.2:

Theorem 5.2. *For every $g \geq 3$ and $n \geq 1$ there exist infinitely many conjugacy classes of malnormal convex-cocompact subgroups H of $MCG(\Sigma_g)$ isomorphic to a free group of rank n whose corresponding extension Γ_H is virtually compact special.*

Proof. In [23] the authors construct subgroups (that we call *MMS subgroups*) satisfying all requirements except possibly malnormality. MMS subgroups are described in [23, Proposition 5.7, Theorem 5.8]. Our goal is to construct (many) malnormal MMS subgroups.

Each MMS subgroup H stabilises a *tight tree of homologous non-separating curves* (a *tight tree* for short), which is a tree T with the following description (see [23, Section 3.1]):

- For each vertex v of T we have a vertex $c_v \in C(\Sigma_g)$, and for each edge e of T , we have a tight geodesic γ_e in $C(\Sigma_g)$ connecting the vertices corresponding to the endpoints of e .
- All multicurves occurring along γ_e (including the endpoints) are formed from non-separating curves. Moreover, all curves appearing along all tight geodesics are homologous to each other.

There are some additional properties of T guaranteeing that the natural subdivision of T isometrically embeds into $C(\Sigma_g)$ [23, Proposition 3.2]. Moreover, the tight tree T stabilised by an MMS subgroup H has the property that whenever u, v, w are consecutive vertices along a tight geodesic as above, then there is a pseudo-Anosov on $\Sigma_g - \{v\}$ which maps u to w . In fact, the tree is constructed starting from a choice of such partial pseudo-Anosovs and any choice of sufficiently large powers of the partial pseudo-Anosovs [23, Section 5.1].

As a result, there are MMS subgroups H that stabilise a tight tree T with the following additional properties. For an edge e of the tree, we call the *projection numbers* the quantities $d_{C(\Sigma_g - v)}(u, w)$ for consecutive u, v, w along γ_e . Then

- for any edge e of T , all projection numbers are different and larger than $10B$, for B as in Theorem 5.1;
- for any two edges e, f of T belonging to different H -orbits, all projection numbers for e are different from all projection numbers for f ;
- the tight geodesics all have length at least 10^3 ;
- $Stab_{MCG(\Sigma_g - v)}(u) \cap Stab_{MCG(\Sigma_g - v)}(w)$ is trivial for all u, v, w that are consecutive in γ_e , where e is any edge of T .

The last item can be arranged using the pseudo-Anosovs constructed in Lemma 3.7.

We now verify malnormality of H . Suppose for some $g \in MCG(\Sigma_g)$ that $gHg^{-1} \cap H$ is infinite. Then T and gT , respectively, contain bi-infinite geodesics α and $g\alpha'$ that 200-fellow-travel.

Fix an edge e of T such that $\gamma_e \subset \alpha$. We claim that for all vertices $c \in \gamma_e$ that are at distance more than 200 from both endpoints of γ_e , the geodesic $g\alpha'$ passes through c . Note that then the third item provides at least 100 such vertices c .

To prove the claim we use the lower bound on projection numbers from the first of the above four items and Theorem 5.1, as follows. First of all, by Theorem 5.1 the endpoints a', b' of γ_e each project to $C(\Sigma_g - c)$ within B of the projection of one of the vertices a, b adjacent to c along γ_e . Again by Theorem 5.1 points a'', b'' in $g\alpha'$ that lie within 200 of a', b' project within $2B$ of the projections of a, b . In particular, a'', b'' project B -far on $C(\Sigma_g - c)$, and once again by Theorem 5.1 we have that $g\alpha'$ passes through c .

With another application of the third item, there exists an edge f of T such that there are distinct triples u, v, w and u_1, v_1, w_1 of γ_e -consecutive vertices, both of which are contained in $g\gamma_f$.

By the second item, e and f are in the same H -orbit. Hence there exists $h \in H$ such that $u, v, w, u_1, v_1, w_1 \in \gamma_e \cap (gh)\gamma_e$.

By the 'projection numbers are different' clause of the first item, $ghu = u, ghv = v, ghw = w$. From the fourth item, it follows that gh is a power of the Dehn twist around v .

The same argument applied to u_1, v_1, w_1 shows that gh is also a power of the Dehn twist around v_1 , so $gh = 1$ and thus $g \in H$, that is, H is malnormal.

To arrange infinitely many conjugacy classes the argument is similar, arranging the projection numbers to be different for tight trees associated to different subgroups. \square

ACKNOWLEDGEMENTS

We are grateful to Michah Sageev for answering a question about MMS subgroups and to Ashot Minasyan for several very helpful comments and corrections. We also thank the referee for a number of suggestions that improved the paper.

JOURNAL INFORMATION

The *Bulletin of the London Mathematical Society* is wholly owned and managed by the London Mathematical Society, a not-for-profit Charity registered with the UK Charity Commission. All surplus income from its publishing programme is used to support mathematicians and mathematics research in the form of research grants, conference grants, prizes, initiatives for early career researchers and the promotion of mathematics.

REFERENCES

1. C. R. Abbott and J. F. Manning, *Acylically hyperbolic groups and their quasi-isometrically embedded subgroups*, arXiv:2105.02333, 2021.
2. I. Agol, *The virtual Haken conjecture*, Doc. Math. **18** (2013), 1045–1087. With an appendix by Agol, Daniel Groves, and Jason Manning.
3. Y. Antolín, A. Minasyan, and A. Sisto, *Commensurating endomorphisms of acylindrically hyperbolic groups and applications*, Groups Geom. Dyn. **10** (2016), no. 4, 1149–1210.
4. Y. Antolín, M. Mj, A. Sisto, and S. J. Taylor, *Intersection properties of stable subgroups and bounded cohomology*, Indiana Univ. Math. J. **68** (2019), no. 1, 179–199.
5. T. Aougab, *Uniform hyperbolicity of the graphs of curves*, Geom. Topol. **17** (2013), no. 5, 2855–2875.
6. J. Behrstock, M. Hagen, A. Martin, and A. Sisto, *A combinatorial take on hierarchical hyperbolicity and applications to quotients of mapping class groups*, arXiv:2005.00567, 2020.
7. M. Bestvina and K. Fujiwara, *Bounded cohomology of subgroups of mapping class groups*, Geom. Topol. **6** (2002), 69–89.
8. B. H. Bowditch, *Tight geodesics in the curve complex*, Invent. Math. **171** (2008), no. 2, 281–300.
9. B. Bowditch, *Uniform hyperbolicity of the curve graphs*, Pacific J. Math. Ind. **269** (2014), no. 2, 269–280.

10. M. R. Bridson, *Finiteness properties for subgroups of $GL(n, \mathbb{Z})$* , Math. Ann. **317** (2000), no. 4, 629–633.
11. F. Dahmani, V. Guirardel, and D. Osin, *Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces*, Mem. Amer. Math. Soc. **245** (2017), no. 1156, v+152.
12. G. Dufour, *Cubulations de variétés hyperboliques compactes*, Ph.D. thesis, Université Paris Sud-Paris XI, 2012.
13. B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, vol. 49, Princeton University Press, Princeton, NJ, 2012.
14. B. Farb and L. Mosher, *Convex cocompact subgroups of mapping class groups*, Geom. Topol. **6** (2002), 91–152.
15. E. K. Grossman, *On the residual finiteness of certain mapping class groups*, J. Lond. Math. Soc. (2) **9** (1974/75), 160–164.
16. M. F. Hagen and D. T. Wise, *Cubulating hyperbolic free-by-cyclic groups: the irreducible case*, Duke Math. J. **165** (2016), no. 9, 1753–1813.
17. U. Hamenstädt, *Word hyperbolic extensions of surface groups*, arxiv:0505244, 2005.
18. S. Hensel, P. Przytycki, and R. C. H. Webb, *1-Slim triangles and uniform hyperbolicity for arc graphs and curve graphs*, J. Eur. Math. Soc. (JEMS) **17** (2015), no. 4, 755–762.
19. A. E. Kent and C. J. Leininger, *Shadows of mapping class groups: capturing convex cocompactness*, Geom. Funct. Anal. **18** (2008), no. 4, 1270–1325.
20. C. J. Leininger and D. B. McReynolds, *Separable subgroups of mapping class groups*, Topology Appl. **154** (2007), no. 1, 1–10.
21. J. Maher and A. Sisto, *Random subgroups of acylindrically hyperbolic groups and hyperbolic embeddings*, Int. Math. Res. Not. IMRN (2019), no. 13, 3941–3980.
22. J. Maher and G. Tiozzo, *Random walks, WPD actions, and the Cremona group*, Proc. Lond. Math. Soc. (3) **123** (2021), no. 2, 153–202.
23. J. F. Manning, M. Mj, and M. Sageev, *Cubulating surface-by-free groups*, arXiv:1908.03545, 2019.
24. H. A. Masur and Y. N. Minsky, *Geometry of the complex of curves. I. Hyperbolicity*, Invent. Math. **138** (1999), no. 1, 103–149.
25. H. A. Masur and Y. N. Minsky, *Geometry of the complex of curves. II. Hierarchical structure*, Geom. Funct. Anal. **10** (2000), no. 4, 902–974.
26. A. Minasyan and D. Osin, *Normal automorphisms of relatively hyperbolic groups*, Trans. Amer. Math. Soc. **362** (2010), no. 11, 6079–6103.
27. A. Minasyan and P. Zalesskii, *Virtually compact special hyperbolic groups are conjugacy separable*, Comment. Math. Helv. **91** (2016), no. 4, 609–627.
28. G. A. Niblo, *Separability properties of free groups and surface groups*, J. Pure Appl. Algebra **78** (1992), no. 1, 77–84.
29. P. Przytycki and A. Sisto, *A note on acylindrical hyperbolicity of mapping class groups*, Hyperbolic geometry and geometric group theory, Advanced Studies Pure Mathematics, vol. 73, Math. Soc. Japan, Tokyo, 2017, pp. 255–264.
30. A. W. Reid, *Surface subgroups of mapping class groups*, Problems on mapping class groups and related topics, Proceedings of Symposia in Pure Mathematics, vol. 74, Amer. Math. Soc., Providence, RI, 2006, pp. 257–268.
31. P. Scott, *Subgroups of surface groups are almost geometric*, J. Lond. Math. Soc. **2** (1978), no. 3, 555–565.
32. D. T. Wise, *The structure of groups with a quasiconvex hierarchy*, Annals of Mathematics Studies, vol. 209, Princeton University Press, Princeton, NJ, 2021.